Targeting the GPIbα binding site of thrombin to simultaneously induce dual anticoagulant...

Search within citing articles

Exosite crosstalk in thrombin

JC Fredenburgh, JI Weitz - Journal of Thrombosis and Haemostasis, 2025 - Elsevier

Thrombin is the central mediator of hemostasis, where it converts fibrinogen to fibrin, activates upstream factors to promote coagulation, activates factor XIII and thrombin …

Save Cite Cited by 11 Related articles All 4 versions

Designing smaller, synthetic, functional mimetics of sulfated glycosaminoglycans as allosteric modulators of coagulation factors

EI Abdelfadiel, R Gunta, BK Villuri… - Journal of Medicinal …, 2023 - pmc.ncbi.nlm.nih.gov

Natural glycosaminoglycans (GAGs) are arguably the most diverse collection of natural products. Unfortunately, this bounty of structures remains untapped. Decades of research …

Save Cite Cited by 29 Related articles All 7 versions

The choice of targets and ligands for site-specific delivery of nanomedicine to atherosclerosis

A Zia, Y Wu, T Nguyen, X Wang, K Peter… - Cardiovascular …, 2020 - academic.oup.com

As nanotechnologies advance into clinical medicine, novel methods for applying nanomedicine to cardiovascular diseases are emerging. Extensive research has been …

Save Cite Cited by 64 Related articles All 13 versions

Allosterism-based simultaneous, dual anticoagulant and antiplatelet action: allosteric inhibitor targeting the glycoprotein Ibα-binding and heparin-binding site of …

AY Mehta, BM Mohammed, EJ Martin… - … of Thrombosis and …, 2016 - Wiley Online Library

Essentials Allosteric inhibition is a promising method for reducing bleeding risk associated with anticoagulants. Sulfated β-O4 lignin (SbO4L) targets exosite 2 of thrombin to induce …

Save Cite Cited by 56 Related articles All 10 versions

Allosteric partial inhibition of monomeric proteases. Sulfated coumarins induce regulation, not just inhibition, of thrombin

S Verespy III, AY Mehta, D Afosah, RA Al-Horani… - Scientific Reports, 2016 - nature.com

Allosteric partial inhibition of soluble, monomeric proteases can offer major regulatory advantages, but remains a concept on paper to date; although it has been routinely …

Save Cite Cited by 47 Related articles All 8 versions

Sulfated non-saccharide glycosaminoglycan mimetics as novel drug discovery platform for various pathologies

DK Afosah, RA Al-Horani - Current medicinal chemistry, 2020 - benthamdirect.com

Glycosaminoglycans (GAGs) are very complex, natural anionic polysaccharides. They are polymers of repeating disaccharide units of uronic acid and hexosamine residues. Owing to …

Save Cite Cited by 25 Related articles All 7 versions

Potent trivalent inhibitors of thrombin through hybridization of salivary sulfopeptides from hematophagous arthropods

SM Agten, EE Watson, J Ripoll-Rozada… - Angewandte Chemie …, 2021 - Wiley Online Library

Blood feeding arthropods, such as leeches, ticks, flies and mosquitoes, provide a privileged source of peptidic anticoagulant molecules. These primarily operate through inhibition of the …

Save Cite Cited by 20 Related articles All 12 versions

Molecular dynamics simulations of human α-thrombin in different structural contexts: evidence for an aptamer-guided cooperation between the two exosites

R Troisi, N Balasco, L Vitagliano… - Journal of Biomolecular …, 2021 - Taylor & Francis

Human α-thrombin (thrombin) is a multifunctional enzyme that plays a pivotal role in the coagulation pathway. Thrombin activity can be effectively modulated by G-quadruplex …

Save Cite Cited by 20 Related articles All 11 versions

A molecular dynamics-based algorithm for evaluating the glycosaminoglycan mimicking potential of synthetic, homogenous, sulfated small molecules

B Nagarajan, NV Sankaranarayanan, BB Patel… - PLoS …, 2017 - journals.plos.org

Glycosaminoglycans (GAGs) are key natural biopolymers that exhibit a range of biological functions including growth and differentiation. Despite this multiplicity of function, natural …

Save Cite Cited by 29 Related articles All 13 versions

Exosite 2-directed ligands attenuate protein C activation by the thrombin–thrombomodulin complex

K Chen, AR Stafford, C Wu, CH Yeh, PY Kim… - Biochemistry, 2017 - ACS Publications

Thrombin activity, inhibition, and localization are regulated by two exosites that flank the active site. Substrates, cofactors, and inhibitors bind to exosite 1 to promote active site …

Save Cite Cited by 22 Related articles All 3 versions