Allosteric partial inhibition of monomeric proteases. Sulfated coumarins induce regulation,...

Search within citing articles

Molecular mapping of α-thrombin (αT)/β2-glycoprotein I (β2GpI) interaction reveals how β2GpI affects αT functions

L Acquasaliente, D Peterle, S Tescari… - Biochemical …, 2016 - portlandpress.com

β2-Glycoprotein I (β2GpI) is the major autoantigen in the antiphospholipid syndrome, a thrombotic autoimmune disease. Nonetheless, the physiological role of β2GpI is still unclear …

☆ Save 99 Cite Cited by 30 Related articles All 7 versions

Long-term MALT1 inhibition in adult mice without severe systemic autoimmunity

A Demeyer, Y Driege, I Skordos, J Coudenys… - Iscience, 2020 - cell.com

The protease MALT1 is a key regulator of NF-κB signaling and a novel therapeutic target in autoimmunity and cancer. Initial enthusiasm supported by preclinical results with MALT1 …

☆ Save 99 Cite Cited by 18 Related articles All 9 versions

The research progress of direct thrombin inhibitors

ZG Sun, JM Zhang, SC Cui, ZG Zhang… - Mini Reviews in …, 2020 - benthamdirect.com

Blood coagulation is the process of changing the blood from the flowing state to the gel state. It is an important part of the hemostatic function. Coagulation is a process by which a …

☆ Save 99 Cite Cited by 18 Related articles All 5 versions

Fully Synthetic Non-Carbohydrate Heparin Mimetics—Perspectives for Therapeutic Anticoagulation and Beyond?

K Nekipelov, V Ferro, G Bendas - Archiv der Pharmazie, 2026 - Wiley Online Library

Heparin is one of the oldest drugs on the market. Although clinically well established as an anticoagulant for decades, owing to its natural origin and heterogeneous …

☆ Save 99 Cite Related articles All 7 versions

Discovery of novel N-acylpyrazoles as potent and selective thrombin inhibitors

KM Short, MA Estiarte, SM Pham, DC Williams… - European Journal of …, 2023 - Elsevier

Direct oral anticoagulants (DOACs), which includes thrombin and factor Xa inhibitors, have emerged as the preferred therapeutics for thrombotic disorders, penetrating a market …

☆ Save 99 Cite Cited by 8 Related articles All 4 versions

A small group of sulfated benzofurans induces steady-state submaximal inhibition of thrombin

DK Afosah, S Verespy III, RA Al-Horani… - Bioorganic & Medicinal …, 2018 - Elsevier

Despite the development of promising direct oral anticoagulants, which are all orthosteric inhibitors, a sizable number of patients suffer from bleeding complications. We have …

☆ Save 99 Cite Cited by 20 Related articles All 7 versions

Allosteric inhibition of factor XIIIa. Non-saccharide glycosaminoglycan mimetics, but not glycosaminoglycans, exhibit promising inhibition profile

RA Al-Horani, R Karuturi, M Lee, DK Afosah, UR Desai - PloS one, 2016 - journals.plos.org

Factor XIIIa (FXIIIa) is a transglutaminase that catalyzes the last step in the coagulation process. Orthostery is the only approach that has been exploited to design FXIIIa inhibitors …

☆ Save 99 Cite Cited by 25 Related articles All 14 versions »

Regulation of thrombin activity with ultrasmall nanoparticles: effects of surface chemistry

AL Lira, RS Ferreira, MLV Oliva, AA Sousa - Langmuir, 2020 - ACS Publications

Nanomaterials displaying well-tailored sizes and surface chemistries can provide novel ways with which to modulate the structure and function of enzymes. Recently, we showed …

☆ Save 99 Cite Cited by 16 Related articles All 7 versions

On the process of discovering leads that target the heparin-binding site of neutrophil elastase in the sputum of cystic fibrosis patients

S Morla, NV Sankaranarayanan… - Journal of medicinal …, 2019 - pmc.ncbi.nlm.nih.gov

Cystic fibrosis (CF) is a disease of dysregulated salt and fluid homeostasis that results in massive accumulation of neutrophil elastase resulting in lung degradation and death. The …

☆ Save 99 Cite Cited by 20 Related articles All 7 versions

Discovery of chromen-7-yl furan-2-carboxylate as a potent and selective factor XIa inhibitor

AJ Obaidullah, RA Al-Horani - Cardiovascular & Hematological …, 2017 - ingentaconnect.com

Background: Direct inhibition of coagulation factor XIa (FXIa) carries a significant promise for developing effective and safe anticoagulants. Method: In this report, we studied 4-methyl-2 …

☆ Save 99 Cite Cited by 21 Related articles All 5 versions