Chemoenzymatically prepared heparan sulfate containing rare 2-O-sulfonated glucuronic acid residues

Search within citing articles

Estimating glycosaminoglycan–protein interaction affinity: Water dominates the specific antithrombin–heparin interaction

A Sarkar, W Yu, UR Desai, AD MacKerell… - …, 2016 - academic.oup.com

Glycosaminoglycan (GAG)–protein interactions modulate many important biological processes. Structure–function studies on GAGs may reveal probes and drugs, but their …

Save Cite Cited by 26 Related articles All 12 versions

Combinatorial virtual library screening analysis of antithrombin binding oligosaccharide motif generation by heparan sulfate 3-*O*-Sulfotransferase 1

NV Sankaranarayanan, Y Bi, B Kuberan, UR Desai - 2020 - spj.science.org

Pharmaceutical heparin's activity arises from a key high affinity and high selectivity antithrombin binding motif, which forms the basis for its use as an anticoagulant. The current …

Save Cite Cited by 18 Related articles All 8 versions

Continuous-flow accelerated sulfation of heparan sulfate intermediates

S Anand, S Mardhekar, R Raigawali, N Mohanta… - Organic …, 2020 - ACS Publications

We report for the first time a continuous-flow strategy to execute O-sulfation modification of heparan sulfate (HS) oligosaccharides. A systematic investigation of the influence of the flow …

Save Cite Cited by 15 Related articles All 6 versions

Fluorous-tag assisted synthesis of a glycosaminoglycan mimetic tetrasaccharide as a high-affinity FGF-2 and midkine ligand

S Maza, N Gandia-Aguado, JL de Paz… - Bioorganic & medicinal …, 2018 - Elsevier

Here, we present the preparation of a sulfated, fully protected tetrasaccharide derivative following the glycosaminoglycan (GAG)-related sequence GlcNAc-β (1→ 4)-Glc-β (1→ 3) …

Save Cite Cited by 20 Related articles All 6 versions

In-depth molecular dynamics study of all possible chondroitin sulfate disaccharides reveals key insight into structural heterogeneity and dynamism

B Nagarajan, NV Sankaranarayanan, UR Desai - Biomolecules, 2022 - mdpi.com

GAGs exhibit a high level of conformational and configurational diversity, which remains untapped in terms of the recognition and modulation of proteins. Although GAGs are …

Save Cite Cited by 9 Related articles All 9 versions

Thermodynamic Affinity and Nature of Forces Defining Glycosaminoglycan-Protein Systems Using Fluorescence Spectroscopy

RS Boothello, U Desai - Glycosaminoglycans: Methods and Protocols, 2021 - Springer

Among the biophysical techniques used to study glycosaminoglycan (GAG)–protein interactions, fluorescence spectroscopy is a quantitative tool that has been extensively used …

Save Cite Related articles All 4 versions

ORIGINAL RESEARCH 2-O, 3-O Desulfated Heparin Blocks High Mobility Group Box 1 Release by Inhibition of p300 Acetyltransferase Activity

S Zheng - academia.edu

High mobility group box 1 (HMGB1) is an alarmin released from macrophages after infection or inflammation and is a biomarker of lung disease progression in patients with cystic …

Save Cite Related articles