Designing allosteric regulators of thrombin. Exosite 2 features multiple subsites that can...

Search within citing articles

An Overview of Thrombin Inhibitors in the Perspective of Structureactivity Relationships

J Wang, X Sun, N Li, R Sheng… - Current Medicinal …, 2023 - benthamdirect.com

Thrombosis is one of the most important pathogenic factors related to cardiovascular diseases. Presently, thrombin inhibitors have gradually gained prominence in clinical …

☆ Save 99 Cite Cited by 4 Related articles All 4 versions

Studies on fragment-based design of allosteric inhibitors of human factor XIa

RS Boothello, NV Sankaranarayanan… - Bioorganic & medicinal …, 2020 - Elsevier

Human factor XIa (hFXIa) has emerged as an attractive target for development of new anticoagulants that promise higher level of safety. Different strategies have been adopted so …

☆ Save 99 Cite Cited by 9 Related articles All 6 versions

Inhibition of thrombin by functionalized C60 nanoparticles revealed via in vitro assays and in silico studies

Y Liu, J Fu, W Pan, Q Xue, X Liu, A Zhang - Journal of Environmental …, 2018 - Elsevier

The studies on the human toxicity of nanoparticles (NPs) are far behind the rapid development of engineered functionalized NPs. Fullerene has been widely used as drug …

☆ Save 99 Cite Cited by 8 Related articles All 9 versions

A glassy carbon electrode modified with in-situ generated chromium-loaded CdS nanoprobes and heparin for ultrasensitive electrochemiluminescent determination of …

Q Wen, P Lu, P Yang - Microchimica Acta, 2016 - Springer

We describe the in-situ electrochemical synthesis of a chromium-loaded CdS nanoprobe (CdS-Cr) for use in ultrasensitive electrochemiluminescence (ECL) detection of thrombin. A …

☆ Save 99 Cite Cited by 9 Related articles All 8 versions

Substantial non-electrostatic forces are needed to induce allosteric disruption of thrombin's active site through exosite 2

AY Mehta, UR Desai - Biochemical and biophysical research …, 2014 - Elsevier

Sulfated β-O4 lignin (SbO4L), a non-saccharide glycosaminoglycan mimetic, was recently disclosed as a novel exosite 2-directed thrombin inhibitor with the capability of mimicking …

☆ Save 99 Cite Cited by 9 Related articles All 10 versions

Competition between the Hiyama and Suzuki–Miyaura Pd-catalyzed cross-coupling reaction mechanisms for the formation of some regioselective derivatives of …

Z Ahmadvand, M Bayat - Journal of Molecular Liquids, 2021 - Elsevier

Pd-NHC type complexes are supposed to have a high degree of participation in 'C–C'cross-coupling reactions, so they are commonly used in industry as well as academia. As regards …

☆ Save 99 Cite Cited by 3 Related articles

One-pot synthesis protocol for highly functionalized benzofuran scaffold

PS Sidhu - Postdoc J, 2013 - scholar.archive.org

Benzofuran scaffold appears frequently in natural products, commercial libraries, biologically and pharmaceutically relevant compounds. The presence of substituent handles …

☆ Save 99 Cite Cited by 5 Related articles »

A simple, general approach of allosteric coagulation enzyme inhibition through monosulfated hydrophobic scaffolds

PS Sidhu, Q Zhou, UR Desai - Bioorganic & Medicinal Chemistry Letters, 2014 - Elsevier

Allosteric inhibition of coagulation enzymes offers the advantage of controlled inhibition. In this study, a small library of mono sulfated indole and benzothiazole based molecules was …

☆ Save 99 Cite Cited by 4 Related articles All 5 versions

Dual-target synergistic antithrombotic mechanism of a dabigatran etexilate analogue (HY023016)

S Wan, S Cui, M Jiang, Q Wu, Y Ji… - Clinical and …, 2022 - Wiley Online Library

Thrombin has long been considered a desirable antithrombotic target, but anti-thrombin therapy without anti-platelet therapy has never achieved the ideal effect. HY023016, derived …

☆ Save 99 Cite Cited by 1 Related articles All 6 versions

Thiocarbamate-Linked Polysulfonate–Peptide Conjugates As Selective Hepatocyte Growth Factor Receptor Binders

S Besret, J Vicogne, F Dahmani, V Fafeur… - Bioconjugate …, 2014 - pmc.ncbi.nlm.nih.gov

The capacity of many proteins to interact with natural or synthetic polyanions has been exploited for modulating their biological action. However, the polydispersity of these …

☆ Save 99 Cite Cited by 2 Related articles All 6 versions