Emerging patterns of tyrosine sulfation and O-glycosylation cross-talk and co-localization

Search within citing articles

Global view of human protein glycosylation pathways and functions

KT Schjoldager, Y Narimatsu, HJ Joshi… - … reviews Molecular cell …, 2020 - nature.com

Glycosylation is the most abundant and diverse form of post-translational modification of proteins that is common to all eukaryotic cells. Enzymatic glycosylation of proteins involves a …

☆ Save 99 Cite Cited by 1601 Related articles All 7 versions

Siglec ligands

A Gonzalez-Gil, RL Schnaar - Cells, 2021 - mdpi.com

A dense and diverse array of glycans on glycoproteins and glycolipids decorate all cell surfaces. In vertebrates, many of these carry sialic acid, in a variety of linkages and glycan …

☆ Save 99 Cite Cited by 145 Related articles All 11 versions »

Direct mapping of tyrosine sulfation states in native peptides by nanopore

H Niu, MY Li, Y Gao, JG Li, J Jiang, YL Ying… - Nature chemical …, 2025 - nature.com

Sulfation is considered the most prevalent post-translational modification (PTM) on tyrosine; however, its importance is frequently undervalued due to difficulties in direct and …

☆ Save 99 Cite Cited by 58 Related articles All 5 versions

Genetic glycoengineering in mammalian cells

Y Narimatsu, C Büll, YH Chen, HH Wandall… - Journal of Biological …, 2021 - jbc.org

Advances in nuclease-based gene-editing technologies have enabled precise, stable, and systematic genetic engineering of glycosylation capacities in mammalian cells, opening up a …

☆ Save 99 Cite Cited by 133 Related articles All 6 versions

Short Mucin-Type *O*-Glycans in Cancer: Biomarker and Therapeutic Potential amid Analytical Barriers

H Sousa, IAB Ribeiro, M Falcão, DF Barreira… - …, 2025 - academic.oup.com

Mucin-type O-glycans are abundant protein modifications that regulate cell signalling, adhesion, and immune interactions. In cancer, their biosynthetic pathways are frequently …

☆ Save 99 Cite Cited by 8 Related articles All 3 versions

G protein-coupled receptors in the sweet spot: glycosylation and other post-translational modifications

CK Goth, UE Petäjä-Repo… - ACS pharmacology & …, 2020 - ACS Publications

Post-translational modifications (PTMs) are a fundamental phenomenon across all classes of life and several hundred different types have been identified. PTMs contribute widely to …

☆ Save 99 Cite Cited by 102 Related articles All 9 versions

O-glycan recognition and function in mice and human cancers

GE Cervoni, JJ Cheng, KA Stackhouse… - Biochemical …, 2020 - portlandpress.com

Protein glycosylation represents a nearly ubiquitous post-translational modification, and altered glycosylation can result in clinically significant pathological consequences. Here we …

☆ Save 99 Cite Cited by 89 Related articles All 8 versions

Total Synthesis of a PSGL-1 Glycopeptide Analogue for Targeted Inhibition of P-Selectin

B Dhakal, A Mandhapati, P Eradi, S Park… - Journal of the …, 2024 - ACS Publications

The high affinity interaction between P-selectin glycoprotein ligand-1 (PSGL-1) and P-selectin is mediated by a multimotif glycosulfopeptide (GSP) recognition domain consisting …

☆ Save 99 Cite Cited by 16 Related articles All 4 versions

Sulfotyrosine residues: interaction specificity determinants for extracellular protein–protein interactions

V Stewart, PC Ronald - Journal of Biological Chemistry, 2022 - jbc.org

Tyrosine sulfation, a post-translational modification, can determine and often enhance protein–protein interaction specificity. Sulfotyrosyl residues (sTyrs) are formed by the …

☆ Save 99 Cite Cited by 40 Related articles All 8 versions

Discovery and characterization of vertebrate sialoglycan-binding proteins

L Adler, S Yehuda, A Varki, V Padler-Karavani - Seminars in immunology, 2025 - Elsevier

All known life forms feature a dense ensemble of cell surface and extracellular glycans. In vertebrates this includes a high density of sialic acids (Sias), a family of acidic sugars …

☆ Save 99 Cite Cited by 3 Related articles All 4 versions