Increased galectin-9 levels correlate with disease activity in patients with DMARD-naïve...

Search within citing articles

Plasma proteomic signatures of social support and their association with cardiovascular disease and mortality: exploratory analyses in a national cohort study

P Qin, J Gong, A Steptoe, D Fancourt - GeroScience, 2026 - Springer

Social support has been related to cardiovascular disease (CVD) incidence and mortality in longitudinal cohort analyses, but the biological pathways underpinning this remain …

☆ Save 99 Cite Related articles All 5 versions

Plasma proteomic signatures of social support and their association with cardiovascular disease and mortality

P Qin, J Gong, A Steptoe, D Fancourt - medRxiv, 2025 - pmc.ncbi.nlm.nih.gov

Background: Social support has been related to cardiovascular disease (CVD) incidence and mortality in longitudinal cohort analyses, but the biological pathways underpinning this …

☆ Save 99 Cite Cited by 2 Related articles All 5 versions

Predictors of remission or combined remission and low disease activity in rheumatoid arthritis patients in Taiwan: A prospective cohort study

PH Tsai, YF Fang, YF Chen, CC Chen… - Journal of Clinical …, 2024 - mdpi.com

Objectives: This study aimed to identify predictors of remission or low disease activity (LDA) in patients with rheumatoid arthritis (RA) and low-ultrasound inflammation. Methods: A total …

☆ Save 99 Cite Cited by 4 Related articles All 8 versions »

Salivary Galectin-9 Levels in Primary Sjögren's Disease: An Observational Cross-Sectional Case–Control Study

E İnanç, S Yolbaş, S Zontul, F Albayram… - Journal of Clinical …, 2026 - mdpi.com

Background/Objectives: Primary Sjögren's disease is a systemic autoimmune disease characterized by chronic inflammation of exocrine glands and heterogeneous clinical …

☆ Save 99 Cite Related articles All 8 versions »

The Role of Serum MCP-1 in the Diagnosis and Assessment of Rheumatoid Arthritis

AM Mahdi, AH Odda, AG Yassin - Karbala Journal of …, 2025 - journals.uokerbala.edu.iq

Background: RA is a chronic autoimmune disease characterized by synovial inflammation and joint destruction. Monocyte chemoattractant protein-1 (MCP-1) plays a key role in …

☆ Save 99 Cite Cited by 1 Related articles All 4 versions »

Differential Expression of Galectin-1 and Galectin-9 in Immune-Mediated Inflammatory Diseases

J Galicia Carreón, M Pardines Ortiz, N Montes… - 2025 - scientiasalut.gencat.cat

Galectin-1 and-9 (Gal1/9) are essential mediators of immune-inflammatory responses, which makes these proteins potential biomarkers for immune-mediated diseases (IMIDs), such as …

☆ Save 99 Cite Related articles All 2 versions »

AB0734 PEPTIDYLARGININE DEIMINASE 4 (PAD4) INHIBITOR 4, 6-BIS (2, 2, 2-TRIFLUOROETHOXY)-1, 3, 5-TRIAZIN-2-AMINE II AMELIORATES COLLAGEN …

ZY Huang, WN Huang, CC Lin - Annals of the Rheumatic Diseases, 2024 - Elsevier

Background: Peptidylarginine deiminase-4 (PAD4) is a calcium-dependent enzyme that catalyzes the conversion of arginine into citrulline in macromolecules within the body …

☆ Save 99 Cite Related articles All 3 versions

AB0735 ALTERED GALECTIN EXPRESSION CORRELATE WITH TNF-MEDIATED INFLAMMATORY CARTILAGE DAMAGE

S Hayer, B Niederreiter, C Strauß… - Annals of the …, 2024 - ard.eular.org

Background: Galectins (Gal) are a family of soluble ß-galactoside-binding proteins that serve as regulators in multiple physiological processes like proliferation or cell adhesion. There is …

☆ Save 99 Cite Related articles All 3 versions