Interaction of thrombin with sucrose octasulfate

Search within citing articles

Potent, selective, allosteric inhibition of human plasmin by sulfated non-saccharide glycosaminoglycan mimetics

DK Afosah, RA Al-Horani… - Journal of medicinal …, 2017 - pmc.ncbi.nlm.nih.gov

Although plasmin inhibitors could be used in multiple disorders, their use has been restricted to preventing blood loss in hemostatic dysregulation because of poor efficacy and …

☆ Save 99 Cite Cited by 39 Related articles All 4 versions

Soluble expression, purification, and characterization of active recombinant human tissue plasminogen activator by auto-induction in *E. coli*

X Long, Y Gou, M Luo, S Zhang, H Zhang, L Bai… - BMC …, 2015 - Springer

Background Human tissue plasminogen activator (tPA) belongs to the serine protease family. It converts plasminogen into plasmin and is used clinically to treat thrombosis …

☆ Save 99 Cite Cited by 41 Related articles All 18 versions

Designing allosteric regulators of thrombin. Monosulfated benzofuran dimers selectively interact with Arg173 of exosite 2 to induce inhibition

MH Abdel Aziz, PS Sidhu, A Liang, JY Kim… - Journal of medicinal …, 2012 - ACS Publications

Earlier, we reported on the design of sulfated benzofuran dimers (SBDs) as allosteric inhibitors of thrombin (Sidhu et al. J. Med. Chem. 2011 54 5522–5531). To identify the site of …

☆ Save 99 Cite Cited by 51 Related articles All 6 versions

Raman spectroscopy of disulfide bridges in thrombin

NN Brandt, AY Chikishev, AV Golovin… - Biomedical …, 2014 - journals.sagepub.com

Disulfide bridges of bovine α-thrombin are studied using the analysis of the Raman spectral features in the spectral interval 500–550 cm− 1. The changes under study are caused by the …

☆ Save 99 Cite Cited by 28 Related articles All 5 versions

Crystal structure of a protein–aromatic foldamer composite: macromolecular chiral resolution

JM Alex, V Corvaglia, X Hu, S Engilberge… - Chemical …, 2019 - pubs.rsc.org

Co-crystallization of a 2 kDa tether-free sulfonated foldamer and the 13 kDa lysine-rich cytochrome c yielded a remarkable biohybrid assembly with chiral resolution of the foldamer …

☆ Save 99 Cite Cited by 20 Related articles All 10 versions

A fluorometric assay of thrombin using magnetic nanoparticles and enzyme-free hybridization chain reaction

Z Huang, D He, HW Li - Microchimica Acta, 2020 - Springer

A fluorescence method based on functionalized magnetic nanoparticles (FMNPs) and hybridization chain reaction (HCR) is developed for the enzyme-free amplified determination …

☆ Save 99 Cite Cited by 19 Related articles All 7 versions

Identification of the site of binding of sulfated, low molecular weight lignins on thrombin

MHA Aziz, PD Mosier, UR Desai - Biochemical and biophysical research …, 2011 - Elsevier

Sulfated, low molecular weight lignins (LMWLs), designed recently as macromolecular mimetics of the low molecular weight heparins (LMWHs), were found to exhibit a novel …

☆ Save 99 Cite Cited by 32 Related articles All 9 versions

Plasmin regulation through allosteric, sulfated, small molecules

RA Al-Horani, R Karuturi, DT White, UR Desai - Molecules, 2015 - mdpi.com

Plasmin, a key serine protease, plays a major role in clot lysis and extracellular matrix remodeling. Heparin, a natural polydisperse sulfated glycosaminoglycan, is known to …

☆ Save 99 Cite Cited by 28 Related articles All 16 versions »

Heparinized Dendritic Mesoporous Silica Nanoparticles for the Selective Adsorption and Isolation of Thrombin

M Qiao, QQ Sun, XQ Wang, MM Liu, YX Sun… - ACS …, 2025 - pmc.ncbi.nlm.nih.gov

A novel solid-phase extractant, heparinized dendritic mesoporous silica nanoparticles (DMSNs-NH-Hep), is prepared by grafting heparin onto amino-terminated dendritic …

☆ Save 99 Cite Related articles All 4 versions

Polysulfated trehalose as a novel anticoagulant agent with dual mode of action

Q Rashid, M Abid, N Gupta, T Tyagi… - BioMed Research …, 2015 - Wiley Online Library

Physiological hemostatic balance is a coordinated outcome of counteracting coagulation and fibrinolytic systems. An imbalance of procoagulant and anticoagulant factors may result …

☆ Save 99 Cite Cited by 17 Related articles All 15 versions