Interaction of thrombin with sucrose octasulfate

Search within citing articles

Proteoglycan sequence

L Li, M Ly, RJ Linhardt - Molecular BioSystems, 2012 - academic.oup.com

Proteoglycans (PGs) are among the most structurally complex biomacromolecules in nature. They are present in all animal cells and frequently exert their critical biological functions …

☆ Save 99 Cite Cited by 160 Related articles All 12 versions

Heparin mimetics with anticoagulant activity

AA Nahain, V Ignjatovic, P Monagle… - Medicinal Research …, 2018 - Wiley Online Library

Heparin, a sulfated polysaccharide belonging to the glycosaminoglycan family, has been widely used as an anticoagulant drug for decades and remains the most commonly used …

☆ Save 99 Cite Cited by 81 Related articles All 14 versions

Synthesis, biological evaluation and reversal of sulfonated di-and triblock copolymers as novel parenteral anticoagulants

J Swieton, J Miklosz, N Bielicka… - Advanced …, 2024 - Wiley Online Library

Despite targeting different coagulation cascade sites, all Food and Drug Administration-approved anticoagulants present an elevated risk of bleeding, including potentially life …

☆ Save 99 Cite Cited by 9 Related articles All 9 versions

Sulfated pentagalloylglucoside is a potent, allosteric, and selective inhibitor of factor XIa

RA Al-Horani, P Ponnusamy, AY Mehta… - Journal of medicinal …, 2013 - pmc.ncbi.nlm.nih.gov

Inhibition of factor XIa (FXIa) is a novel paradigm for developing anticoagulants without major bleeding consequences. We present the discovery of sulfated pentagalloylglucoside …

☆ Save 99 Cite Cited by 110 Related articles All 7 versions

On the specificity of heparin/heparan sulfate binding to proteins. Anion-binding sites on antithrombin and thrombin are fundamentally different

PD Mosier, C Krishnasamy, GE Kellogg, UR Desai - PLoS One, 2012 - journals.plos.org

Background The antithrombin–heparin/heparan sulfate (H/HS) and thrombin–H/HS interactions are recognized as prototypic specific and non-specific glycosaminoglycan …

☆ Save 99 Cite Cited by 74 Related articles All 19 versions »

Designing allosteric inhibitors of factor XIa. Lessons from the interactions of sulfated pentagalloylglucopyranosides

RA Al-Horani, UR Desai - Journal of Medicinal Chemistry, 2014 - pmc.ncbi.nlm.nih.gov

We recently introduced sulfated pentagalloylglucopyranoside (SPGG) as an allosteric inhibitor of factor XIa (FXIa)(Al-Horani et al.,). To better understand the SPGG–FXIa …

☆ Save 99 Cite Cited by 71 Related articles All 5 versions

Allosteric partial inhibition of monomeric proteases. Sulfated coumarins induce regulation, not just inhibition, of thrombin

S Verespy III, AY Mehta, D Afosah, RA Al-Horani… - Scientific Reports, 2016 - nature.com

Allosteric partial inhibition of soluble, monomeric proteases can offer major regulatory advantages, but remains a concept on paper to date; although it has been routinely …

☆ Save 99 Cite Cited by 47 Related articles All 8 versions

Raman study of the cleavage of disulphide bonds in albumin, chymotrypsin, and thrombin

NN Brandt, AY Chikishev, VN Kruzhilin - Vibrational Spectroscopy, 2017 - Elsevier

The structural changes resulting from the reduction of disulphide bonds in three globular proteins (albumin, chymotrypsin and thrombin) in aqueous solutions are investigated using …

☆ Save 99 Cite Cited by 36 Related articles All 6 versions

Allosteric inhibition of factor XIa. Sulfated non-saccharide glycosaminoglycan mimetics as promising anticoagulants

RA Al-Horani, D Gailani, UR Desai - Thrombosis research, 2015 - Elsevier

Recent development of sulfated non-saccharide glycosaminoglycan mimetics, especially sulfated pentagalloyl glucopyranoside (SPGG), as potent inhibitors of factor XIa (FXIa)(J …

☆ Save 99 Cite Cited by 53 Related articles All 9 versions

Highly sensitive chemiluminescence biosensor for protein detection based on the functionalized magnetic microparticles and the hybridization chain reaction

N Li, J Chen, M Luo, C Chen, X Ji, Z He - Biosensors and Bioelectronics, 2017 - Elsevier

An ultrasensitive chemiluminescence (CL) biosensor for the detection of protein is developed in this study based on the functionalized magnetic microparticles (MMPs) and the …

☆ Save 99 Cite Cited by 42 Related articles All 7 versions