Novel reversible fluorescent glycan linker for functional glycomics

Search within citing articles

Exploiting glycan arrays and biosensors to diagnose and understand diseases

AS Lee, HR Heo, YH Song, CS Kim, JH Seo… - Biotechnology …, 2025 - Elsevier

Aberrant glycosylation patterns, often observed in disease states compared to those in normal states, are involved in disease-related processes such as cancer cell development …

☆ Save 99 Cite Cited by 2 Related articles All 6 versions

The Role of Glycans and Glycosyltransferases Involved in N-glycan Branching in Cancer, COPD, Alzheimer's Disease, and Redox Regulation

N Taniguchi, Y Ohkawa, M Nakano, J Gu… - JoVE (Journal of …, 2026 - jove.com

Glycosyltransferases that biosynthesize glycans and their genes (glycogenes) play important roles in health and disease. In general, pathophysiological changes are defined …

☆ Save 99 Cite Cited by 2 Related articles All 2 versions

Integrated analysis of natural glycans using a versatile pyrazolone-type heterobifunctional tag ANPMP

Y Liu, T Yang, J Rong, J Yuan, L Man, M Wei… - Carbohydrate …, 2024 - Elsevier

Glycans mediate various biological processes through carbohydrate-protein interactions, and glycan microarrays have become indispensable tools for understanding these …

☆ Save 99 Cite Cited by 5 Related articles All 3 versions

A mild protecting-group free strategy for neoglycoconjugate synthesis

P Raju, C Dong, CR Garen, MT Woodside… - Bioconjugate …, 2025 - ACS Publications

The synthesis of neoglycoconjugates has paved the way for the discovery of novel probes that mimic natural glycoconjugates and can provide designed research tools and …

☆ Save 99 Cite Related articles All 7 versions

A universal strategy of glyconanoparticle preparation using a bifunctional linker for lectin sensing and cell imaging

Q Sha, J Fei, C Tu, BF Liu, Z Hu, X Liu - Microchimica Acta, 2022 - Springer

Abstract Glyconanoparticles (G-NPs), biofunctional nanomaterials that can fully combine the unique properties of nanoparticles (NPs) with the bioactivities of carbohydrates, have …

☆ Save 99 Cite Cited by 7 Related articles All 9 versions

A Versatile Urea Type Linker for Functionalizing Natural Glycans and Its Validation in Glycan Arrays

S Serna, R Artschwager… - … A European Journal, 2023 - Wiley Online Library

The isolation from organisms and readily available glycoproteins has become an increasingly convenient source of N-glycans for multiple applications including glycan …

☆ Save 99 Cite Cited by 3 Related articles All 6 versions

Regeneration of free reducing glycans from reductive amination-tagged glycans by oxone

Q Zhang, S Wang, Z Li, Y Lasanajak… - The Journal of …, 2022 - pmc.ncbi.nlm.nih.gov

Glycans are usually fluorescently tagged by reductive amination for analytic tools. However, free reducing glycan regeneration is sometimes important and necessary for further …

☆ Save 99 Cite Cited by 4 Related articles All 6 versions

Novel multifunctional glycan probes to elucidate specificities of diverse glycan recognition systems

J Yu, AC Decout, A Di Maio, W Chai, T Stehle… - bioRxiv, 2023 - biorxiv.org

Glycan microarrays of sequence-defined glycans are the most widely used approach for high-throughput studies of interactions of glycans with glycan binding proteins. Currently …

☆ Save 99 Cite Cited by 2 Related articles All 4 versions »

Polysaccharides of Fungal Origin: Focus on the Capsule of the Pathogen *Cryptococcus neoformans*

MP Wear, A Casadevall - Polysaccharides of Microbial Origin: Biomedical …, 2021 - Springer

Cryptococcus neoformans was first isolated as a human pathogen in the 1890s. Subsequent study has shown that the polysaccharide capsule of C. neoformans is a key virulence factor …

☆ Save 99 Cite Cited by 3 Related articles All 3 versions

Development of a multifunctional neoglycoside auxiliary for applications in glycomics research

T Cheewawisuttichai, M Brichacek - Organic & Biomolecular Chemistry, 2021 - pubs.rsc.org

A novel, multifunctional, tetrazine-containing neoglycoside auxiliary has been synthesized in three steps and 28% overall yield. The oxyamine was conjugated with unprotected …

☆ Save 99 Cite Cited by 3 Related articles All 5 versions