Rational design of potent, small, synthetic allosteric inhibitors of thrombin

Search within citing articles

De novo synthesis of benzofurans *via* trifluoroacetic acid catalyzed cyclization/oxidative aromatization cascade reaction of 2-hydroxy-1,4-diones

Q Sha, H Liu - Organic & Biomolecular Chemistry, 2019 - pubs.rsc.org

A facile, one-pot method for the de novo synthesis of benzofurans from 2-hydroxy-1, 4-diones is described. Using trifluoroacetic acid (TFA) as the catalyst and N-bromobutanimide …

☆ Save 99 Cite Cited by 17 Related articles All 6 versions

Novel heparin mimetics reveal cooperativity between exosite 2 and sodium-binding site of thrombin

MHA Aziz, UR Desai - Thrombosis research, 2018 - Elsevier

Introduction Thrombin is a primary target of most anticoagulants. Yet, thrombin's dual and opposing role in pro-as well as anti-coagulant processes imposes considerable challenges …

☆ Save 99 Cite Cited by 12 Related articles All 8 versions

Blood compatibility of heparin-inspired, lactose containing, polyureas depends on the chemistry of the polymer backbone

Y Huang, MA Shaw, MR Warmin, ES Mullins… - Polymer …, 2016 - pubs.rsc.org

The influence of polymer chemical structure and degree of sulfation of polyurea glycopolymers on blood compatibility is reported. Polymers were prepared using lactose …

☆ Save 99 Cite Cited by 17 Related articles All 4 versions

Synthesis and biological evaluation of some dibenzofuran-piperazine derivatives

L Yurttaş, U Abu Mohsen, Y Ozkan… - Journal of Enzyme …, 2016 - Taylor & Francis

In the present paper, a novel series of dibenzofuran-piperazine derivatives were synthesized via the treatment of N-(2-methoxy-3-dibenzofuranyl)-2-chloroacetamide with …

☆ Save 99 Cite Cited by 15 Related articles All 12 versions

Homogeneous, synthetic, non-saccharide glycosaminoglycan mimetics as potent inhibitors of human cathepsin G

DK Afosah, RM Fayyad, VR Puliafico, S Merrell… - Biomolecules, 2023 - mdpi.com

Cathepsin G (CatG) is a pro-inflammatory neutrophil serine protease that is important for host defense, and has been implicated in several inflammatory disorders. Hence, inhibition …

☆ Save 99 Cite Cited by 4 Related articles All 10 versions »

On scaffold hopping: challenges in the discovery of sulfated small molecules as mimetics of glycosaminoglycans

PS Sidhu, PD Mosier, Q Zhou, UR Desai - Bioorganic & medicinal …, 2013 - Elsevier

The design of sulfated, small, nonsaccharide molecules as modulators of proteins is still in its infancy as standard drug discovery tools such as library of diverse sulfated molecules and …

☆ Save 99 Cite Cited by 23 Related articles All 7 versions

Thrombin Exosite Maturation and Ligand Binding at ABE II Help Stabilize PAR Binding Competent Conformation at ABE I

R Billur, TM Sabo, MC Maurer - Biochemistry, 2019 - pmc.ncbi.nlm.nih.gov

Thrombin, derived from zymogen prothrombin (ProT), is a serine protease involved in procoagulation, anticoagulation, and platelet activation. Thrombin's actions are regulated …

☆ Save 99 Cite Cited by 10 Related articles All 7 versions

An Overview of Thrombin Inhibitors in the Perspective of Structureactivity Relationships

J Wang, X Sun, N Li, R Sheng… - Current Medicinal …, 2023 - benthamdirect.com

Thrombosis is one of the most important pathogenic factors related to cardiovascular diseases. Presently, thrombin inhibitors have gradually gained prominence in clinical …

☆ Save 99 Cite Cited by 4 Related articles All 4 versions

Inhibition of thrombin by functionalized C60 nanoparticles revealed via in vitro assays and in silico studies

Y Liu, J Fu, W Pan, Q Xue, X Liu, A Zhang - Journal of Environmental …, 2018 - Elsevier

The studies on the human toxicity of nanoparticles (NPs) are far behind the rapid development of engineered functionalized NPs. Fullerene has been widely used as drug …

☆ Save 99 Cite Cited by 8 Related articles All 9 versions

Deciphering conformational changes associated with the maturation of thrombin anion binding exosite I

R Billur, D Ban, TM Sabo, MC Maurer - Biochemistry, 2017 - pmc.ncbi.nlm.nih.gov

Thrombin participates in procoagulation, anticoagulation, and platelet activation. This enzyme contains anion binding exosites, ABE I and ABE II, which attract regulatory …

☆ Save 99 Cite Cited by 6 Related articles All 5 versions