Substantial non-electrostatic forces are needed to induce allosteric disruption of thrombin's...

Search within citing articles

Designing smaller, synthetic, functional mimetics of sulfated glycosaminoglycans as allosteric modulators of coagulation factors

El Abdelfadiel, R Gunta, BK Villuri… - Journal of Medicinal …, 2023 - pmc.ncbi.nlm.nih.gov

Natural glycosaminoglycans (GAGs) are arguably the most diverse collection of natural products. Unfortunately, this bounty of structures remains untapped. Decades of research …

☆ Save Cite Cited by 29 Related articles All 7 versions

Sulfated non-saccharide glycosaminoglycan mimetics as novel drug discovery platform for various pathologies

DK Afosah, RA Al-Horani - Current medicinal chemistry, 2020 - benthamdirect.com

Glycosaminoglycans (GAGs) are very complex, natural anionic polysaccharides. They are polymers of repeating disaccharide units of uronic acid and hexosamine residues. Owing to …

☆ Save Cite Cited by 25 Related articles All 7 versions

A unique nonsaccharide mimetic of heparin hexasaccharide inhibits colon cancer stem cells via p38 MAP kinase activation

RS Boothello, NJ Patel, C Sharon… - Molecular cancer …, 2019 - aacrjournals.org

Targeting of cancer stem cells (CSC) is expected to be a paradigm-shifting approach for the treatment of cancers. Cell surface proteoglycans bearing sulfated glycosaminoglycan (GAG) …

☆ Save Cite Cited by 45 Related articles All 8 versions

Allosteric inhibition of α-thrombin enzymatic activity with ultrasmall gold nanoparticles

AL Lira, RS Ferreira, RJS Torquato, MLV Oliva… - Nanoscale …, 2019 - pubs.rsc.org

The catalytic activity of enzymes can be regulated by interactions with synthetic nanoparticles (NPs) in a number of ways. To date, however, the potential use of NPs as …

☆ Save Cite Cited by 43 Related articles All 12 versions

Allosterism-based simultaneous, dual anticoagulant and antiplatelet action: allosteric inhibitor targeting the glycoprotein Ibα-binding and heparin-binding site of …

AY Mehta, BM Mohammed, EJ Martin… - … of Thrombosis and …, 2016 - Wiley Online Library

Essentials Allosteric inhibition is a promising method for reducing bleeding risk associated with anticoagulants. Sulfated β-O4 lignin (SbO4L) targets exosite 2 of thrombin to induce …

☆ Save Cite Cited by 56 Related articles All 10 versions »

Allosteric partial inhibition of monomeric proteases. Sulfated coumarins induce regulation, not just inhibition, of thrombin

S Verespy III, AY Mehta, D Afosah, RA Al-Horani… - Scientific Reports, 2016 - nature.com

Allosteric partial inhibition of soluble, monomeric proteases can offer major regulatory advantages, but remains a concept on paper to date; although it has been routinely …

☆ Save Cite Cited by 47 Related articles All 8 versions

Development of orally active thrombin inhibitors for the treatment of thrombotic disorder diseases

LW He, WC Dai, NG Li - Molecules, 2015 - pmc.ncbi.nlm.nih.gov

Thrombotic disorders represent the major share of the various cardiovascular diseases, and significant progress has been made in the development of synthetic thrombin inhibitors as …

☆ Save Cite Cited by 21 Related articles All 11 versions

Novel heparin mimetics reveal cooperativity between exosite 2 and sodium-binding site of thrombin

MHA Aziz, UR Desai - Thrombosis research, 2018 - Elsevier

Introduction Thrombin is a primary target of most anticoagulants. Yet, thrombin's dual and opposing role in pro-as well as anti-coagulant processes imposes considerable challenges …

☆ Save Cite Cited by 12 Related articles All 8 versions

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and …

SM Rudman, J Heavyside, DJ Rennison, D Schluter - boris-portal.unibe.ch

Effectiveness of vertical ridge augmentation interventions. A systematic review and meta‐analysis
Page 1 Accepted Article This article has been accepted for publication and undergone full …

☆ Save Cite Related articles All 11 versions »