Unique repertoire of anti-carbohydrate antibodies in individual human serum

Search within citing articles

Use of Microbial Microarrays to Define Antibody Specificity

CJ Stowell, V Ortiz, CL Cepeda, VC Lee… - Methods in Transfusion …, 2026 - Springer

The formation of alloantibodies against red blood cell (RBC) antigens is the most frequent immune barrier to blood transfusion. Among these antibodies, those that occur naturally …

☆ Save 99 Cite Related articles All 3 versions

Structural insight into terminal galactose recognition by two non-HBGA binding GI. 3 noroviruses

C Wang, H Kang, M Tan, J Cong, D Su, X Li… - Journal of …, 2022 - journals.asm.org

Human noroviruses (huNoVs) cause epidemic acute gastroenteritis using histo-blood group antigens (HBGAs) as host receptors or attachment factors to initiate an infection. While most …

☆ Save 99 Cite Cited by 6 Related articles All 4 versions

Prospects and challenges of the study of anti-glycan antibodies and microbiota for the monitoring of gastrointestinal cancer

EP Smorodin - International Journal of Molecular Sciences, 2021 - mdpi.com

Over the past decades, a large amount of data has been accumulated in various subfields of glycobiology. However, much clinically relevant data and many tools are still not widely used …

☆ Save 99 Cite Cited by 5 Related articles All 6 versions »

Characterization of human medullary thyroid carcinoma glycosphingolipids identifies potential cancer markers

K Säljö, A Thornell, C Jin, O Norlén… - International Journal of …, 2021 - mdpi.com

Medullary thyroid carcinoma (MTC) accounts for only 1–2% of thyroid cancers; however, metastatic MTC is a mortal disease with no cure. In this study, glycosphingolipids were …

☆ Save 99 Cite Cited by 5 Related articles All 9 versions »

Identification of genes for variable regions of immunoglobulins that recognize sialylated glycans

T Okuda, M Kitamara, S Kasahara, K Kato - Biochemical and biophysical …, 2021 - Elsevier

We previously reported an antibody (clone ID: FR9, IgM-κ) that recognizes the sialyl oligosaccharide Neu5Acα2, 6Galβ1, 4GlcNAc as an epitope on glycoproteins and …

☆ Save 99 Cite Cited by 5 Related articles All 7 versions

STRUCTURE-FUNCTION STUDIES OF AN RTX-ADHESIN FROM VIBRIO VULNIFICUS AND ITS ROLE IN BINDING HOST CELL SURFACES

BP Soares - 2026 - queensu.scholaris.ca

Numerous Gram-negative pathogenic bacteria employ repeat-in-toxin-adhesins to colonize a niche and initiate biofilm formation. Surface adhesion can occur via specific ligand-binding …

☆ Save 99 Cite Related articles All 2 versions

Measurement of low avidity, polyreactive immunoglobulin G antibodies with increased sensitivity by using low ionic strength buffers

A Engelmaier, H Arno Butterweck, A Weber - J Anal Tech Res, 2022 - academia.edu

Low avidity polyreactive immunoglobulin G (IgG) antibodies have a broad range of affinity with dissociation constants of 10-5 to 10-8. Both their low concentration and avidity require …

☆ Save 99 Cite Cited by 2 Related articles »

Profiling of Anti-glycan IgA and IgG in Oral Fluid Using a Glycan Microarray

NV Shilova, SM Polyakova, MV Lavrenteva… - Dynamics of Bacteria …, 2025 - Springer

Secretory immunoglobulins (sIg), primarily IgA, control the microflora of the mucous membrane, protecting it from pathogens, as well as promoting its colonization by …

☆ Save 99 Cite Related articles All 5 versions

The usefulness of rare blood group systems in the risk determination for severe COVID-19

TG Konstantinidis, V Iliadi, G Martinis, M Panopoulou - Pathophysiology, 2021 - mdpi.com

The newly identified human coronavirus was named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), based on a detailed analysis of clinical manifestation. It was …

☆ Save 99 Cite Cited by 2 Related articles All 10 versions »

A review of glycoconjugate vaccines: The current state of the technology, considerations for the future, and the gold standard carrier protein, CRM (197)

O Meeds - PQDT-Global, 2023 - search.proquest.com

Conjugation of bacterial capsular polysaccharides to immunogenic carrier proteins has been employed for almost five decades to protect infants and adults against bacterial …

☆ Save 99 Cite Related articles