



# Epidemiological trends of foodborne *Campylobacter* outbreaks in the United States of America, 1998–2016

Azam Ali Sher <sup>a, b</sup>  , Muhammad Adnan Ashraf <sup>c</sup> , Bahar E Mustafa <sup>d</sup> , Muhammad Mohsin Raza <sup>e</sup> 

Show more 

 Outline |  Share  Cite

<https://doi.org/10.1016/j.fm.2021.103751>

[Get rights and content](#)

## Highlights

- *Campylobacter* foodborne outbreaks were found to be increasing across the USA.
- Besides dairy products (32%), chicken items (17%) and fresh produce (6%) were commonly associated with *Campylobacter* outbreaks.
- Vegetable and chicken item associated outbreaks were more in summer while milk associated outbreaks were frequent in winter.
- Dine-out (39%) and private residence places (17%) were frequently associated with *Campylobacter* food outbreaks.
- The active disease surveillance should be enhanced by every single state to reduce the burden of *Campylobacter* infections.

Full Text

Help

## Abstract

*Campylobacter* is a major cause of foodborne diarrheal infections in the United States of America (USA). This study aimed to elucidate the patterns of *Campylobacter* foodborne outbreaks temporally and spatially concerning food vehicles. We collected the data of foodborne outbreaks from 1998 to 2016 reported to the Centers for Disease Control and Prevention. The incidence rate of outbreaks for each food source was calculated and analyzed for each variable including season, food location, and census region. Overall, 465 single-state outbreaks and 8003 cases were reported during 1998–2016. Outbreaks were frequently attributed to dairy products (32%), chicken (17%) and vegetables (6%). Binomial regression analysis showed that compared to chicken items, the highest rate ratio of outbreaks was associated with dairy products (1.86) followed by vegetables (1.35) and meat products (0.76). More outbreaks were reported in the summer (35%) followed by the spring (26%) and fall (22%) season. We found that the highest number of outbreaks occurred in the West 159 (34%) and Midwest 137 (29%) census regions. The study highlights the role of dairy, chicken, and vegetables as food vehicles in *Campylobacter* outbreaks. Findings from this study can help in devising strategies to mitigate the increasing occurrence of *Campylobacter* foodborne outbreaks.

[Previous](#)[Next](#)

## Keywords

Food sources; *Campylobacter* outbreaks; Foodborne disease; Disease incidence; Risk factors

---

### Recommended articles

[Full Text](#)[Help](#)

## Cited by (9)

[Survive and thrive: Control mechanisms that facilitate bacterial adaptation to survive manufacturing-related stress](#)

2022, International Journal of Food Microbiology

[Show abstract](#)

[Screening, identification, and application of nucleic acid aptamers applied in food safety biosensing](#)

2022, Trends in Food Science and Technology

[Show abstract](#) ✓

## [Improved removal performance of Gram-negative and Gram-positive bacteria in sand filtration system with arginine modified biochar amendment](#)

2022, Water Research

[Show abstract](#) ✓

## [Outbreaks of foodborne Salmonella enteritidis in the United States between 1990 and 2015: An analysis of epidemiological and spatial-temporal trends](#)

2021, International Journal of Infectious Diseases

[Show abstract](#) ✓

## [Surveillance for foodborne disease outbreaks in Zhejiang Province, China, 2015–2020](#)

2022, BMC Public Health

## [Draft Genome Sequence of Campylobacter jejuni ST-508 Strain Shizu21005, Isolated from an Asymptomatic Food Handler in Japan, 2021](#)

2022, Microbiology Resource Announcements



[View all citing articles on Scopus](#)

[View full text](#)

© 2021 Elsevier Ltd. All rights reserved.

[Full Text](#)

[Help](#)



Copyright © 2022 Elsevier B.V. or its licensors or contributors.  
ScienceDirect® is a registered trademark of Elsevier B.V.

RELX™