

# Fourier Analysis on Coset Spaces

Brian Forrest  
presented by Alan Kydd

April 2, 2004

## Introduction

$G$  a locally compact group,  $H$  a closed subgroup of  $G$ . We define and study natural analogs of the Fourier and Fourier-Stieltjes algebras for  $G/H$ , and show that when  $H$  is compact,  $A(G/H)$  can be used to study the nature of  $G/H$  in a manner similar to the group case.

## Table of Contents

1. Defining  $A(G/H)$  and  $B(G/H)$
2. Structure of  $A(G/K)$  for compact subgroup  $K$
3. Weak amenability of  $A(G)$

## Defining $A(G/H)$ and $B(G/H)$

### Notation

Let  $H$  be a closed subgroup of  $G$ .  $q : G \rightarrow G/H$  denotes the canonical quotient map.  $\tilde{x}$  denotes the left coset  $xH$ . We have an isomorphism between  $C(G/H)$  and  $C(G : H) = \{f \in C(G/H) : f(xh) = f(x) \forall x \in G, h \in H\}$  via the map  $\tilde{f} \mapsto f$ , where  $f = \tilde{f} \circ q$ . We denote the equivalence class of all continuous unitary representations of  $G$  by  $\Sigma_G$ . For  $\pi \in \Sigma_G$ , we let  $A_\pi$  denote the closed linear span of the coefficient functions of  $\pi$ , and we denote the weak-\* closure of  $A_\pi$  by  $B_\pi$ . For  $\rho$ , the left regular representation of  $G$  on  $L_2(G)$ ,  $A_\rho$  is usually denoted  $A(G)$ .

### Definition 1

$$B(G : H) = \{u \in B(G) : u(xh) = u(x) \forall x \in G, h \in H\},$$

$$A(G : H) = \{u \in B(G : H) : q(\text{supp } u) \text{ compact in } G/H\}^{-\|\cdot\|_{B(G)}}.$$

**Proposition 2**    (i)  $B(G : H), A(G : H)$  are closed subalgebras of  $B(G)$ . Moreover,  $A(G : H)$  is a closed ideal in  $B(G : H)$ .

(ii)  $B(G : H)$  is unital.

(iii)  $A(G : H) \cap A(G) \neq \{0\}$  iff  $H$  is compact.

(iv)  $A(G : H) = B(G : H)$  iff  $G/H$  is compact.

We know that  $A(G : H)$  is isometrically isomorphic to  $A(G/H)$  when  $H$  is compact and normal. We can do better:

**Proposition 3** *Let  $H$  be a closed normal subgroup of  $G$ . Then  $B(G : H)$  and  $A(G : H)$  are isometrically isomorphic to  $B(G/H)$  and  $A(G/H)$  respectively.*

**Theorem 4** *Let  $H$  be a closed subgroup of  $G$ . Then there exists a projection  $P : B(G) \rightarrow B(G : H)$  with  $\|P\| \leq 1$ .*

In general,  $P$  does not map  $A(G)$  onto  $A(G : H)$ . However, for compact subgroup  $K$ , we have

$$P(f)(x) = P_K(f)(x) := \int_K f(xk)dk.$$

Thus we have the following corollary:

**Corollary 5** *Let  $K$  be a compact subgroup of  $G$ . Then  $P_K$  is a continuous projection of  $B(G)$  onto  $B(G : K)$ . The restriction of  $P_K$  to  $A(G)$  is a projection of  $A(G)$  onto  $A(G : K)$ .*

The analog of the Fourier Algebra for the coset space  $G/H$  is usually considered to be the space  $A_{\pi_H}$ , where  $\pi_H$  is the quasi-regular representation of  $G$  determined by  $H$ . This definition has two major problems.

**Problem I.**  $A_{\pi_H}$  is in general not an algebra.

For example, when  $K$  is a compact subgroup,  $A_{\pi_H}$  is an algebra iff  $A_{\pi_H} = A_{\pi_{K_1}}$ , where  $K_1 = \cap_{x \in G} xKx^{-1}$ . Since  $K_1$  is normal,  $A_{\pi_{K_1}} = A(G/K_1)$  (Arsac, 1976).

We can say more:

**Proposition 6** *Let  $K$  be a compact subgroup of  $G$ . Then  $A(G : K) = A_{\pi}$  for some  $\pi \in \Sigma_G$  iff  $K$  is normal.*

**Problem II.** It is possible to have two distinct closed subgroups  $H_1, H_2$ , and yet  $A_{\pi_{H_1}} = A_{\pi_{H_2}}$ , even when these subgroups are compact (Arsac, 1976).

**Proposition 7** *Let  $K_1, K_2$  be compact subgroups of  $G$ . Then  $A(G : K_1) = A(G : K_2)$  iff  $K_1 = K_2$ . If  $G$  is a [SIN]-group and  $H_1, H_2$  are closed subgroups of  $G$  with  $H_1 \neq H_2$ , then  $A(G : H_1) \neq A(G : H_2)$ .*

**Corollary 8** *Let  $K_1, K_2$  be compact subgroups of  $G$ . Then  $B(G : K_1) = B(G : K_2)$  iff  $K_1 = K_2$ . If  $G$  is a [SIN]-group, then  $B(G : H_1) = B(G : H_2)$  for  $H_1, H_2$  closed subgroups of  $G$ , iff  $H_1 = H_2$ .*

*Proof:* in either case,  $f_1$  as constructed above is in  $B(G : K_2) [B(G : H_2)]$ , but not in  $B(G : K_1) [B(G : H_1)]$ .  $\square$

In light of the above problems for  $A_{\pi_H}$ ,  $A(G : H)$  and  $B(G : H)$  are more useful analogs for  $G/H$  of the Fourier and Fourier-Stieltjes algebras.

**Definition 9** *We define  $A(G/H)$ , the Fourier algebra of the coset space  $G/H$ , to be the subalgebra of  $C(G/H)$  identified with  $A(G : H)$ .*

**Definition 10** *We define  $B(G/H)$ , the Fourier-Stieltjes algebra of the coset space  $G/H$ , to be the subalgebra of  $C(G/H)$  identified with  $B(G : H)$ .*

When  $H$  is a compact subgroup,  $A(G/H)$  and  $B(G/H)$  have many of the same properties of  $A(G)$ ,  $B(G)$ .



**Definition 11**  $AP(G/H)$  is the set of all  $f \in C(G/H)$  such that the set  $\{xf : x \in G\}$  is relatively compact in the norm topology of  $C(G/H)$  (Skantharajah, 1985).

**Definition 12**  $WAP(G/H)$  is the set of all  $f \in C(G/H)$  such that the set  $\{xf : x \in G\}$  is relatively compact in the weak topology of  $C(G/H)$  (Skantharajah, 1985).

**Proposition 13** Let  $H$  be a closed subgroup of  $G$ . Then,

- (i)  $B(G/H) \subseteq WAP(G/H)$ , and  $B(G/H) \cap AP(G/H)$  is the space identified with  $B(G : H) \cap AP(G)$ .
- (ii)  $B(G/H) \cap AP(G/H)$  is a complemented subalgebra of  $B(G/H)$  with the Radon-Nikodym property.

Recall that  $A(G)$  is sup-norm dense in  $C_0(G)$ . For a compact subgroup  $K$ ,  $A(G/K)$  is also sup-norm dense in  $C_0(G/K)$ . However, when  $H$  is not compact, there may exist  $f \in C_0(G/H)$  such that  $f \notin WAP(G/H)$  (Chou). Thus  $A(G/H)$  may not separate the points of  $G/H$ . The proof of Proposition 7 gives the following result.

**Theorem 14** *Let  $G$  be a [SIN]-group with a closed subgroup  $H$ . Then  $A(G/H)$  separates points in  $G/H$ .*

This next proposition extends a result of Herz (1973).

**Proposition 15** *Let  $K$  be a compact subgroup of  $G$ . Let  $H$  be a closed subgroup of  $G$  such that  $K \subseteq H$ . Then every  $\tilde{u} \in A(H/K)$  extends to a function  $\tilde{u}_1 \in A(G/K)$  with  $\|\tilde{u}\|_{A(H/K)} = \|\tilde{u}_1\|_{A(G/K)}$ .*

We can extend a  $u \in B(H)$  to some  $v \in B(G)$  when  $G$  is a [SIN]-group or if  $H$  is normal (Cowling, Rodway, 1979).

**Proposition 16** *Let  $H$  be a closed subgroup of  $G$ . Assume that either  $G$  is a [SIN]-group or that  $H$  is normal. Let  $H_1$  be another closed subgroup containing  $H$ . Then every  $u \in B(H_1/H)$  extends to a  $v \in B(G/H)$  with the same norm.*

## Structure of $A(G/K)$ for compact subgroup $K$

### Notation

Let  $\mathcal{A}$  be a semisimple commutative Banach algebra.  $\Delta(\mathcal{A})$  denotes the maximal ideal space of  $\mathcal{A}$ . Given any closed set  $A$  of  $\Delta(\mathcal{A})$ , we define the following ideals:

$$I(A) = \{u \in \mathcal{A} : u(x) = 0 \forall x \in A\}$$

$$j(A) = \{u \in I(A) : \text{supp } u \text{ is compact}\}$$

$$J(A) = \text{the norm closure of } j(A) \text{ in } I(A).$$

$A$  is said to be of spectral sythesis if  $I(A) = J(A)$ .  $A$  is said to be of weak spectral sythesis if for each  $u \in I(A)$ , there exists a positive integer  $n$  such that  $u^n \in J(A)$ . We say that (weak) spectral sythesis fails if there exists a closed subset  $A$  of  $\Delta(\mathcal{A})$  that is not a set of (weak) spectral sythesis.

**Definition 17** *A multiplier of  $\mathcal{A}$  is a linear operator  $T$  on  $\mathcal{A}$  for which  $T(uv) = uT(v)$ . We denote the set of all such mps by  $\mathcal{M}(\mathcal{A})$ , a Banach space with the operator norm.*

**Definition 18** *Let  $\mathcal{X}$  be a Banach  $\mathcal{A}$ -bimodule. A derivation of  $\mathcal{A}$  on  $\mathcal{X}$  is a linear map  $D : \mathcal{A} \rightarrow \mathcal{X}$  such that  $D(uv) = uD(v) + D(u)v$  for every  $u, v \in \mathcal{A}$ .*

**Theorem 19**  $A(G/K)$  is a regular commutative Banach algebra with  $\Delta A(G/K) = G/K$ .

*Proof outline:* Let  $\tilde{x}_0 \in G/K$ .  $\delta_{\tilde{x}_0}(\tilde{u}) = \tilde{u}(\tilde{x}_0)$  is continuous multiplicative linear functional on  $A(G/K)$ .

Suppose  $\tilde{\Phi} \in \Delta(A(G/K))$ . We can identify  $\tilde{\Phi}$  with  $\Phi \in A(G : K)^*$ .  $A(G : K)$  is complemented in  $A(G)$  by Corollary 5. Thus there exists  $\Gamma \in VN(G)$  with  $P_K^*(\Phi) = \Gamma$  and  $\Gamma|_{A(G:K)} = \Phi$ .  $\Phi \neq 0$ , so  $\Gamma \neq 0$ . It can be shown that  $\text{supp}\Gamma = x_0K$  for some  $x_0 \in G$ .  $x_0K$  is a set of spectral synthesis for  $A(G)$  (Forrest, 1992), therefore  $\Gamma$  is the weak-\* norm limit of  $\Psi = \sum_{i=1}^n a_i L_{x_i}$ , where  $x_i \in x_0K$ . But  $L_{x_i}|_{A(G:K)} = \Phi = \delta_{\tilde{x}_0}$ . It can be shown that the map  $\tilde{x}_0 \mapsto \delta_{\tilde{x}_0}$  is a homeomorphism of  $G/K$  onto  $\Delta(A(G/K))$ .  $\square$

**Theorem 20** *Let  $G$  be a locally compact group with compact subgroup  $K$ . The following are equivalent:*

- (i)  $G$  is amenable*
- (ii)  $G/K$  is an amenable coset space*
- (iii)  $A(G/K)$  has a bounded approximate identity consisting of functions with compact support in  $G/K$*
- (iv)  $A(G/K)$  weakly factorizes*

**Corollary 21** *Let  $G$  be an amenable locally compact group with a compact subgroup  $K$ . Then  $\mathcal{M}(A(G/K)) = B(G/K)$  and the usual norms agree.*

**Proposition 22** *Let  $K$  be a compact subgroup of  $G$ . Let  $\tilde{E} \subset G/K$  be a set for which (weak) spectral synthesis fails in  $A(G/K)$ . Then (weak) spectral synthesis fails for  $q^{-1}(\tilde{E})$  in  $A(G/K)$ . In particular, if (weak) spectral synthesis fails for  $A(G/K)$ , then (weak) spectral synthesis fails for  $A(G)$ .*

**Corollary 23** *Let  $G$  be a locally compact group for which  $A(G)$  admits (weak) spectral synthesis. Then  $G$  is totally disconnected.*

**Corollary 24** *Let  $G$  be a locally compact group with a compact subgroup  $K$ . Then each singleton  $\{x\} \subset G/K$  is a set of spectral synthesis for  $A(G/K)$ . Furthermore, if  $G$  is amenable, then every finite subset of  $G/K$  is a set of spectral synthesis.*

**Proposition 25** *Let  $G$  be amenable with compact subgroup  $K$ . Let  $\{x_1, \dots, x_n\}$  be a finite subset of  $G/K$ . Then  $I = I_{G/K}\{x_1, \dots, x_n\}$  has a bounded approximate identity  $\{u_\alpha\}$  in  $A(G/K) \cap C_c(G/K)$ .*

**Theorem 26** *Let  $K$  be a compact subgroup of  $G$ . The following are equivalent:*

- (i)  $G$  is amenable
- (ii) If  $I$  is a cofinite ideal of  $A(G/K)$ , then  $I = I(\{x_1, \dots, x_n\})$  where  $n = \text{codim}(I)$
- (iii) Every cofinite ideal in  $A(G/K)$  has a bounded approximate identity
- (iv) Each homomorphism of  $A(G/K)$  with finite dimensional range is continuous



This theorem is a generalization of Forrest (1988)

**Theorem 27** *Let  $K$  be a compact subgroup of  $G$ . The following are equivalent:*

(i)  *$G$  is amenable*

(ii) *Every derivation from  $A(G/K)$  into a Banach  $A(G/K)$ -bimodule is continuous.*

## Weak amenability of $A(G)$

**Definition 28** *A commutative Banach algebra  $\mathcal{A}$  is weakly amenable if every continuous derivation from  $\mathcal{A}$  into a commutative Banach  $\mathcal{A}$ -bimodule is identically zero.*

$A(G)$  is weakly amenable if  $G$  is discrete (Forrest, 1988). If  $G$  is the rotation group on  $\mathbb{R}^3$ , then  $A(G)$  is not weakly amenable (Johnson, 1994). Very little is known about the class of groups  $G$  for which  $A(G)$  is amenable. We show that this class contains all totally disconnected groups.

**Theorem 29** *Let  $H$  be an open subgroup of  $G$ . Then  $A(G/H)$  is weakly amenable.*

*Proof:* Let  $D : A(G/H) \rightarrow \mathcal{X}$  be a continuous derivation into a commutative Banach  $A(G/H)$ -bimodule. Let  $\tilde{u}$  be an idempotent in  $A(G/H)$ . Then  $D(\tilde{u}) = D(\tilde{u}^n) = nD(\tilde{u})$  for  $n \geq 2$ . Thus  $D(\tilde{u}) = 0$ .  $H$  is open, so the linear span of the idempotents is dense in  $A(G/H)$ . Hence  $D$  is identically zero.  $\square$

**Lemma 30** *Let  $G$  be totally disconnected. Let  $u \in A(G)$  and  $\epsilon > 0$ . Then there exists an open compact subgroup  $K$  and a  $v \in A(G : K)$  such that  $\|u - v\|_{A(G)} < \epsilon$ .*

**Theorem 31** *Let  $G$  be disconnected. Then  $A(G)$  is weakly amenable.*

*Proof:* Let  $D : A(G) \rightarrow \mathcal{X}$  be a continuous derivation into a commutative Banach  $A(G)$ -bimodule. Let  $K$  be a compact open subgroup of  $G$ . The restriction of  $D$  to  $A(G : K)$  determines a derivation of  $A(G/K)$ . By Theorem 29,  $D$  is zero on each  $A(G : K)$ . By Lemma 30 each  $u \in A(G)$  can be approximated within  $\epsilon$  by some  $v \in A(G : K)$  for some open compact subgroup  $K$ . Thus  $D = 0$ .  $\square$

**Proposition 32** *Let  $G_1, G_2$  be such that  $A(G_i)$  is weakly amenable for  $i = 1, 2$ . Then  $A(G_1 \times G_2)$  is also weakly amenable.*

**Corollary 33** *Let  $G = G_1 \times G_2$  where  $G_1$  is abelian and  $G_2$  is totally disconnected. Then  $A(G)$  is weakly amenable.*