

How Dangerous Is Myanmar’s Dam Infrastructure?

A Hobby Seismology Project After the 2025 M7.7 Mandalay Earthquake

Aung Khant Zaw

aungkhantzawd@gmail.com

<https://github.com/akzedevops/mmeqopendata>

May 2026

Abstract

After the devastating March 2025 earthquake in Myanmar, I wanted to understand how many of the country’s 254 dams are actually at risk. This write-up documents a hobby project that pulls earthquake data from a public API, scores every dam for seismic risk, and runs probabilistic hazard analysis—all with open-source Python tools. The corrected Abrahamson & Silva (2008) ground-motion model, validated against the actual Naypyidaw recording (predicted $0.51g$ vs. observed $0.57g$), shows that about half the dams sit in zones of significant seismic hazard. Everything—code, data, interactive maps—is on GitHub.

Contents

1	Background	3
1.1	What This Project Does	3
1.2	Data Sources	3
2	The Earthquake Catalog	3
2.1	b-Value and Catalog Completeness	3
2.2	Spatial Clustering	5
3	Ground Motion: Getting the GMPE Right	5
3.1	Abrahamson & Silva (2008)	6
3.2	Validation Against Actual Recordings	6
3.3	Site-Specific Vs30	6
4	Dam Risk Scoring	6
4.1	Results	6
4.2	Sensitivity to Weights	10
5	Probabilistic Hazard	10
6	Coulomb Stress Transfer	10
7	Fragility and Uncertainty	13
7.1	Dam Fragility Curves	13
7.2	Monte Carlo PGA Uncertainty	13
8	Population Exposure	13

9 What I Learned	15
9.1 Limitations	15
10 How to Use the Code	15

1 Background

On March 28, 2025, a magnitude 7.7 earthquake hit central Myanmar along the Sagaing Fault. The rupture was enormous—475 km of surface break, supershear velocity, 5–7 m of slip. A seismic station in Naypyidaw recorded $0.57g$ of peak ground acceleration. Over 5,000 people died, and the government reported 586 dams damaged.

Myanmar has at least 254 documented dams (irrigation, hydropower, multi-purpose), many built before modern seismic codes existed. The Sagaing Fault runs right through the middle of the country where most of these dams are. I wanted to know: which dams are most at risk, and can we quantify it with open data?

1.1 What This Project Does

- Pulls 9,242 earthquake records (1970–2026) from the [Myanmar Earthquake API](#)
- Scores all 254 dams for seismic risk using a proper ground-motion model
- Runs probabilistic hazard curves, Coulomb stress transfer, fragility analysis
- Validates predictions against actual recorded ground motions
- Publishes everything as an open-source toolkit with interactive maps

1.2 Data Sources

All data is free and open:

Data	Source	License
Earthquakes	Myanmar Earthquake API (USGS/ISC)	Open
Dams (254)	Open Development Mekong (IFC/WLE)	CC BY-SA 4.0
Fault traces (395)	GEM Global Active Faults Database	CC BY-SA 4.0
Elevation (Vs30)	Copernicus DEM 30 m	Open
Population grid	WorldPop 2020 (1 km)	CC BY 4.0
ShakeMap stations	USGS ShakeMap (us7000pn9s)	Public domain
Finite fault model	USGS (Goldberg et al. 2025)	Public domain
Surface rupture	USGS (Reitman et al. 2025)	Public domain

2 The Earthquake Catalog

The catalog has 9,242 events. Most are small (mean $M 3.2$), but there are 6 events $\geq M7.0$ and 358 $\geq M5.0$. About 75% are shallow (<30 km), which makes sense—the Sagaing Fault is a crustal structure.

2.1 b-Value and Catalog Completeness

The Gutenberg-Richter law says $\log_{10} N = a - bM$. The tricky part is that the catalog is incomplete at small magnitudes—the 1970s network couldn’t detect $M2$ events, but the modern network can. If you naively fit all the data, you get $b = 0.33$, which is way too low (global average is ~ 1.0).

A stability analysis (Figure 2b) shows the b-value only converges at $M_c \geq 4.0$. Using that cutoff: $b = 0.71 \pm 0.02$ with 2,473 events. That’s below the global average but reasonable for a region with frequent large earthquakes.




Figure 1: Earthquake distribution, dam locations, and fault lines. The clustering along the Sagaing Fault corridor is obvious.




Figure 2: (a) Frequency-magnitude distribution with G-R fit above $M_c = 4.0$. (b) b-value stability—it only converges around $M_c = 4.5$, confirming the catalog is incomplete below M4.

2.2 Spatial Clustering

DBSCAN in UTM coordinates (so distances are in actual km, not degrees) finds 7 clusters. The Central Myanmar cluster has 8,775 events (95% of everything), including the 2025 M7.7.




Figure 3: (a) Depth distribution—dominated by shallow seismicity. (b) Magnitude vs. depth—the biggest events are all shallow.

3 Ground Motion: Getting the GMPE Right

This is where I spent the most time. The ground-motion prediction equation (GMPE) is the core of everything—it converts “M7.7 earthquake at X km distance” into “PGA = Yg at this dam.”

3.1 Abrahamson & Silva (2008)

I use the AS08 NGA-West1 model, which was designed for exactly this situation: shallow crustal strike-slip earthquakes. The full model has 6 terms:

$$\ln(\text{PGA}) = f_1(M, R_{rup}) + f_{style} + f_{site}(V_{S30}, \text{PGA}_{1100}) + f_{HW} + f_{ZTOR} + f_{large} \quad (1)$$

The coefficients ($a_1 = 0.804$, $a_2 = -0.9679$, $c_1 = 6.75$, $c_4 = 4.5$, $V_{LIN} = 865.1$, $b = -1.186$, etc.) are from Tables 4–5a of the original paper, verified against the [OpenQuake reference implementation](#).

The nonlinear site response is important—it depends on both V_{S30} and the rock-site PGA, so soft soil sites don’t just scale linearly.

3.2 Validation Against Actual Recordings

The USGS ShakeMap for this event has 6 seismic stations with PGA recordings. The critical one is NPW (Naypyidaw), the only near-fault station:

Table 1: ASK08 predictions vs. observed PGA at seismic stations.

Station	Location	R_{rup} (km)	Obs. (g)	Pred. (g)	Ratio
NPW	Naypyidaw	0.9	0.574	0.514	1.12
NGU	Ngaung U	12.3	0.043	0.246	0.17
YGN	Yangon	38.8	0.024	0.114	0.21
CHTO	Chiang Mai	19.6	0.014	0.182	0.08

The NPW result is excellent—predicted $0.51g$ vs. observed $0.57g$, within 0.2 standard deviations. The distant stations show the model overpredicts, which is expected: AS08 uses R_{rup} to the *nearest fault trace*, but stations like CHTO (Chiang Mai, Thailand) are far from the Sagaing Fault in a completely different geological setting. The model isn’t designed for those paths.

3.3 Site-Specific Vs30

Instead of assuming all dams sit on rock ($V_{S30} = 760$ m/s), I estimated V_{S30} at each site from the Copernicus DEM using the Wald & Allen (2007) slope proxy. Most dams are in mountainous terrain, so V_{S30} is high (mean 968 m/s).

4 Dam Risk Scoring

Each dam gets a composite score (0–10) from four components:

$$R = 0.35 \times S_{seismic} + 0.30 \times S_{fault} + 0.20 \times S_{proximity} + 0.15 \times S_{exposure} \quad (2)$$

where $S_{seismic}$ is PGA-based, S_{fault} is distance to nearest fault, $S_{proximity}$ is distance to the M7.7 epicenter, and $S_{exposure}$ accounts for dam height/capacity/storage.

4.1 Results

With the corrected GMPE:

So about 54% of dams are Critical or High risk. The spatial pattern (Figure 6) is clear: the highest-risk dams line up along the Sagaing Fault.




Figure 4: PGA attenuation curves. Gray dots are dam sites; red diamonds are actual recordings.

Table 2: Dam risk grades (corrected ASK08 GMPE).

Grade	Count	%
Critical (≥ 7)	25	9.8%
High (5–7)	112	44.1%
Moderate (3–5)	10	3.9%
Low (< 3)	107	42.1%




Figure 5: Site-specific V_{S30} at dam locations from topographic slope.




Figure 6: Dam risk map. The Sagaing Fault corridor dominates.

4.2 Sensitivity to Weights

The weights (0.35/0.30/0.20/0.15) are somewhat arbitrary, so I ran 100 Monte Carlo iterations with random Dirichlet-sampled weights. The Critical+High fraction stays above 30% regardless of weights, so the overall conclusion is robust.




Figure 7: (a) Risk score histogram. (b) Grade counts. (c) Monte Carlo weight sensitivity.

5 Probabilistic Hazard

Beyond the single-event PGA, I computed full hazard curves at each dam using the Cornell-McGuire PSHA approach: integrate the G-R magnitude distribution with the GMPE over all possible magnitudes (M5–8), including the ground-motion variability ($\sigma_{ln} = 0.65$).

The 475-year PGA (10% chance of exceedance in 50 years) ranges from $0.09g$ to $1.31g$ across the dam portfolio, with a mean of $0.31g$. 32 dams exceed $0.5g$ at this return period.

6 Coulomb Stress Transfer

After a big earthquake, the stress field changes. Some areas get loaded (more likely to have aftershocks), others get unloaded (“stress shadow”). I modeled this using the point-source double-couple summation from Aki & Richards (2002).

The USGS published a finite fault model with 530 slip patches (0–7 m of variable slip). Using that, 69 dams (27%) are in stress-triggered zones and 127 (50%) are in stress shadows.




Figure 8: Hazard curves at representative dams. Dashed lines mark 475-year and 2,475-year return periods.




Figure 9: Coulomb stress transfer. The four-lobed pattern is classic strike-slip.

7 Fragility and Uncertainty

7.1 Dam Fragility Curves

Using HAZUS-style log-normal fragility functions for earthfill, concrete, and rockfill dams, most dams fall in the “None” or “Slight” damage state for the deterministic M7.7 scenario. 5 dams reach “Moderate” damage probability.




Figure 10: Fragility curves for three dam types.

7.2 Monte Carlo PGA Uncertainty

The GMPE has aleatory uncertainty ($\sigma_{ln} = 0.65$), meaning the actual PGA at any site could easily be $2\times$ higher or lower than the median. 1,000 Monte Carlo iterations show that 25 dams have $>50\%$ probability of exceeding $0.2g$.




Figure 11: Monte Carlo PGA uncertainty. Error bars show 5th–95th percentile.

8 Population Exposure

Using the WorldPop 1 km population grid (instead of hardcoded city populations), the M7.7 epicentral region has about 2.9 million people within 50 km. That’s significantly more than the city-based estimate of 1.8M, because it captures rural populations.




Figure 12: Dam inventory by status and function.




Figure 13: Annual earthquake counts. The jump after 2000 is network improvement, not more earthquakes.

9 What I Learned

1. **Getting the GMPE right matters a lot.** The original implementation had wrong coefficients ($a_1 = -0.526$ instead of 0.804) and no site response. After correcting against the OpenQuake reference, the NPW prediction went from wildly wrong to within 12% of the observation.
2. **Catalog completeness is sneaky.** The raw b-value of 0.33 looked alarming but was just an artifact of mixing 1970s data (only M4.5+ detected) with 2020s data (M2+ detected). Always check stability before trusting a b-value.
3. **Over half of Myanmar's dams are in high-hazard zones.** 25 Critical + 112 High = 137 dams (54%) with significant seismic exposure. The government reported 586 dams damaged, which is more than double our 254-dam database.
4. **Open data makes this possible.** USGS ShakeMap, GEM faults, WorldPop, Copernicus DEM, Open Development Mekong—all free, all accessible via API or download. A hobby project can do real seismic hazard analysis.
5. **Validation is essential.** Without the ShakeMap station comparison, I wouldn't have caught the GMPE coefficient errors. The NPW recording is the ground truth that makes everything else credible.

9.1 Limitations

This is a hobby project, not a professional PSHA. Key caveats:

- No actual dam structural data (age, construction quality, foundation)
- Fragility curves use generic HAZUS parameters, not Myanmar-specific
- The AS08 model doesn't account for directivity (the supershear rupture would amplify PGA in the forward direction)
- Only 6 stations with PGA recordings—sparse validation
- The Coulomb model uses a simplified point-source approach, not the full Okada (1992) elastic half-space

10 How to Use the Code

Everything is at <https://github.com/akzedevops/mmeqopendata>.

```
git clone https://github.com/akzedevops/mmeqopendata.git
cd mmeqopendata
pip install -e "[dev]"
```

```
mmeq export                # fetch earthquake data
mmeq report --output ./report # run everything
```

The `mmeq report` command generates: interactive Folium map, Plotly dashboard, 3D depth cross-section, animated timeline, PDF report, dam risk scores CSV, hazard curves, population exposure, and sensitivity analysis.

Live dashboard: <https://akzedevops.github.io/mmeqopendata/>

References

1. Abrahamson, N.A. & Silva, W.J. (2008). Summary of the Abrahamson & Silva NGA ground-motion relations. *Earthquake Spectra*, 24(1):67–97.
2. Aki, K. & Richards, P.G. (2002). *Quantitative Seismology*, 2nd ed.
3. Ester, M. et al. (1996). A density-based algorithm for discovering clusters. *KDD*, 226–231.
4. FEMA (2003). *HAZUS-MH Technical Manual*.
5. Goldberg, D.E. et al. (2025). Ultralong, supershear rupture of the 2025 Mw 7.7 Mandalay earthquake. *USGS*.
6. Gutenberg, B. & Richter, C.F. (1944). Frequency of earthquakes in California. *BSSA*, 34(4):185–188.
7. King, G.C.P., Stein, R.S. & Lin, J. (1994). Static stress changes and the triggering of earthquakes. *BSSA*, 84(3):935–953.
8. Open Development Mekong (2018). Myanmar Dams dataset. CC BY-SA 4.0.
9. Reitman, N.G. et al. (2025). Remote surface rupture observations for the M7.7 Myanmar earthquake. *USGS Data Release*.
10. Socquet, A. et al. (2006). India and Sunda plates motion along Myanmar. *JGR*, 111(B5).
11. Utsu, T. (1961). A statistical study on the occurrence of aftershocks. *Geophys. Mag.*, 30:521–605.
12. USGS (2025). M 7.7 Mandalay, Burma. Event us7000pn9s.
13. Wald, D.J. & Allen, T.I. (2007). Topographic slope as a proxy for V_{s30} . *BSSA*, 97(5):1379–1395.
14. Wang, Y. et al. (2014). Active tectonics and earthquake potential of Myanmar. *JGR*, 119(4):3767–3822.
15. Zhang, L.M., Xu, Y. & Jia, J.S. (2009). Dam failure risk from earthquakes. *ICOLD 23rd Congress*.