

# pylab4-1

February 10, 2020

## 0.1 Numerical Integration Methods

Session 1 Jeff Shen | 1004911526

Stacy Ossipov | 1004877779

10 Feb 2020

```
[1]: import numpy as np
import matplotlib.pyplot as plt
%matplotlib inline
import pandas as pd
```

```
[2]: # velocity function
def v(x):
    return np.diff(x)

# acceleration function
def a(x):
    return np.diff(np.diff(x))
```

```
[3]: # change in time (between samples) in seconds
dt = 0.01

#initial time in seconds
t0 = 0

# total/end time in seconds
t = 10

# period. seconds / # of oscillations
p = t/14.5


# mass in kg
m = 0.2

# angular frequency
omega = 2*np.pi / p
```

```
# k (spring constant)
k = np.square(omega) * m
```

```
[4]: data = pd.read_csv('data.txt', delimiter='\t')
data = data['Sample'][250:1250]
data = data.reset_index().Sample
```

```
[5]: plt.plot(data)
plt.title('Position vs time of measured data')
plt.xlabel('Time (centiseconds)')
plt.ylabel('Position (centimeters)')
plt.savefig('timepos0.png')
```




```
[6]: # initializing arrays for numerical integration
ys = np.zeros(1000)
vs = np.zeros(1000)

ys[0] = data[0]
vs[0] = v(data)[0]
```


```
[7]: # numerical integration
for i in np.arange(999):
    ys[i+1] = ys[i] + dt * vs[i]
```

```
vs[i+1] = vs[i] - dt * np.square(omega) * ys[i]
```


```
[8]: plt.plot(np.arange(0, t, dt), ys)
plt.title('Time vs position of simulated data (Forward Euler)')
plt.xlabel('Time (s)')
plt.ylabel('Position (cm)')
plt.savefig('timepos1.png')
```



```
[9]: plt.plot(np.arange(0, t, dt), vs)
plt.title('Time vs velocity of simulated data (Forward Euler)')
plt.xlabel('Time (s)')
plt.ylabel('Velocity (cm/s)')
plt.savefig('timevel1.png')
```




```
[10]: plt.plot(ys, vs)
plt.title('Position vs velocity of simulated data (Forward Euler)')
plt.xlabel('position (cm)')
plt.ylabel('velocity (cm/s)')
plt.savefig('phase1.png')
```



```
[11]: # calculating energy
e = 1/2 * m * np.square(vs) + 1/2 * k * np.square(ys)
```

```
[12]: plt.plot(e)
plt.title('Energy plot of simulated data (Forward Euler)')
plt.xlabel('Time (centiseconds)')
plt.ylabel(r'Energy ($kg\,cm^2\,s^{-2}$)')
plt.savefig('eplot1.png')
```




```
[13]: # initializing second array
ys2 = np.zeros(1000)
vs2 = np.zeros(1000)


ys2[0] = data[0]
vs2[0] = v(data)[0]
```

```
[14]: # Euler-Cromer method
for i in np.arange(999):
    ys2[i+1] = ys2[i] + dt * vs2[i]
    vs2[i+1] = vs2[i] - dt * k / m * ys2[i+1]
```

```
[15]: # position plot of euler-cromer
plt.plot(np.arange(0, t, dt), ys2)
plt.title('Time vs position of simulated data (Euler-Cromer)')
plt.xlabel('Time (s)')
plt.ylabel('Position (cm)')
plt.savefig('timepos2.png')
```




```
[16]: # velocity plot of euler-cromer
plt.plot(np.arange(0, t, dt), vs2)
plt.title('Time vs velocity of simulated data (Euler-Cromer)')
plt.xlabel('Time (s)')
plt.ylabel('Velocity (cm/s)')
plt.savefig('timevel2.png')
```




```
[17]: plt.plot(ys2, vs2)
plt.title('Position vs velocity of simulated data (Euler-Cromer)')
plt.xlabel('Position (cm)')
plt.ylabel('Velocity (cm/s)')
plt.savefig('phase2.png')
```





```
[18]: # calculating energy
e2 = 1/2 * m * np.square(vs2) + 1/2 * k * np.square(ys2)
```

```
[19]: plt.plot(e2)
plt.title('Energy plot of simulated data (Euler-Cromer)')
plt.xlabel('Time (centiseconds)')
plt.ylabel(r'Energy ($kg\,cm^2\,s^{-2}$)')
plt.savefig('eplot2.png')
```



```
[20]: print('Values from experiment:')
      print(f'Period: {p:.2f} s')
      print(f'Frequency: {1/p:.2f} cycles/s')
      print(f'Amplitude: {ys2.max() - ys2.mean():.2f} cm')
      print(f'Spring constant: {k:.2f} kg cm2 / s2')

```

Values from experiment:  
 Period: 0.69 s  
 Frequency: 1.45 cycles/s  
 Amplitude: 21.60 cm  
 Spring constant: 16.60 kg cm<sup>2</sup> / s<sup>2</sup>