

Concept	1-Categorical construction	$\infty$ -Categorical construction	Intuition
Accessible category	$\mathcal{C}$ is locally small, admits $\kappa$ -filtered colimits, and there is a set of $\kappa$ -compact objects that generate the category under $\kappa$ -filtered colimits. ([nLa25a], Def 2.1)	$\mathcal{C}$ is locally small, admits $\kappa$ -filtered colimits, the full subcategory $\mathcal{C}^\kappa \subseteq \mathcal{C}$ of $\kappa$ -compact objects is essentially small, and $\mathcal{C}^\kappa$ generates $\mathcal{C}$ under small, $\kappa$ -filtered colimits <sup>1</sup> . ([Lur09], Prop 5.4.2.2)	[todo]
$F$ -Cartesian edge	$F : X \rightarrow S$ a functor, $f : x \rightarrow y$ a morphism in $X$ is $F$ -cartesian if the induced map $X_{/f} \rightarrow X_{/y} \times_{S_{/F(y)}} S_{/F(f)}$ is an isomorphism of categories. ([nLa25b], Prop 2.4)	$F : X \rightarrow S$ an inner fibration of simplicial sets, $f : x \rightarrow y$ an edge in $X$ is $F$ -cartesian if the induced map $X_{/f} \rightarrow X_{/y} \times_{S_{/F(y)}} S_{/F(f)}$ is a trivial Kan fibration. ([Lur09], Def 2.4.1.1)	In the model structure on <b>sSet</b> , the fibrations are Kan fibrations and the weak equivalences are weak homotopy equivalences ([Lur09], A.2.7). A trivial fibration in a model category is a map which is both a fibration and a weak equivalence, which in <b>sSet</b> is equivalent to the definition given in this table. Thus, being related by a Kan fibration is a higher categorical notion of ‘sameness’. <b>Why not a categorical equivalence?</b> [Lur09] Rem 1.2.5.5 implies this is stronger, which would match more with the fact that the 1-categorical version is defined in terms of an isomorphism (not equivalence) of categories.
Category	Collection of objects $\mathcal{C}$ , set $\text{Hom}(X, Y)$ for every $X, Y \in \mathcal{C}$ , associative composition and identity morphisms	Simplicial set $\mathcal{C} : \Delta^{\text{op}} \rightarrow \mathbf{Set}$ satisfying the inner horn filling condition. ([Lur09], Def 1.1.2.4)	Category with objects $\mathcal{C}_0$ , morphisms $\mathcal{C}_1$ , morphisms between morphisms $\mathcal{C}_2$ , etc. Inner horn filling defines a non-unique composition.
$F$ -Cocartesian edge	$F : X \rightarrow S$ a functor, $f : x \rightarrow y$ a morphism in $X$ is $F$ -cocartesian if the induced map $X_{f/} \rightarrow X_{x/} \times_{S_{F(y)/}} S_{F(f)/}$ is an isomorphism of categories. ([nLa25b], Prop 2.4)	$F : X \rightarrow S$ an inner fibration of simplicial sets, $f : x \rightarrow y$ an edge in $X$ is $F$ -cocartesian if the induced map $X_{f/} \rightarrow X_{x/} \times_{S_{F(y)/}} S_{F(f)/}$ is a trivial Kan fibration. ([Lur09], Def 2.4.1.1 / Prop 2.4.1.8)	Note that the definitions of an inner fibration and a Kan fibration are invariant under taking opposites. For other intuition, see: $F$ -cartesian edge.
Colimit	A colimit for $F : J \rightarrow \mathcal{C}$ is an initial cocone on $F$ .	A colimit for $F : X \rightarrow \mathcal{C}$ ( $X$ a simplicial set, $\mathcal{C}$ an $\infty$ -category) is an initial object of $\mathcal{C}_{F/}$ . ([Lur09], Def 1.2.13.4)	The obvious extension of the definition of the undercategory $\mathcal{C}_{C/}$ for $C : \{*\} \rightarrow \mathcal{C}$ to $\mathcal{C}_{F/}$ for an arbitrary functor $F : J \rightarrow \mathcal{C}$ ends up being exactly <b>Cocone</b> ( $F$ ).
$\kappa$ -Compact object	Let $C \in \mathcal{C}$ , and let $j_C : \mathcal{C} \rightarrow \mathbf{Set}$ denote the functor represented by $C$ . If $\mathcal{C}$ admits $\kappa$ -filtered colimits, then $C$ is $\kappa$ -compact if $j_C$ commutes with filtered colimits. ([Lur09], 5.3.4)	Let $C \in \mathcal{C}$ , and let $j_C : \mathcal{C} \rightarrow \hat{\mathcal{S}}$ denote the functor <b>represented by</b> $C$ . If $\mathcal{C}$ admits $\kappa$ -filtered colimits, then $C$ is $\kappa$ -compact if $j_C$ <b>preserves</b> $\kappa$ -filtered colimits. <sup>2</sup> ([Lur09], Def 5.3.4.5)	[todo]
Essentially small category	$\mathcal{C}$ equivalent to a small category.	$\mathcal{C}$ categorically equivalent to a small $\infty$ -category. [Lur25, Def 03SJ]	-
Essentially surjective functor	$F : \mathcal{C} \rightarrow \mathcal{D}$ is essentially surjective if for every $D \in \mathcal{D}$ , there exists some $C \in \mathcal{C}$ with $FC \cong D$ .	$F : \mathcal{C} \rightarrow \mathcal{D}$ is essentially surjective if $\text{h}F : \text{h}\mathcal{C} \rightarrow \text{h}\mathcal{D}$ is essentially surjective. ([Lur09], Def 1.2.10.1)	Essentially surjective up to homotopy.

<sup>1</sup>I.e. for any full subcategory  $\mathcal{C}' \subseteq \mathcal{C}$  containing every element of  $\mathcal{C}^\kappa$ , if  $\mathcal{C}'$  is stable under small  $\kappa$ -filtered colimits, then  $\mathcal{C}' = \mathcal{C}$ . ([Lur09], p328)

<sup>2</sup>Lurie introduces the term  $\kappa$ -continuous for such functors, but in ordinary category theory this generally means a functor which preserves  $\kappa$ -small limits; a functor preserving  $\kappa$ -filtered colimits is called  $\kappa$ -finitary. I have thus steered clear of this term.

Faithful functor	$F : \mathcal{C} \rightarrow \mathcal{D}$ is faithful if $\text{Hom}(X, Y) \rightarrow \text{Hom}(FX, FY)$ is injective for all $X, Y \in \mathcal{C}$ .	$F : \mathcal{C} \rightarrow \mathcal{D}$ is faithful if $hF : h\mathcal{C} \rightarrow h\mathcal{D}$ is faithful. ([Lur09], Def 1.2.10.1)	Faithful up to homotopy.
$\kappa$ -Filtered category	For a regular cardinal $\kappa$ , $\mathcal{C}$ is $\kappa$ -filtered if, for every $\kappa$ -small category $J$ and every functor $F : J \rightarrow \mathcal{C}$ , there exists a cocone on $F$ .	For a regular cardinal $\kappa$ , $\mathcal{C}$ is $\kappa$ -filtered if, for every $\kappa$ -small simplicial set $X$ and every map $f : X \rightarrow \mathcal{C}$ , there exists a map $\bar{f} : K^\triangleright \rightarrow \mathcal{C}$ extending $f$ . ([Lur09], Def 5.3.1.7)	A cocone on $F$ is a collection of compatible maps $(\lambda_j : F(j) \rightarrow C)$ . Define $\bar{F} : J \star [0] \rightarrow \mathcal{C}$ to be $F$ on $J$ , send the cone point to $C$ , and send the unique morphisms $*_j$ from $j \in J$ to the cone point to the $\lambda_j$ . Conversely, if you have some $\bar{F}$ extending $F$ , define $\lambda_j := F(*_j)$ .
Final object	Object $C \in \mathcal{C}$ such that for any other object $C' \in \mathcal{C}$ , there exists a unique morphism $C' \rightarrow C$ .	Object $C \in \mathcal{C}$ such that $C$ is final in $h\mathcal{C}$ , regarded as an enriched category over $\mathcal{H}$ . ([Lur09], Def 1.2.12.1)	Object $C \in \mathcal{C}$ such that for any other object $C' \in \mathcal{C}$ , there exists a unique (up to homotopy) morphism $C' \rightarrow C$ .
Full functor	$F : \mathcal{C} \rightarrow \mathcal{D}$ is full if $\text{Hom}(X, Y) \rightarrow \text{Hom}(FX, FY)$ is surjective for all $X, Y \in \mathcal{C}$ .	$F : \mathcal{C} \rightarrow \mathcal{D}$ is full if $hF : h\mathcal{C} \rightarrow h\mathcal{D}$ is full. ([Lur09], Def 1.2.10.1)	Full up to homotopy.
Functor	Functor.	Natural transformation of simplicial sets. ([Lur09], 1.2.7)	-
Groupoid	Category whose morphisms are all invertible.	Kan complex.	Not only can you find (non-unique) ‘composites’, but you can also fill in diagrams like $\begin{array}{ccc} C & \xrightarrow{\text{id}} & C \\ f \downarrow & \nearrow & \\ D & & \end{array}$ $\begin{array}{ccc} C & \xrightarrow{\text{id}} & D \\ \downarrow & \nearrow f & \\ C & & \end{array}$
Initial object	Object $C \in \mathcal{C}$ such that for any other object $C' \in \mathcal{C}$ , there exists a unique morphism $C \rightarrow C'$ .	Object $C \in \mathcal{C}$ such that $C$ is initial in $h\mathcal{C}$ , regarded as an enriched category over $\mathcal{H}$ . ([Lur09], Def 1.2.12.1)	Object $C \in \mathcal{C}$ such that for any other object $C' \in \mathcal{C}$ , there exists a unique (up to homotopy) morphism $C \rightarrow C'$ .
Join	$\mathcal{C} \star \mathcal{D}$ has objects $\text{ob } \mathcal{C} \sqcup \text{ob } \mathcal{D}$ , and $\text{Hom}_{\mathcal{C} \star \mathcal{D}}(X, Y)$ is given by: $\begin{cases} \text{Hom}_{\mathcal{C}}(X, Y) & X, Y \in \mathcal{C}, \\ \text{Hom}_{\mathcal{D}}(X, Y) & X, Y \in \mathcal{D}, \\ \emptyset & X \in \mathcal{D}, Y \in \mathcal{C}, \\ * & X \in \mathcal{C}, Y \in \mathcal{D}. \end{cases} \quad ([\text{Lur09}], 1.2.8)$	$\mathcal{C} \star \mathcal{D}$ has $n$ -simplices $(\mathcal{C} \star \mathcal{D}) = \mathcal{C}_n \cup \mathcal{D}_n \cup \bigcup_{i+j=n-1} \mathcal{C}_i \times \mathcal{D}_j$ . The $i$ th boundary map $d_i : (\mathcal{C} \star \mathcal{D})_n \rightarrow (\mathcal{C} \star \mathcal{D})_{n-1}$ is defined on $\mathcal{C}_n$ and $\mathcal{D}_n$ using the $i$ th boundary map on $\mathcal{C}$ and $\mathcal{D}$ . Given $\sigma \in S_j, \tau \in T_k$ , $d_i(\sigma, \tau)$ is given by $\begin{cases} (d_i\sigma, \tau) & i \leq j, j \neq 0, \\ (\sigma, d_{i-j-1}\tau) & i > j, k \neq 0. \end{cases}$ If $j = 0$ , then $d_0(\sigma, \tau) = \tau$ , and if $k = 0$ , then $d_n(\sigma, \tau) = \sigma$ . ([Lur09], Def 1.2.8.1 / [nLa25c])	Objects are in both cases disjoint unions of objects from the two categories being joined. Morphisms are also exactly the same in both cases (you get all the morphisms from $\mathcal{C}$ and $\mathcal{D}$ , plus a morphism from $c \rightarrow d$ for every pair $(c, d) \in \mathcal{C}_0 \times \mathcal{D}_0$ ). Whenever you have an $n$ -simplex in $\mathcal{C}$ and an $m$ -simplex in $\mathcal{D}$ , you get an $(m+n+1)$ -simplex in $\mathcal{C} \star \mathcal{D}$ , so in particular $\Delta^n \star \Delta^m \cong \Delta^{m+n+1}$ .
Left cone	$\mathcal{C}^\triangleleft := [0] \star \mathcal{C}$ .	$\mathcal{C}^\triangleleft := \Delta^0 \star \mathcal{C}$ . ([Lur09], Not 1.2.8.4)	$\mathcal{C}$ with extra vertex (cone point) added, as well as a map from that cone point to every other vertex in $\mathcal{C}$ (plus obligatory degenerate simplices).

Left dualisable object	Object $C \in \mathcal{C}$ such that there exists some $C^* \in \mathcal{C}$ and maps $e : C \otimes C^* \rightarrow \mathbb{1}$ , $c : \mathbb{1} \rightarrow C^* \otimes C$ such that the composites $C \rightarrow C \otimes C^* \otimes C \rightarrow C$ and $C^* \rightarrow C^* \otimes C \otimes C^* \rightarrow C^*$ are equal to the identity. <sup>3</sup>	Object $C \in \mathcal{C}$ such that there exists some $C^* \in \mathcal{C}$ and maps $e : C \otimes C^* \rightarrow \mathbb{1}$ , $c : \mathbb{1} \rightarrow C^* \otimes C$ such that the composites $C \rightarrow C \otimes C^* \otimes C \rightarrow C$ and $C^* \rightarrow C^* \otimes C \otimes C^* \rightarrow C^*$ are homotopic to the identity.	$C$ has a left dual up to homotopy.
Left Kan extension (along the inclusion of a full subcategory)	<p>Given a commutative diagram <math display="block">\begin{array}{ccc} \mathcal{C}^0 &amp; \xrightarrow{F_0} &amp; \mathcal{D} \\ \downarrow \iota &amp; \nearrow F &amp; \\ \mathcal{C} &amp; &amp; \end{array},</math></p> <p><math>F</math> is a left Kan extension of <math>F_0</math> along <math>\iota</math> if there is a natural transformation <math>\eta : F_0 \rightarrow F\iota</math> such that for any other pair <math>(G : \mathcal{C} \rightarrow \mathcal{D}, \gamma : F_0 \rightarrow G\iota)</math>, there exists a unique natural transformation <math>\alpha : F \rightarrow G</math> such that <math>\gamma = (\alpha * \iota) \circ \eta</math>. ([Rie16], Def 6.1.1)</p>	<p>Given a commutative diagram <math display="block">\begin{array}{ccc} \mathcal{C}^0 &amp; \xrightarrow{F_0} &amp; \mathcal{D} \\ \downarrow \iota &amp; \nearrow F &amp; \\ \mathcal{C} &amp; &amp; \end{array},</math></p> <p><math>F</math> is a left Kan extension of <math>F_0</math> along <math>\iota</math> if for all <math>C \in \mathcal{C}</math>, the induced diagram <math display="block">\begin{array}{ccc} \mathcal{C}_{/C}^0 &amp; \xrightarrow{F_C} &amp; \mathcal{D} \\ \downarrow &amp; \nearrow &amp; \\ (\mathcal{C}_{/C}^0)^{\triangleright} &amp; &amp; \end{array}</math> exhibits <math>FC</math> as a colimit of <math>F_C</math>. ([Lur09], Def 4.3.2.2)</p>	In the 1-categorical case, the colimits of $\mathcal{C}_{/C}^0 \rightarrow \mathcal{C}^0 \xrightarrow{F_0} \mathcal{D}$ for each $C \in \mathcal{C}$ (if they all exist) define the left Kan extension of $F_0$ along $\iota$ ([Rie16], Thm 6.2.1) <sup>4</sup> . This is the case if and only if $F$ is a <i>pointwise</i> Kan extension ([Rie16], Thm 6.3.7), so really the higher categorical version generalises pointwise left Kan extensions (along the inclusion of a full subcategory).
Limit	A limit for $F : J \rightarrow \mathcal{C}$ is a terminal cone on $F$ .	A limit for $F : X \rightarrow \mathcal{C}$ ( $X$ a simplicial set, $\mathcal{C}$ an $\infty$ -category) is a final object of $\mathcal{C}_{/F}$ . ([Lur09], Def 1.2.13.4)	The obvious extension of the definition of the overcategory $\mathcal{C}_{/C}$ for $C : \{*\} \rightarrow \mathcal{C}$ to $\mathcal{C}_{/F}$ for an arbitrary functor $F : J \rightarrow \mathcal{C}$ ends up being exactly <b>Cone</b> ( $F$ ).
Locally small category	For every $X, Y \in \mathcal{C}$ , $\text{Hom}(X, Y)$ is a set.	For every $X, Y \in \mathcal{C}$ , the space $\text{Hom}(X, Y)$ is essentially small. ([Lur09], Prop 5.4.1.7)	-

<sup>3</sup>The opposite convention (i.e. that this defines a *right* dualisable object) is taken in [Eti+15], and by many other authors. I'm attempting to follow Lurie's conventions as far as possible.

<sup>4</sup>I think Lurie is saying exactly the same thing in the  $\infty$ -categorical case, just in a slightly confusing way.

Monoidal category	<p>Opfibration of categories <math>p : \mathcal{C}^\otimes \rightarrow \Delta^{\text{op}}</math> such that, for each <math>n \geq 0</math>, the associated functors <math>\mathcal{C}_{[n]}^\otimes \rightarrow \mathcal{C}_{\{i,i+1\}}^\otimes</math> determine an equivalence of categories <math>\mathcal{C}_{[n]}^\otimes \rightarrow \mathcal{C}_{\{0,1\}}^\otimes \times \dots \times \mathcal{C}_{\{n-1,n\}}^\otimes \simeq (\mathcal{C}_{[1]}^\otimes)^n</math>.</p>	<p>Cocartesian fibration of simplicial sets <math>p : \mathcal{C}^\otimes \rightarrow N(\Delta)^{\text{op}}</math> such that, for each <math>n \geq 0</math>, the associated functors <math>\mathcal{C}_{[n]}^\otimes \rightarrow \mathcal{C}_{\{i,i+1\}}^\otimes</math> determine an equivalence<sup>5</sup> of <math>\infty</math>-categories</p> $\mathcal{C}_{[n]}^\otimes \rightarrow \mathcal{C}_{\{0,1\}}^\otimes \times \dots \times \mathcal{C}_{\{n-1,n\}}^\otimes \simeq (\mathcal{C}_{[1]}^\otimes)^n$ <p>(<a href="#">[Lur07]</a>, Def 1.1.2)<sup>6</sup></p>	<p>In the 1-categorical case, you recover the original category by setting <math>\mathcal{C} := \mathcal{C}_{[1]}^\otimes</math>. The unit is <math>\mathcal{C}_{[0]}^\otimes</math>, the tensor product <math>\otimes</math> is induced by the outer inclusion <math>\{0 &lt; 2\} = [1] \subseteq [2]</math>, the unitors and associators come from the commutative diagrams</p> $  \begin{array}{ccccc}  \{0, 1\} & \longleftarrow & \{0, 1, 2\} & \longrightarrow & \{1, 2\} \\  & \nwarrow & \uparrow & \nearrow & \\  & & \{0, 2\} & & \\  \\   \{0, 3\} & \longrightarrow & \{0, 1, 3\} & & \\  \downarrow & & \downarrow & & \\  \{0, 2, 3\} & \longrightarrow & \{0, 1, 2, 3\} & &   \end{array}  $ <p>in <math>\Delta</math>. Conversely, given a monoidal category <math>\mathcal{C}</math>, define <math>\mathcal{C}^\otimes</math> to have objects finite sequences <math>[C_1, \dots, C_n]</math> of objects of <math>\mathcal{C}</math>, and a morphism <math>[C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]</math> to be a map <math>[m] \rightarrow [n]</math> and a collection of morphisms <math>C_{f(i-1)+1} \otimes \dots \otimes C_{f(i)} \rightarrow C'_i</math> for <math>1 \leq i \leq m</math>.<sup>7</sup> Then the forgetful functor to <math>\Delta^{\text{op}}</math> is the required cocartesian fibration. (<a href="#">[Lur07]</a>, p5-6)</p>
-------------------	--	--	--

<sup>5</sup>Weak or categorical?

<sup>6</sup>I consider this a temporary definition, because I can't yet reconcile it with [\[Lur17\]](#), Def 4.1.1.10.

<sup>7</sup>You “tensor along the gap”, if that makes any sense.

(Coloured) operad	<ul style="list-style-type: none"> <li>• A collection of objects <math>\mathcal{O}</math>.</li> <li>• For every finite set <math>I</math>, every <math>I</math>-indexed collection of objects <math>\{X_i\}_{i \in I}</math> of <math>\mathcal{O}</math>, and every <math>Y \in \mathcal{O}</math>, a set <math>\text{Hom}(\{X_i\}_{i \in I}, Y)</math>.</li> <li>• For every map of finite sets <math>I \rightarrow J</math> having fibres <math>\{I_j\}_{j \in J}</math>, every finite collection of objects <math>\{X_i\}_{i \in I}</math>, every finite collection of objects <math>\{Y_j\}_{j \in J}</math>, and every object <math>Z \in \mathcal{O}</math>, a composition map <math>\prod_j \text{Hom}(\{X_i\}_{i \in I_j}, Y_j) \times \text{Hom}(\{Y_j\}_j, Z) \rightarrow \text{Hom}(\{X_i\}_i, Z)</math>, which is associative.</li> <li>• Units <math>\text{id}_X \in \text{Hom}(\{X\}, X)</math>. ([Lur17], Def 2.1.1.1)</li> </ul>	<p>Functor <math>p : \mathcal{O}^\otimes \rightarrow N(\mathbf{Fin}_*)</math> between <math>\infty</math>-categories which satisfies the following conditions:</p> <ul style="list-style-type: none"> <li>• For every inert morphism <math>f : \langle m \rangle \rightarrow \langle n \rangle</math> in <math>N(\mathbf{Fin}_*)</math> and every object <math>C \in \mathcal{O}_{\langle m \rangle}^\otimes</math>, there exists a <math>p</math>-cocartesian morphism <math>\bar{f} : C \rightarrow C'</math> in <math>\mathcal{O}^\otimes</math> lifting <math>f</math>.</li> <li>• Let <math>C \in \mathcal{O}_{\langle m \rangle}^\otimes</math> and <math>C' \in \mathcal{O}_{\langle n \rangle}^\otimes</math> be objects, let <math>f : \langle m \rangle \rightarrow \langle n \rangle</math> be a morphism in <math>\mathbf{Fin}_*</math>, and let <math>\text{Hom}_{\mathcal{O}^\otimes}^f(C, C')</math> be the union of those connected components of <math>\text{Hom}_{\mathcal{O}^\otimes}(C, C')</math> which lie over <math>f \in \text{Hom}_{\mathbf{Fin}_*}(\langle m \rangle, \langle n \rangle)</math>. Choose <math>p</math>-cocartesian morphisms <math>C' \rightarrow C'_i</math> lying over the inert morphisms <math>\rho^i : \langle n \rangle \rightarrow \langle 1 \rangle</math> for <math>1 \leq i \leq n</math>. Then the induced map</li> </ul> $\text{Hom}_{\mathcal{O}^\otimes}^f(C, C') \rightarrow \prod_{1 \leq i \leq n} \text{Hom}_{\mathcal{O}^\otimes}^{\rho^i f}(C, C'_i)$ <p>is a homotopy equivalence.</p> <ul style="list-style-type: none"> <li>• For every finite collection of objects <math>C_1, \dots, C_n \in \mathcal{O}_{\langle 1 \rangle}^\otimes</math>, there exists an object <math>C \in \mathcal{O}_{\langle n \rangle}^\otimes</math> and a collection of <math>p</math>-cocartesian morphisms <math>C \rightarrow C_i</math> covering <math>\rho^i : \langle n \rangle \rightarrow \langle 1 \rangle</math>.</li> </ul>	[todo]
(Coloured) operad map	[todo]	<p>For <math>\infty</math>-operads <math>p : \mathcal{O}^\otimes \rightarrow N(\mathbf{Fin}_*)</math>, <math>q : \mathcal{O}'^\otimes \rightarrow N(\mathbf{Fin}_*)</math>, an <math>\infty</math>-operad map <math>\mathcal{O}^\otimes \rightarrow \mathcal{O}'^\otimes</math> is a map of simplicial sets <math>f : \mathcal{O}^\otimes \rightarrow \mathcal{O}'^\otimes</math> such that the diagram below commutes,</p> $\begin{array}{ccc} \mathcal{O}^\otimes & \xrightarrow{\quad} & \mathcal{O}'^\otimes \\ & \searrow \quad \swarrow & \\ & N(\mathbf{Fin}_*) & \end{array}$ <p>and the functor <math>f</math> carries inert morphisms<sup>8</sup> in <math>\mathcal{O}^\otimes</math> to inert morphisms in <math>\mathcal{O}'^\otimes</math>.</p>	[todo]
Opposite category	$\mathcal{C}^{\text{op}}$ has the same objects as $\mathcal{C}$ , and $\text{Hom}_{\mathcal{C}^{\text{op}}}(X, Y) = \text{Hom}_{\mathcal{C}}(Y, X)$ .	$\mathcal{C}_n^{\text{op}} = \mathcal{C}([n]^{\text{op}})$ , where $\{0 < 1 < \dots < n\}^{\text{op}} = \{0 > 1 > \dots > n\}$ . ([Lur09], 1.2.1)	A map $x \rightarrow y$ is an edge $\Delta^1 \rightarrow \mathcal{C}$ where $0 \mapsto x$ and $1 \mapsto y$ . In $\mathcal{C}^{\text{op}}$ 0 and 1 swap roles, so we instead get a map $y \rightarrow x$ .

<sup>8</sup>I.e.  $p$ -cocartesian morphisms  $g$  with  $p(g) : \langle m \rangle \rightarrow \langle n \rangle$  such that for each  $i \in \langle n \rangle^\circ$ , the inverse image  $g^{-1}\{i\}$  has exactly one element ([Lur17] Def 2.1.1.8 and 2.1.2.3).

Overcategory	<p>For <math>C \in \mathcal{C}</math>, the category <math>\mathcal{C}_{/C}</math> satisfies the following universal property: for any category <math>\mathcal{D}</math>, there is a bijection</p> $\mathrm{Hom}(\mathcal{D}, \mathcal{C}_{/C}) \simeq \mathrm{Hom}_C(\mathcal{D} \star [0], \mathcal{C}),$ <p>where the subscript on the right indicates that we consider only those functors <math>\mathcal{D} \star [0] \rightarrow \mathcal{C}</math> whose restriction to <math>[0]</math> coincides with <math>C</math>. ([Lur09], 1.2.9)</p>	<p>For <math>f : S \rightarrow \mathcal{C}</math>, <math>S</math> a simplicial set and <math>\mathcal{C}</math> an <math>\infty</math>-category, the <math>\infty</math>-category <math>\mathcal{C}_{/f}</math> satisfies the following universal property: for any simplicial set <math>X</math>, there is a bijection</p> $\mathrm{Hom}(X, \mathcal{C}_{/f}) \simeq \mathrm{Hom}_f(X \star S, \mathcal{C}),$ <p>where the subscript on the right indicates that we consider only those functors <math>X \star S \rightarrow \mathcal{C}</math> whose restriction to <math>S</math> coincides with <math>f</math>. Explicitly,</p> $(\mathcal{C}_{/f})_n := \mathrm{Hom}_f(\Delta^n \star S, \mathcal{C}).$ <p>([Lur09], Prop 1.2.9.2)</p>	<p>If <math>S = \Delta^0</math>, writing <math>C \in \mathcal{C}</math> for the object picked out by <math>f</math>, we have <math>(\mathcal{C}_{/C})_n = \mathrm{Hom}_C(\Delta^n \star \Delta^0, \mathcal{C}) \cong \mathrm{Hom}_C(\Delta^{n+1}, \mathcal{C})</math> (where the subscript indicates that we only consider morphisms sending the <math>(n+1)</math>st vertex to <math>C</math>). In other words, the objects are maps to <math>C</math>, the morphisms are commuting triangles over <math>C</math>, and so on; these are exactly the objects and morphisms in the 1-categorical case.</p>
Presentable category	[todo]	[todo]	[todo]
Presheaf	[todo]	[todo]	[todo]
Representable functor	[todo]	[todo]	[todo]
Right cone	$\mathcal{C}^{\triangleright} := \mathcal{C} \star [0]$ .	$\mathcal{C}^{\triangleright} := \mathcal{C} \star \Delta^0$ . ([Lur09], Not 1.2.8.4)	$\mathcal{C}$ with extra vertex (cone point) added, as well as a map from every other vertex in $\mathcal{C}$ to that cone point (plus obligatory degenerate simplices).
Right dualisable object	Object $C \in \mathcal{C}$ such that there exists some $*C \in \mathcal{C}$ and maps $e : *C \otimes C \rightarrow \mathbb{1}$ , $c : \mathbb{1} \rightarrow C \otimes *C$ such that the composites $C \rightarrow C \otimes *C \otimes C \rightarrow C$ and $*C \rightarrow *C \otimes C \otimes *C \rightarrow *C$ are equal to the identity.	Object $C \in \mathcal{C}$ such that there exists some $*C \in \mathcal{C}$ and maps $e : *C \otimes C \rightarrow \mathbb{1}$ , $c : \mathbb{1} \rightarrow C \otimes *C$ such that the composites $C \rightarrow C \otimes *C \otimes C \rightarrow C$ and $*C \rightarrow *C \otimes C \otimes *C \rightarrow *C$ are homotopic to the identity.	$C$ has a right dual up to homotopy.
Small category	$\mathcal{C}$ has a set's worth of objects, and between any two objects, a set's worth of morphisms.	$\mathcal{C}$ has a set's worth of nondegenerate simplices. <sup>9</sup> [Lur25, Section 03PP]	-

<sup>9</sup>In other words,  $\mathcal{C}$  is a simplicial set. So, by our definition, “ $\infty$ -category” means the same as “small  $\infty$ -category”.

Subcategory	Subcategory $\mathcal{C}' \subseteq \mathcal{C}$ .	<p>Subsimplicial set <math>\mathcal{C}' \subseteq \mathcal{C}</math> arising as a pullback</p> $\begin{array}{ccc} \mathcal{C}' & \longrightarrow & \mathcal{C} \\ \downarrow \lrcorner & & \downarrow \\ N(\mathbf{h}\mathcal{C})' & \longrightarrow & N(\mathbf{h}\mathcal{C}) \end{array} \quad \text{where } (\mathbf{h}\mathcal{C})' \subseteq \mathbf{h}\mathcal{C} \text{ is a subcategory. ([Lur09], 1.2.11)}$	<p>Expected definition: Subsimplicial set <math>\mathcal{C}' \subseteq \mathcal{C}</math> satisfying inner horn filling. These are actually equivalent: suppose we have a such a subsimplicial set. Then</p> $\begin{array}{ccc} \mathcal{C}' & \longrightarrow & \mathcal{C} \\ \downarrow \lrcorner & & \downarrow \\ N(\mathbf{h}\mathcal{C})' & \longrightarrow & N(\mathbf{h}\mathcal{C}) \end{array} \quad \text{is a pullback. Conversely, suppose we're given a pullback as in the box to the left. Then the diagram}$ $\begin{array}{ccccc} \Delta^n & \xleftrightarrow{\quad} & \Lambda_i^n & & \\ & \searrow \text{dotted} & \downarrow & \searrow & \\ & & \mathcal{C}' & \xrightarrow{\quad} & \mathcal{C} \\ & \searrow \text{dotted} & \downarrow \lrcorner & & \downarrow \\ & & N(\mathbf{h}\mathcal{C})' & \longrightarrow & N(\mathbf{h}\mathcal{C}) \end{array}$ <p>shows that <math>\mathcal{C}'</math> also satisfies inner horn filling, where the outer two dotted maps come from horn filling for <math>\mathcal{C}</math> and <math>N(\mathbf{h}\mathcal{C})'</math>, and the inner dotted map comes from the fact that the square is a pullback.</p>
Symmetric monoidal category	Opfibration of categories $p : \mathcal{C}^\otimes \rightarrow \mathbf{Fin}_*$ such that, for each $n \geq 0$ , the associated functors $\mathcal{C}_{\langle n \rangle}^\otimes \rightarrow \mathcal{C}_{\{*,i\}}^\otimes$ determine an equivalence of categories $\mathcal{C}_{\langle n \rangle}^\otimes \rightarrow \mathcal{C}_{\{*,1\}}^\otimes \times \dots \times \mathcal{C}_{\{*,n\}}^\otimes \simeq (\mathcal{C}_{\langle 1 \rangle}^\otimes)^n$ .	<p>Cocartesian fibration of simplicial sets <math>p : \mathcal{C}^\otimes \rightarrow N(\mathbf{Fin}_*)</math> such that, for each <math>n \geq 0</math>, the associated functors <math>\mathcal{C}_{\langle 1 \rangle}^\otimes \rightarrow \mathcal{C}_{\{*,i\}}^\otimes</math> determine an equivalence<sup>10</sup> of <math>\infty</math>-categories</p> $\mathcal{C}_{\langle n \rangle}^\otimes \rightarrow \mathcal{C}_{\{*,1\}}^\otimes \times \dots \times \mathcal{C}_{\{*,n\}}^\otimes \simeq (\mathcal{C}_{\langle 1 \rangle}^\otimes)^n$ <p>([Lur17], Def 2.0.0.7)</p>	<p>In the 1-categorical case, you recover the original category by setting <math>\mathcal{C} := \mathcal{C}_{\langle 1 \rangle}^\otimes</math>. The unit is <math>\mathcal{C}_{\langle 0 \rangle}^\otimes</math>, the tensor product <math>\otimes</math> is induced by <math>\langle 2 \rangle \rightarrow \langle 1 \rangle</math>, the symmetric isomorphism comes from the swapping map <math>\langle 2 \rangle \rightarrow \langle 2 \rangle</math>, the unitors and associators come from similar commutative diagrams to those in the intuition for a monoidal <math>\infty</math>-category. Conversely, given a symmetric monoidal category <math>\mathcal{C}</math>, define <math>\mathcal{C}^\otimes</math> to have objects finite sequences <math>[C_1, \dots, C_n]</math> of objects of <math>\mathcal{C}</math>, and a morphism <math>[C_1, \dots, C_n] \rightarrow [C'_1, \dots, C'_m]</math> to be a map <math>\langle n \rangle \rightarrow \langle m \rangle</math> and a collection of morphisms <math>\bigotimes_{i \in \alpha^{-1}\{j\}} C_i \rightarrow C'_j</math> for <math>1 \leq j \leq m</math>. Then the forgetful functor to <math>\mathbf{Fin}_*</math> is the required cocartesian fibration. ([Lur17], p165-168)</p>
Symmetric monoidal functor	[todo]	<p>Let <math>p : \mathcal{C}^\otimes \rightarrow N(\mathbf{Fin}_*)</math>, <math>q : \mathcal{D} \rightarrow N(\mathbf{Fin}_*)</math> be symmetric monoidal <math>\infty</math>-categories. Then an <math>\infty</math>-operad map <math>f \in \mathbf{Alg}_{\mathcal{C}}(\mathcal{D})</math> is a symmetric monoidal functor if it carries <math>p</math>-cocartesian morphisms to <math>q</math>-cocartesian morphisms. ([Lur17], Def 2.1.3.7)</p>	[todo]
Topos	[todo]	[todo]	[todo]

<sup>10</sup>Weak or categorical?

Undercategory	<p>For <math>C \in \mathcal{C}</math>, the category <math>\mathcal{C}_{C/}</math> satisfies the following universal property: for any category <math>\mathcal{D}</math>, there is a bijection</p> $\mathrm{Hom}(\mathcal{D}, \mathcal{C}_{C/}) \simeq \mathrm{Hom}_C([0] \star \mathcal{D}, \mathcal{C}),$ <p>where the subscript on the right indicates that we consider only those functors <math>[0] \star \mathcal{D} \rightarrow \mathcal{C}</math> whose restriction to <math>[0]</math> coincides with <math>C</math>. ([Lur09], 1.2.9)</p>	<p>For <math>f : S \rightarrow \mathcal{C}</math>, <math>S</math> a simplicial set and <math>\mathcal{C}</math> an <math>\infty</math>-category, the <math>\infty</math>-category <math>\mathcal{C}_{f/}</math> satisfies the following universal property: for any simplicial set <math>X</math>, there is a bijection</p> $\mathrm{Hom}(X, \mathcal{C}_{f/}) \simeq \mathrm{Hom}_f(S \star X, \mathcal{C}),$ <p>where the subscript on the right indicates that we consider only those functors <math>S \star X \rightarrow \mathcal{C}</math> whose restriction to <math>S</math> coincides with <math>f</math>. Explicitly,</p> $(\mathcal{C}_{f/})_n := \mathrm{Hom}_f(S \star \Delta^n, \mathcal{C}).$ <p>([Lur09], Prop 1.2.9.2)</p>	<p>If <math>S = \Delta^0</math>, writing <math>C \in \mathcal{C}</math> for the object picked out by <math>f</math>, we have <math>(\mathcal{C}_{C/})_n = \mathrm{Hom}_C(\Delta^0 \star \Delta^n, \mathcal{C}) \cong \mathrm{Hom}_C(\Delta^{n+1}, \mathcal{C})</math> (where the subscript indicates that we only consider morphisms sending the 0th vertex to <math>C</math>). In other words, the objects are maps from <math>C</math>, the morphisms are commuting triangles under <math>C</math>, and so on; these are exactly the objects and morphisms in the 1-categorical case.</p>
---------------	--	---	---



Equivalences		
Name	Between	Definition
Strong equivalence	Topological categories $\mathcal{C}, \mathcal{D}$	$\mathcal{C} \rightarrow \mathcal{D}$ is an equivalence in the sense of enriched category theory. ([Lur09], Def 1.1.3.1)
(Weak) equivalence	Topological categories $\mathcal{C}, \mathcal{D}$	The induced functor $\mathbf{h}\mathcal{C} \rightarrow \mathbf{h}\mathcal{D}$ is an equivalence of $\mathcal{H}$ -enriched categories. ([Lur09], Def 1.1.3.6)
Categorical equivalence	Simplicial sets $X, S$	The induced functor $\mathbf{h}X \rightarrow \mathbf{h}S$ is an equivalence of $\mathcal{H}$ -enriched categories. ([Lur09], Def 1.1.5.14)
Weak (homotopy) equivalence	Simplicial sets $X, S$	The induced map $ X  \rightarrow  S $ is a weak homotopy equivalence of topological spaces. ([Lur09], 1.1.4)
Equivalence	Simplicial categories $\mathcal{C}, \mathcal{D}$	The induced functor $\mathbf{h}\mathcal{C} \rightarrow \mathbf{h}\mathcal{D}$ is an equivalence of $\mathcal{H}$ -enriched categories. ([Lur09], Def 1.1.4.4)

Fibrations and anodyne morphisms		
Name	Describes	Definition
Acyclic Kan fibration	$f : X \rightarrow S$ map of simplicial sets	see: trivial Kan fibration. ([nLa23])
Anodyne	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>p : Y \rightarrow T</math> a Kan fibration,</p> $\begin{array}{ccc} X & \longrightarrow & Y \\ f \downarrow & \nearrow & \downarrow p \\ S & \longrightarrow & T \end{array}$ <p>there exists a dotted lift. ([Lur09], Ex 2.0.0.1)</p>
Cartesian fibration	$f : X \rightarrow S$ map of simplicial sets	$f$ is an inner fibration such that for every edge $g : x \rightarrow y$ of $S$ and every vertex $\tilde{y}$ of $X$ with $f(\tilde{y}) = y$ , there exists an $f$ -cartesian edge $\tilde{g} : \tilde{x} \rightarrow \tilde{y}$ with $f(\tilde{g}) = g$ . ([Lur09], Def 2.4.2.1)
Categorical fibration	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>p : Y \rightarrow T</math> both a cofibration and a categorical equivalence,</p> $\begin{array}{ccc} Y & \longrightarrow & X \\ p \downarrow & \nearrow & \downarrow f \\ T & \longrightarrow & S \end{array}$ <p>there exists a dotted lift. ([Lur09], p90)</p>
Cocartesian fibration	$f : X \rightarrow S$ map of simplicial sets	$f$ is an inner fibration such that for every edge $g : x \rightarrow y$ of $S$ and every vertex $\tilde{x}$ of $X$ with $f(\tilde{x}) = x$ , there exists an $f$ -cocartesian edge $\tilde{g} : \tilde{x} \rightarrow \tilde{y}$ with $f(\tilde{g}) = g$ . ([Lur09], Def 2.4.2.1)
Cofibration	$f : X \rightarrow S$ map of simplicial sets	$f$ is a monomorphism. ([Lur09], A.2.7)
Inner anodyne	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>p : Y \rightarrow T</math> an inner fibration,</p> $\begin{array}{ccc} X & \longrightarrow & Y \\ f \downarrow & \nearrow & \downarrow p \\ S & \longrightarrow & T \end{array}$ <p>there exists a dotted lift. ([Lur09], Def 2.0.0.3)</p>

Inner fibration	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>0 &lt; i &lt; n</math>,</p> $\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \Delta^n & \longrightarrow & S \end{array}$ <p>there exists a dotted lift.</p>
Isofibration	$F : \mathcal{C} \rightarrow \mathcal{D}$ map of $\infty$ -categories	<p><math>F</math> is an inner fibration such that for all <math>C \in \mathcal{C}</math> and every isomorphism <math>u : D \rightarrow FC</math> in <math>\mathcal{D}</math> (i.e. <math>[u]</math> is an isomorphism in <math>\mathbf{h}\mathcal{D}</math>) there exists an isomorphism <math>\bar{u} : \bar{D} \rightarrow C</math> in <math>\mathcal{C}</math> such that <math>F(\bar{u}) = u</math>. [Lur25, Def 01EN]</p>
(Kan) fibration	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>0 \leq i \leq n</math>,</p> $\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \Delta^n & \longrightarrow & S \end{array}$ <p>there exists a dotted lift. ([Lur09], A.2.7)</p>
Left anodyne	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>p : Y \rightarrow T</math> a left fibration,</p> $\begin{array}{ccc} X & \longrightarrow & Y \\ f \downarrow & \nearrow & \downarrow p \\ S & \longrightarrow & T \end{array}$ <p>there exists a dotted lift. ([Lur09], Def 2.0.0.3)</p>
Left fibration	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>0 \leq i &lt; n</math>,</p> $\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \Delta^n & \longrightarrow & S \end{array}$ <p>there exists a dotted lift. ([Lur09], Def 2.0.0.3)</p>
Right anodyne	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>p : Y \rightarrow T</math> a right fibration,</p> $\begin{array}{ccc} X & \longrightarrow & Y \\ f \downarrow & \nearrow & \downarrow p \\ S & \longrightarrow & T \end{array}$ <p>there exists a dotted lift. ([Lur09], Def 2.0.0.3)</p>
Right fibration	$f : X \rightarrow S$ map of simplicial sets	<p>For every solid arrow diagram as below, with <math>0 &lt; i \leq n</math>,</p> $\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \Delta^n & \longrightarrow & S \end{array}$ <p>there exists a dotted lift. ([Lur09], Def 2.0.0.3)</p>

Serre fibration	$f : Y \rightarrow Z$ map of topological spaces	For every solid arrow diagram as below, <div style="text-align: center;"> <math display="block">\begin{array}{ccc} \{0\} \times  \Delta^n  &amp; \longrightarrow &amp; Y \\ \downarrow &amp; \nearrow &amp; \downarrow f \\ [0, 1] \times  \Delta^n  &amp; \longrightarrow &amp; Z \end{array}</math> </div> there exists a dotted lift. [Lur25, Def 021R]
Trivial (Kan) fibration	$f : X \rightarrow S$ map of simplicial sets	For every solid arrow diagram as below, <div style="text-align: center;"> <math display="block">\begin{array}{ccc} \partial\Delta^n &amp; \longrightarrow &amp; X \\ \downarrow &amp; \nearrow &amp; \downarrow f \\ \Delta^n &amp; \longrightarrow &amp; S \end{array}</math> </div> there exists a dotted lift. ([Lur25, Def 006W]/[Lur09], Def 2.0.0.2)

Nerves		
Name	Domain object	Definition
Nerve	Category $\mathcal{C}$	$(N\mathcal{C})_n = \{n\text{-composable strings of morphisms in } \mathcal{C}\}$ . ([Lur09], p9)
Simplicial nerve	Simplicial category $\mathcal{C}$	$(N\mathcal{C})_n = \text{Hom}_{\mathbf{Cat}_\Delta}(\mathfrak{C}[\Delta^n], \mathcal{C})$ , where $\mathfrak{C}[\Delta^n]$ is the category whose objects are the same as $[n]$ , and $\text{Hom}_{\mathfrak{C}[\Delta^n]}(i, j) = \emptyset$ for $i < j$ and $N(P_{ij})$ for $i \geq j$ (where $P_{ij} = \{I \subseteq [n] : (i, j \in I) \wedge (\forall k \in I, i \leq k \leq j)\}$ ). ([Lur09], Def 1.1.5.5)
Topological nerve	Topological category $\mathcal{C}$	The simplicial nerve of $\text{Sing } \mathcal{C}$ . ([Lur09], Def 1.1.5.5)

Homotopy categories	
Domain object	Definition
$\infty$ -Category $\mathcal{C}$	The objects of $\text{h}\mathcal{C}$ are the vertices of $\mathcal{C}$ , and $\text{Hom}_{\text{h}\mathcal{C}}(X, Y)$ is the set of homotopy classes of edges $X \rightarrow Y$ in $\mathcal{C}$ . ([Lur09], Prop 1.2.3.9)
Simplicial category $\mathcal{C}$	$\text{h} \mathcal{C} $ . ([Lur09], 1.1.4)
Topological category $\mathcal{C}$	$\text{h}\mathcal{C}$ has the same objects as $\mathcal{C}$ , and $\text{Hom}_{\text{h}\mathcal{C}}(X, Y) = [\text{Hom}_{\mathcal{C}}(X, Y)]$ . ([Lur09], 1.1.3)

Objects	
Name	Definition
$\mathbf{Alg}_{\mathcal{C}}(\mathcal{D})$ ( $p : \mathcal{C}^\otimes \rightarrow N(\mathbf{Fin}_*)$ , $q : \mathcal{D}^\otimes \rightarrow N(\mathbf{Fin}_*)$ $\infty$ -operads)	The full subcategory of $\mathbf{Fun}_{N(\mathbf{Fin}_*)}(\mathcal{C}^\otimes, \mathcal{D}^\otimes)$ <sup>11</sup> spanned by the $\infty$ -operad maps. ([Lur17], Def 2.1.2.7)

<sup>11</sup>This notation is defined nowhere I can see, but must just mean maps  $\mathcal{C}^\otimes \rightarrow \mathcal{D}^\otimes$  making the obvious triangle commute (but is this commuting on the nose, or only up to homotopy?).

<b>Assoc</b> (the associative operad)	The coloured operad with a single object $\mathbf{a}$ , and for every finite set $I$ , $\text{Hom}(\{\mathbf{a}\}_I, \mathbf{a})$ is the set of linear orderings on $I$ . Given a map of finite sets $\alpha : I \rightarrow J$ together with operations $\phi_j \in \text{Hom}(\{\mathbf{a}\}_{\alpha(i)=j}, \mathbf{a})$ and $\psi \in \text{Hom}(\{\mathbf{a}_j, \mathbf{a}\})$ , we identify each $\phi_j$ with a linear ordering $\leq_j$ on the set $\alpha^{-1}\{j\}$ and $\psi$ with a linear ordering $\leq'$ on the set $J$ . The composition of $\psi$ with $\{\phi_j\}$ corresponds to the linear ordering $\leq$ on the set $I$ which is defined by: $i \leq i'$ if either $\alpha(i) <_j \alpha(i')$ or $\alpha(i) = j = \alpha(i')$ and $i \leq_j i'$ . ([Lur17], Def 4.1.1.1)
$\text{Assoc}^\otimes$ (the associative $\infty$ -operad)	$N(\mathbf{Assoc}^\otimes)$ . ([Lur17], Def 4.1.1.3)
$\mathbf{Assoc}^\otimes$	The category whose objects are the objects of $\mathbf{Fin}_*$ , and a morphism $m \rightarrow n$ is given by a map $\alpha : \langle m \rangle \rightarrow \langle n \rangle$ in $\mathbf{Fin}_*$ together with a collection of linear orderings $\leq_j$ on $\alpha^{-1}\{j\}$ , for $1 \leq j \leq n$ . Composition of morphisms is determined by the composition laws on $\mathbf{Fin}_*$ and on $\mathbf{Assoc}$ . [Lur17], Def 4.1.1.3
$\mathbf{Fin}_*$	The category whose objects are the sets $\langle n \rangle = \{*, 1, 2, \dots, n\}$ , and a morphism $\langle m \rangle \rightarrow \langle n \rangle$ is a map $\alpha : \langle m \rangle \rightarrow \langle n \rangle$ such that $\alpha(*) = *$ .
<b>Kan</b>	The full subcategory of $\mathbf{sSet}$ spanned by the collection of small Kan complexes. ([Lur09], Def 1.2.16.1)
<b>KAN</b>	The category of all Kan complexes. ([Lur09], Rem 5.1.6.1)
$\mathcal{S}$ (the $\infty$ -category of spaces)	The simplicial <sup>12</sup> nerve $N(\mathbf{Kan})$ . ([Lur09], Def 1.2.16.1)
$\widehat{\mathcal{S}}$	The simplicial nerve $N(\mathbf{KAN})$ . ([Lur09], Rem 5.1.6.1)

<sup>12</sup> $\mathbf{sSet}$  is a simplicial category, with  $\text{Hom}(X, S)_n = \text{Hom}_{\mathbf{sSet}}(\Delta^n \times X, S)$ . The subcategory  $\mathbf{Kan}$  inherits this structure.

## References

- [Eti+15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor Categories*. 2015.
- [Lur07] Jacob Lurie. *Derived Algebraic Geometry II: Noncommutative Algebra*. 2007. arXiv: [math/0702299](https://arxiv.org/abs/math/0702299) [[math.CT](https://arxiv.org/abs/math/0702299)]. URL: <https://arxiv.org/abs/math/0702299>.
- [Lur09] Jacob Lurie. *Higher Topos Theory*. 2009.
- [Lur17] Jacob Lurie. *Higher Algebra*. 2017. URL: <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Lur25] Jacob Lurie. *Kerodon*. <https://kerodon.net>. 2025.
- [nLa23] nLab (Urs Schreiber). *acyclic Kan fibration*. <https://ncatlab.org/nlab/show/acyclic+Kan+fibration>. Revision 5. 2023.
- [nLa25a] nLab authors. *accessible category*. <https://ncatlab.org/nlab/show/accessible+category>. Revision 69. 2025.
- [nLa25b] nLab authors. *Cartesian morphism*. <https://ncatlab.org/nlab/show/Cartesian+morphism>. Revision 52. 2025.
- [nLa25c] nLab authors. *join of simplicial sets*. <https://ncatlab.org/nlab/show/join+of+simplicial+sets>. Revision 62. 2025.
- [Rie16] Emily Riehl. *Category Theory in Context*. 2016. URL: <https://emilyriehl.github.io/files/context.pdf>.