



## FERTILIZER RECOMMENDATION

**Customer:** Ronie  
**Address:** Lagangilang  
**Analysis Requested by:** LGU Lagangilang  
**Sample Submitted by:** LGU Lagangilang

**Date Received:** October 20, 2023  
**Date Analyzed:** 11/17/2023 to 12/28/2023  
**Date Reported:** May 1, 2026  
**Sample Condition:** Wet

**TEST REQUEST NO.:** RSL-2023-181  
**SAMPLE IDENTIFICATION:** -  
**LABORATORY CODE:** FM-LG23-1

| BEFORE PLANTING: TIMING OF ORGANIC AND INORGANIC FERTILIZER APPLICATIONS |                                                                          |              |              | TARGET YIELD                         |                                                  |                     |
|--------------------------------------------------------------------------|--------------------------------------------------------------------------|--------------|--------------|--------------------------------------|--------------------------------------------------|---------------------|
| TIMING                                                                   | MATURITY                                                                 |              |              | Combo 1<br>3-4 tons                  | Combo 2<br>5-6 tons                              | Combo 3<br>7-8 tons |
|                                                                          | 100-110 Days                                                             | 111-120 Days | 121-130 Days |                                      |                                                  |                     |
| 30 Days before transplanting                                             | Rice Straw: Spread in the rice field or use in the production of compost |              |              | 3- 4 tons rice straw                 | 5-6 tons rice straw                              | 7-8 tons rice straw |
| 14 days before transplanting                                             | Organic: Chicken Manure, Commercial Organic fertilizer, o Vermicompost   |              |              | 10 bags                              | 10 bags                                          | 10 bags             |
| Seeding<br>7-10 days after seeding                                       | N/A                                                                      | N/A          | N/A          | N/A                                  | Apply in the seedbed<br>1-2 kg ZnSO <sub>4</sub> |                     |
| Seeding<br>10-14 days after seeding                                      | N/A                                                                      | N/A          | N/A          | Apply in the seedbed 2-4 kg 14-14-14 |                                                  |                     |

| NUTRIENT REQUIREMENT AND RECOMMENDATION FOR HYBRID RICE                                                      |                                                            |                                                                                                                                                                     |    |    |
|--------------------------------------------------------------------------------------------------------------|------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|----|
| Nutrient Requirement, kg/ha                                                                                  |                                                            | N                                                                                                                                                                   | P  | K  |
| (Wet season)                                                                                                 |                                                            | 90                                                                                                                                                                  | 70 | 50 |
| (Dry season)                                                                                                 |                                                            | 100                                                                                                                                                                 | 70 | 50 |
| Fertilizer Requirement (kg/ha)                                                                               |                                                            | Time and Method of Application                                                                                                                                      |    |    |
| Wet Season                                                                                                   | Dry Season                                                 |                                                                                                                                                                     |    |    |
| **Basal: 10-20 bags organic fertilizer. Apply the recommended organic fertilizer during the last harrowing** |                                                            |                                                                                                                                                                     |    |    |
| 1st Top dress:<br>14-14-14= 357 kg/ha<br>0-20-0= 100 kg/ha                                                   | 1st Top dress:<br>14-14-14= 357 kg/ha<br>0-20-0= 100 kg/ha | Broadcast 1/2 of the nitrogenous fertilizer with phosphorus and potash<br>0-14 days after transplanting/ 10-14 days after seeding.                                  |    |    |
| ZnSO4                                                                                                        | 5 kg/ha                                                    | Apply within 15 days after transplanting; can be sprayed or either mixed<br>with fertilizer during top dressing.                                                    |    |    |
| 2nd Top Dress:<br>46-0-0= 43 kg/ha                                                                           | 2nd Top Dress:<br>46-0-0= 54 kg/ha                         | Topdress with remaining 1/4 of the nitrogen 22-26 days after<br>transplanting/ 32-36 days after seeding.                                                            |    |    |
| 3rd Top Dress:<br>46-0-0= 43 kg/ha                                                                           | 3rd Top Dress:<br>46-0-0= 54 kg/ha                         | Topdress with remaining 1/4of the nitrogen 32-36 days after<br>transplanting/ 48-52 days after seeding; add remaining potash from<br>first top dress if there's any |    |    |

Fertilizer recommendation based on BFS Recommendation and Protocol for Rice -2022 (Bureau of Soils and Water Management)

**Notes:** For further concerns to the test report, you may email us at [rsldacar@gmail.com](mailto:rsldacar@gmail.com) or contact **0912-877-3603**.

Fertilizer Recommendation by:

Reviewed by:

Noted by:

**JEROME S. LUCYAO**  
Agriculturist I  
Registration No. 26242  
Validity Date: 10/25/2026

**HAILY G. PATNA-AN**  
Agriculturist II  
Registration No. 9877  
Validity Date: 02/12/2025

**ALONA JEAN I. ASTRERA**  
Chemist III