

Operative Report

Dana Keller | MRN DEMO-BC-001 | DOB 1974-02-11 | Female

Operation date	2026-08-07
Preoperative diagnosis	Left-breast invasive carcinoma after neoadjuvant systemic therapy
Postoperative diagnosis	Same
Procedures	Left image-guided partial mastectomy; sentinel lymph-node biopsy; targeted excision of clipped left axillary node
Surgeon	Alex Morgan, MD
Anesthesia	General

Operative findings

The localized lesion in the upper outer left breast was excised with the localization marker and biopsy clip contained within the specimen. Three sentinel/targeted axillary nodes were removed, including the previously clipped node. No grossly abnormal additional axillary nodes were identified.

Procedure

A curvilinear incision was made over the localized breast lesion. A partial mastectomy specimen was removed with orienting sutures and sent fresh to pathology. Additional cavity margins were obtained. Through a separate axillary incision, sentinel lymphatic mapping was performed and three nodes were excised, including the clipped target node. Hemostasis was achieved and both incisions were closed in layers.

Estimated blood loss / complications

Estimated blood loss: 20 mL. No intraoperative complication. Patient transferred to recovery in stable condition.

Specimens submitted

Left partial mastectomy specimen; additional superior, inferior, medial, lateral, anterior and posterior margins; three left axillary sentinel/targeted lymph nodes.