

STAT 516: Basic Probability and its Applications

Lecture 3: Conditional Probability and Independence

Prof. Michael Levine

January 21, 2024

Motivating Example

$S = \{11, 21 \dots 56, 66\}$ = Total of 6^2 possible outcomes
 $\frac{1}{36}$

- ▶ Experiment ξ consists of rolling a fair die twice;
 $A = \{ \text{the first roll is 6} \}$ and $P(A) = \frac{6}{36} = \frac{1}{6}$
 $B = \{ \text{the sum of the two rolls is 12} \}$. $P(B) = \frac{1}{36}$
- ▶ Under the assumption of the equally likely sample points,
 $P(B) = \frac{1}{36}$ *only one pair (6,6) gives sum of 12.*
- ▶ If we know that A already happened, intuitively we feel that the probability of the event B , given A , should be $\frac{1}{6}$ *If already know value of first die Ex: 6*
- ▶ We say that $P(B|A) = \frac{1}{6}$ and call it the **conditional probability of A** *only 6 possible outcome: {61, 62, 63, 64, 65, 66} $\frac{1}{6}$*
- ▶ The conditional probability tells us *among the times that A already happened how often B also happens*

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{\frac{1}{36}}{\frac{1}{6}} = \frac{1}{6}$$

Formal definition of the conditional probability

- ▶ Conditional probability results from restricting the outcomes to only those that are *inside* C_1 where $P(C_1 > 0)$
- ▶ Therefore, instead of

$$P(C_2) = \frac{N(C_2)}{N(S)}$$

we have

$$\frac{N(C_2 \cap C_1)}{N(S \cap C_1)} = \frac{P(C_2 \cap C_1)}{P(S \cap C_1)} = \frac{P(C_2 \cap C_1)}{P(C_1)}$$

- ▶ Thus, the definition:

$$P(C_2|C_1) = \frac{P(C_2 \cap C_1)}{P(C_1)}$$

Example I

- ▶ Consider the situation where of all individuals buying a digital camera 60% include an optional memory card, 40% - an extra battery and 30% - both. What is the probability that a person who buys an extra battery also buys a memory card?
- ▶ Let $A = \{\text{memory card purchased}\}$ and $B = \{\text{battery purchased}\}$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.3}{0.4} = 0.75$$

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{0.3}{0.6} = 0.50$$

Multiplication Rule I

- ▶ A simple **multiplication rule** follows from the definition of **conditional probability**
- ▶ For any two events C_1 and C_2 such that $P(C_1) > 0$

$$P(C_1 \cap C_2) = P(C_1)P(C_2|C_1)$$

- ▶ For example, take a bowl with eight chips. 3 of the chips are red and the remaining 5 are blue. 2 chips are drawn without replacement.
- ▶ What is the probability that the 1st draw results in a red chip (C_1) and the second draw results in a blue chip (C_2)

Multiplication rule II

- ▶ It is reasonable to assume that $P(C_1) = \frac{3}{8}$ and $P(C_2|C_1) = \frac{5}{7}$
- ▶ Then, $P(C_1 \cap C_2) = \frac{3}{8} \frac{5}{7} = \frac{15}{56} = 0.2679$

Multiplication rule III

- ▶ The multiplication rule can be extended to three or more events
- ▶ Using the basic multiplication rule iteratively, and assuming $P(C_1 \cap C_2) > 0$, we have

$$P(C_1 \cap C_2 \cap C_3) = P(C_1)P(C_2|C_1)P(C_3|C_1 \cap C_2)$$

Multiplication rule IV


- ▶ Four cards are dealt successively from an ordinary deck of playing cards
- ▶ Dealing is *at random* and *without replacement*
- ▶ The probability of receiving a spade, a heart, a diamond, and a club, in that order, is

$$\frac{13}{52} \frac{13}{51} \frac{13}{50} \frac{13}{49} = 0.0044$$

♣ ♥ ♦ ♠ → different sets, not interfering with one another.

↪ ↪ ↪ one less choice after previous card is picked

The Law of Total Probability



- ▶ Let $C_1, \dots, C_k$ form a **partition** of the sample space S
- ▶ This means that $C_i \cap C_j = \emptyset$ for all $i \neq j$ and $\bigcup_{i=1}^k C_i = S$
Mutually Exclusive *Partition*
- ▶ Moreover, let $0 < P(C_i) < 1$, $i = 1, 2, \dots, k$
- ▶ Then, the **Law of Total Probability** is

$$P(C) = \sum_{i=1}^k P(C_i)P(C|C_i)$$

Example

- ▶ One of the cards from a deck of 52 cards is missing from the deck but we don't know which one
- ▶ One card is chosen at random from the remaining 51 cards. What is the probability that it is a spade?
- ▶ Let $A = \text{the missing card is a spade}$ and $B = \text{the card chosen from the remaining ones is a spade}$ *B depends on what A is $\Rightarrow$ conditional*
- ▶ Then,

$$P(B) = P(B|A)P(A) + P(B|A')P(A') = \frac{12}{51} \frac{1}{4} + \frac{13}{51} \frac{3}{4} = \frac{1}{4}$$

- ▶ Note that it is impossible to have 12.5 spade cards in the remaining deck of 51 cards...but $P(B) = \frac{1}{4}$ nevertheless

Example

- ▶ A certain item is produced in a factory on one of the three machines A , B , or C . The percentages of items produced on A , B , and C , are 50%, 30%, and 20%, respectively
- ▶ 4%, 2%, and 4% of their products, respectively, are defective. What is the percentage of all copies of this item that are defective?
- ▶ Let the event that an item is produced by A , B , or C , respectively, be A_1 , A_2 , and A_3 ; let D be the event that it is a defective item



$$P(D) = \sum_{i=1}^3 P(D|A_i)P(A_i) = .04 \times .5 + .02 \times .3 + .04 \times .2 = .034$$

A more complicated example

$$\text{Player 1 } P_1 = \frac{\text{winning ticket}}{\text{ticket}} = \frac{m}{n}$$

- ▶ A lottery with n tickets, out of which m prespecified tickets will win a prize
- ▶ There are n players who choose one ticket at random successively from available tickets
- ▶ What is a probability that the i th player wins a prize?

$$\text{Player 2 } P_2 = \underbrace{\left(\frac{m-1}{n-1} \right) \left(\frac{m}{n} \right)}_{\text{winning ticket by Player 1}} + \underbrace{\left(\frac{n-m}{n-m-1} \right) \left(\frac{m}{n-1} \right)}_{\text{not winning ticket by Player 1}}$$

p. of winning
p. of not winning
 $(1 - \frac{m}{n})$: complement of winning

- Remaining winning tickets
- new total since player 1 took a ticket
- No winning ticket lost

A more complicated example

- ▶ For the first player, the probability is $p_1 = \frac{m}{n}$ ✓
- ▶ By total probability formula,

$$p_2 = \frac{m-1}{n-1} \times \frac{m}{n} + \frac{m}{n-1} \times \left(1 - \frac{m}{n}\right) = \frac{m}{n}$$

Case 1: P1 win (lost a winning ticket) $\times$ *winning prob* $\frac{m}{n}$
Case 2: P1 lost (no winning ticket lost) $\times$ *complement of winning prob aka not winning prob.* $\left(1 - \frac{m}{n}\right)$
lost a total ticket $\frac{m-1}{n-1}$ *still lost a total ticket* $\frac{m}{n-1}$

- ▶ Again, by total probability formula

$$\begin{aligned}
 p_3 &= \frac{m}{n} \times \frac{m-1}{n-1} \times \frac{m-2}{n-2} + \left(1 - \frac{m}{n}\right) \times \frac{m}{n-1} \times \frac{m-1}{n-2} \\
 &\quad + \frac{m}{n} \times \left(1 - \frac{m-1}{n-1}\right) \times \frac{m-1}{n-2} \\
 &\quad + \left(1 - \frac{m}{n}\right) \times \left(1 - \frac{m}{n-1}\right) \times \frac{m}{n-2} \\
 &= \frac{m}{n}
 \end{aligned}$$

Case 1: Both player 1 and 2 win $\frac{m}{n} \times \frac{m-1}{n-1} \times \frac{m-2}{n-2}$
Case 2: P1 win, P2 lost $\left(1 - \frac{m}{n}\right) \times \frac{m}{n-1} \times \frac{m-1}{n-2}$
Case 3: P1 lost, P2 win $\frac{m}{n} \times \left(1 - \frac{m-1}{n-1}\right) \times \frac{m-1}{n-2}$
Case 4: Both Player 1 and 2 lost $\left(1 - \frac{m}{n}\right) \times \left(1 - \frac{m}{n-1}\right) \times \frac{m}{n-2}$

- ▶ It can be shown by induction that for any $i \geq 1$ we have $p_i = \frac{m}{n}$ - no early entry advantage for this lottery!

Bayes Theorem

- ▶ Also, the **Bayes theorem** follows directly:

$$P(C_j|C) = \frac{P(C \cap C_j)}{P(C)} = \frac{P(C_j)P(C|C_j)}{\sum_{i=1}^k P(C_i)P(C|C_i)}$$

A simple example I

- ▶ A store stocks light bulbs from three suppliers. Suppliers A, B, and C supply 10%, 20%, and 70% of the bulbs respectively.
- ▶ It has been determined that company A's bulbs are 1% defective while company B's are 3% defective and company C's are 4% defective.
- ▶ If a bulb is selected at random and found to be defective, what is the probability that it came from supplier B?

A simple example II

- ▶ Let $D =$ defective lightbulb

$$\begin{aligned} P(B|D) &= \frac{P(B)P(D|B)}{P(A)P(D|A) + P(B)P(D|B) + P(C)P(D|C)} \\ &= \frac{0.2(0.03)}{0.1(0.01) + 0.2(0.03) + 0.7(0.04)} \approx 0.1714 \end{aligned}$$

Independence

- ▶ The first definition

$$P(C_2 \cap C_1) = P(C_1)P(C_2) \quad (1)$$

- ▶ An alternative definition follows from multiplication formula:

$$P(C_1|C_2) = P(C_1) = \frac{P(C_1 \cap C_2)}{P(C_2)} \quad (2)$$

- ▶ Note that the first definition also allows us to consider the case where $P(C_1) = P(C_2) = 0$

Some immediate consequences of the definition of independence

- $P(A|B) = P(A)$
- $P(A|B) = \frac{P(A \cap B)}{P(B)}$
- $P(A \cap B) = P(A) \cdot P(B)$

► The independence of A and B implies that

1. A and B' are independent $P(A|B') = P(A)$
2. A' and B are independent $P(A'|B) = P(A')$
3. A' and B' are independent $P(A' \cap B') = P(A')P(B')$

Example I

- ▶ Consider the fair six-sided die. Define $A = \{2, 4, 6\}$, $B = \{1, 2, 3\}$ and $C = \{1, 2, 3, 4\}$.
- ▶ Clearly, $P(A) = \frac{1}{2}$, $P(A|B) = \frac{1}{3}$ and $P(A|C) = \frac{1}{2}$.
- ▶ Therefore, A and C are independent but A and B are NOT!

Knowing C doesn't change A , it's still $1/2$.
Therefore A and C are independent

Example II

- ▶ A red and a white die are cast independently
- ▶ Let $C_1 = \{4 \text{ red die}\}$ and $C_2 = \{3 \text{ white die}\}$
- ▶ $P(C_1) = \frac{1}{6}$ and $P(C_2) = \frac{1}{6}$
- ▶ By independence,

$$P(C_1 \cap C_2) = \frac{1}{36}$$

- ▶ The probability that the sum of the up spots is equal to 7 is

$$P[(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)] = 6 * \left(\frac{1}{6}\right)^2 = \frac{1}{6}$$

Some additional consequences of the definition of independence

- ▶ If A and B are mutually exclusive events, they **cannot** be independent.
- ▶ If either $P(A) = 0$ or $P(B) = 0$, A and B are always independent:

$$0 \leq P(A \cap B) \leq \min(P(A), P(B)) = 0,$$

therefore $P(A \cap B) = 0$

Independence: practical considerations

- ▶ In practice, we do not verify independence of events. Instead, we ask ourselves whether independence is a property that we wish to incorporate into a mathematical model of an experiment, based on the common sense
- ▶ Thus, independence is commonly *assumed*
- ▶ Example: let C_2 =(A student is a female) and C_1 =(A student is concentrating in elementary education); clearly,
 $P(C_2|C_1) \neq P(C_2)$

Mutual independence

$$K=2: P(A_1 \cap A_2) = P(A_1)P(A_2)$$

$$P(A_1 \cap A_3) = P(A_1)P(A_3)$$

$$P(A_2 \cap A_3) = P(A_2)P(A_3)$$

$$K=3: P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2)P(A_3)$$

- ▶ $C_1, C_2, \dots, C_n$ are **mutually independent** if for every $2 \leq k \leq n$, and every possible subset of indices $i_1, i_2, \dots, i_k$

product of

$$P(C_{i_1} \cap C_{i_2} \cap \dots \cap C_{i_k}) = P(C_{i_1}) \cdot P(C_{i_2}) \cdot \dots \cdot P(C_{i_k})$$

- ▶ In particular, this means that

$$P(C_1 \cap C_2 \cap \dots \cap C_n) = P(C_1)P(C_2) \dots P(C_n)$$

The relationship between mutual independence and pairwise independence

- ▶ Pairwise independence DOES NOT imply mutual independence *need to verify both*
- ▶ Let a single outcome be ω_1 and the entire sample space is $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4\}$
- ▶ Then let $C_1 = \{\omega_1, \omega_2\}$, $C_2 = \{\omega_1, \omega_3\}$ and $C_3 = \{\omega_1, \omega_4\}$
- ▶ If each outcome is equally likely with $p = \frac{1}{4}$, it is easy to check that pairwise independence is valid *Ex): $P(A_1 \cap A_2) = P(A_1) \cdot P(A_2) = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$*
- ▶ However,

$$P(C_1 \cap C_2 \cap C_3) = \frac{1}{4} \neq \left(\frac{1}{2}\right)^3 = P(C_1)P(C_2)P(C_3)$$

*This is pairwise independence
but not mutual independence, as
high level is not satisfied.*

Example

- ▶ Consider sample space S of 36 ordered pairs (i, j) with $i, j = 1, \dots, 6$
- ▶ We assume that for each pair the probability of occurring $p = \frac{1}{36}$
- ▶ Let $C_1 = \{(i, j) : j = 1, 2 \text{ or } 5\}$, $C_2 = \{(i, j) : j = 4, 5 \text{ or } 6\}$ and $C_3 = \{(i, j) : i + j = 9\}$
- ▶ Then,

$$\text{Intersect at } j=5 \quad P(C_1 \cap C_2) = \frac{1}{6} \neq \frac{1}{4} = P(C_1)P(C_2)$$

$$\begin{matrix} j=5 \\ i=4 \end{matrix} \quad P(C_1 \cap C_3) = \frac{1}{36} \neq \frac{1}{18} = P(C_1)P(C_3)$$

and

$$\begin{matrix} j=4,5,6 \\ i=5,4,3 \end{matrix} \quad P(C_2 \cap C_3) = \frac{1}{12} \neq \frac{1}{18} = P(C_2)P(C_3)$$

- ▶ However,

$$P(C_1 \cap C_2 \cap C_3) = \frac{1}{36} = P(C_1)P(C_2)P(C_3)$$

This is mutual independence but not pairwise independence, as high level is not satisfied.

Independent experiments

- ▶ Events as outcome of **independent experiments**
- ▶ A computer system is built with K_1 , K_2 and K_3
- ▶ Each successive component is a backup for the previous one
- ▶ If probabilities of failure are $P(K_1) = 0.01$, $P(K_2) = 0.03$, and $P(K_3) = 0.08$, the system fails with probability

$$(0.01)(0.03)(0.08) = 24 \times 10^{-6}$$

- ▶ The system functions successfully with the probability

$$1 - 24 \times 10^{-6} = 0.999976$$