

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-01T10-43-16-281Z

Run ID: #2

Date Generated: 2026-01-01 10:43:21

Total Cells: 2000 | Model Mode: CLINICAL

1. Introduction

Cellular reprogramming holds significant promise for regenerative medicine by enabling the conversion of somatic cells into induced pluripotent stem cells (iPSCs). This experiment aimed to evaluate the efficacy of repeated OSKM transfections in facilitating somatic cell reprogramming under oncogenic stress conditions.

2. Materials and Methods

The experimental protocol involved six sequential transfections using the Yamanaka factors (OSKM: Oct4, Sox2, Klf4, c-Myc) to induce pluripotency in somatic cells. No single-cell variational inference (scVI) analyses were performed to assess the molecular signatures of reprogrammed cells.

Experimental Interventions:

- Oncogenic Stress Induced
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)


3. Results

The final cell population analysis revealed 0 somatic cells, 0 iPSCs, 0 neuronal cells, 0 cardiomyocytes, and 2000 tumor cells. This outcome indicates a complete conversion of initial cell populations into tumorigenic cells, with no successful induction of pluripotency or differentiation into targeted cell lineages.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	0	0.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	2000	100.0%

4. Discussion

The exclusive emergence of tumor cells suggests oncogenic stress, combined with repeated OSKM transfections, predominantly promotes tumorigenesis rather than reprogramming to a pluripotent state. This highlights the potential risks associated with unchecked oncogene activation in reprogramming protocols. Future investigations should focus on modifying the reprogramming conditions to mitigate oncogenic pathways while enhancing iPSC generation efficiency.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.