

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-01T10-44-18-505Z

Run ID: #2

Date Generated: 2026-01-01 10:44:23

Total Cells: 2000 | Model Mode: CLINICAL

1. Introduction

Cellular reprogramming has emerged as a pivotal technique in regenerative medicine, aiming to revert somatic cells to a pluripotent state. This experiment was designed to evaluate the efficacy of repetitive Yamanaka factor (OSKM) transfection in inducing pluripotency in differentiated cells subjected to oncogenic stress.

2. Materials and Methods

The experiment employed six sequential transfections with Yamanaka factors (OSKM) on cells pre-stressed with oncogenic factors to induce cellular reprogramming. Computational analysis was planned to use single-cell variational inference (scVI) but was not executed, focusing instead on direct population outcomes.

Experimental Interventions:

- Oncogenic Stress Induced
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)


3. Results

Post-intervention analysis revealed a complete lack of somatic, induced pluripotent stem (iPSC), neurogenic, and cardiogenic cell populations, with a dominant emergence of 2000 tumor cells. The absence of the desired pluripotent and differentiated cell types indicates a failure in reprogramming under the given conditions, with a pronounced shift towards tumorigenesis.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	0	0.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	2000	100.0%

4. Discussion

The results underscore the challenge of balancing oncogenic stress with reprogramming efficiency, as evidenced by the sole proliferation of tumor cells. The findings suggest that oncogenic stress may have overpowered the reprogramming signals, leading to malignant transformation rather than pluripotency. These outcomes highlight the necessity for optimizing stress levels and transfection protocols to achieve successful cellular reprogramming without oncogenic risks.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.