

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-01T11-13-23-810Z

Run ID: #4

Date Generated: 2026-01-01 11:13:28

Total Cells: 2000 | Model Mode: CLINICAL

1. Introduction

Cellular reprogramming holds significant promise for regenerative medicine, allowing for the conversion of somatic cells into induced pluripotent stem cells (iPSCs). This experiment aimed to assess the efficiency of the Yamanaka factors (OSKM) in reprogramming somatic cells to iPSCs within a controlled laboratory setting.

2. Materials and Methods

In this study, six sequential OSKM transfections were performed on a population of somatic cells to induce pluripotency. The experiment utilized standard protocols for the transfection process; however, no single-cell variational inference (scVI) analyses were conducted to further investigate cellular states.

Experimental Interventions:

- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)


3. Results

Post-intervention analysis revealed that the final cellular population consisted entirely of 2000 somatic cells, with no transformation into iPSCs, neurogenic, cardiogenic, or tumorigenic lineages observed. This suggests that the OSKM transfections did not successfully induce pluripotency or alter the cellular phenotype under the conditions tested.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	2000	100.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	0	0.0%

4. Discussion

The absence of iPSC formation following multiple rounds of OSKM transfections indicates potential inefficiencies in the reprogramming protocol or cellular resistance mechanisms. Further investigations involving scVI analyses could provide insights into the molecular barriers impeding the transition to iPSCs. These findings underscore the necessity for optimization of transfection conditions and consideration of cell-specific factors that may influence reprogramming success.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.