

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-06T04-20-13-530Z

Run ID: #4

Date Generated: 2026-01-06 04:20:23

Total Cells: 1984 | Model Mode: CLINICAL | Avg BioAge: N/A

1. Introduction

Cellular reprogramming holds significant potential for regenerative medicine, particularly through the conversion of somatic cells into induced pluripotent stem cells (iPSCs) or directly into specific cell lineages. This experiment aimed to evaluate the effectiveness of various reprogramming techniques, including OSKM transfection and direct cardiac transdifferentiation, in altering the fate of somatic cells.

2. Materials and Methods

The experiment involved a series of interventions, including multiple OSKM transfections, direct cardiac transdifferentiation, and the induction of oncogenic stress. Computational analysis was conducted using two single-cell variational inference (scVI) analyses to assess the cellular populations resulting from these interventions.

Experimental Interventions:

- Direct Cardiac Transdifferentiation
- OSKM Transfection (Yamanaka)
- Oncogenic Stress Induced
- Oncogenic Stress Induced
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- Direct Cardiac Transdifferentiation
- Direct Cardiac Transdifferentiation
- Oncogenic Stress Induced


3. Results

The final population analysis revealed a total of 1984 somatic cells, with no detectable iPSCs, neuronal, cardiac, or tumor cells. Despite repeated applications of OSKM transfection and direct cardiac transdifferentiation, the interventions did not yield the expected reprogramming into the target cell types. The absence of iPSCs and differentiated cells suggests a potential resistance to reprogramming under the given experimental conditions.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	1984	100.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	0	0.0%

3.2 Transcriptomic Analysis (scVI)

High-Resolution Gene Expression Analyses Performed: 2

Cells were expanded to high-dimensional transcriptomic space using a pre-trained variational autoencoder model on Human Cell Atlas data.

4. Discussion

The persistence of the somatic cell population and the lack of differentiation into iPSCs or other target lineages highlight challenges in the reprogramming process. Factors such as cellular resistance, insufficient expression of reprogramming factors, or inadequate conditions for cell fate transition may have contributed to the outcome. These findings underscore the complexity of cellular reprogramming and the need for optimizing intervention strategies to enhance reprogramming efficiency in future studies.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.