

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-01T10-44-34-074Z

Run ID: #2

Date Generated: 2026-01-01 10:44:44

Total Cells: 2000 | Model Mode: CLINICAL

1. Introduction

Cellular reprogramming using Yamanaka factors (OSKM: Oct4, Sox2, Klf4, and c-Myc) is a pivotal method for converting somatic cells into induced pluripotent stem cells (iPSCs). This experiment aimed to evaluate the effects of repeated OSKM transfections under oncogenic stress conditions on cellular fate determination.

2. Materials and Methods

The experimental setup involved six sequential transfections of the Yamanaka factors into a somatic cell population under induced oncogenic stress. Computational analysis, specifically scVI, was not employed during this experimental run, focusing instead on direct population outcomes.

Experimental Interventions:

- Oncogenic Stress Induced
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)


3. Results

Post-intervention, the final cellular composition comprised entirely of tumor cells, with a total count of 2000, while no somatic, iPSC, neurogenic, or cardiogenic cells were detected. This suggests a complete conversion or replacement of the initial cell types with tumorigenic cells, under the applied conditions.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	0	0.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	2000	100.0%

4. Discussion

The absence of iPSC and other differentiated cell types, combined with the exclusive presence of tumor cells, indicates a potential proclivity towards tumorigenesis under the stress and repeated OSKM transfections. These findings suggest that oncogenic stress might override reprogramming efforts, highlighting the risk of tumor formation in such cellular interventions. Further studies should investigate the underlying mechanisms and explore strategies to mitigate tumorigenic outcomes during reprogramming.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.