

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-01T10-44-34-349Z

Run ID: #2

Date Generated: 2026-01-01 10:44:50

Total Cells: 2000 | Model Mode: CLINICAL

1. Introduction

Cellular reprogramming through the introduction of Yamanaka factors (OSKM: Oct4, Sox2, Klf4, and Myc) has shown potential in converting somatic cells into induced pluripotent stem cells (iPSCs). This experiment aimed to evaluate the impact of repeated OSKM transfections on cellular fate under conditions of oncogenic stress, seeking to elucidate potential pathways for iPSC generation or tumorigenesis.

2. Materials and Methods

The experimental protocol involved the repeated transfection of somatic cells with the OSKM factors, performed six times, under oncogenic stress conditions. No scVI computational analyses were conducted on the resultant cell populations, allowing for direct observation of phenotypic outcomes.

Experimental Interventions:

- Oncogenic Stress Induced
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)


3. Results

The final cellular populations revealed a complete absence of somatic, iPSC, neurogenic, and cardiogenic cells, with an exclusive emergence of 2000 tumor cells. The repeated OSKM transfections under oncogenic stress did not facilitate the generation of iPSCs or other differentiated cell types, but rather, led exclusively to tumor formation.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	0	0.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	2000	100.0%

4. Discussion

The results suggest that the repeated application of OSKM factors under oncogenic stress conditions promotes tumorigenesis rather than reprogramming to a pluripotent or differentiated state. This underscores the critical role of cellular context and stress factors in influencing the reprogramming outcome. The findings highlight the necessity for careful modulation of oncogenic signals during reprogramming efforts to avoid adverse oncogenic transformations.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.