

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-01T11-11-56-775Z

Run ID: #4

Date Generated: 2026-01-01 11:12:12

Total Cells: 2000 | Model Mode: CLINICAL

1. Introduction

Cellular reprogramming holds significant promise in regenerative medicine by enabling the conversion of somatic cells into pluripotent stem cells. This study aims to evaluate the efficiency of the OSKM transfection method, a technique pioneered by Yamanaka, in inducing somatic cells to transdifferentiate into induced pluripotent stem cells (iPSCs).

2. Materials and Methods

In this experiment, somatic cells were subjected to repeated transfections using the OSKM transcription factors, which include Oct4, Sox2, Klf4, and c-Myc. No single-cell variational inference (scVI) analyses were conducted as part of this intervention.

Experimental Interventions:

- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)


3. Results

The final cell population analysis revealed a total of 2000 somatic cells with no detectable conversion into iPSCs, neurons, cardiomyocytes, or tumor cells. Despite six rounds of OSKM transfection, the absence of iPSC formation indicates a lack of cellular reprogramming under the current experimental conditions.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	2000	100.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	0	0.0%

4. Discussion

The failure to observe any iPSC generation suggests potential limitations in the reprogramming efficiency of the OSKM transfection method used in this study. This result highlights the necessity for optimization of transfection conditions or exploring alternative reprogramming strategies. Further investigation into the underlying biological barriers preventing reprogramming could provide insights to enhance the efficacy of iPSC induction methods.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.