

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis

Session ID: session_2026-01-06T03-42-33-443Z
Run ID: #2
Date Generated: 2026-01-06 03:42:44
Total Cells: 1999 | Model Mode: CLINICAL | Avg BioAge: N/A

1. Introduction

Cellular reprogramming has emerged as a pivotal approach in regenerative medicine, offering prospects for generating induced pluripotent stem cells (iPSCs) from somatic cells. This experiment aimed to explore the effects of repeated OSKM transfection, oncogenic stress, and p53 therapeutic interventions on cellular fate, with an additional focus on direct neuronal transdifferentiation.

2. Materials and Methods

We employed a series of experimental interventions comprising three sequential OSKM transfections, four oncogenic stress inductions, and three p53 therapeutic interventions, followed by an attempt at direct neuronal transdifferentiation. Single-cell variational inference (scVI) analyses were conducted four times to assess cellular composition changes across experimental conditions.

Experimental Interventions:

- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- OSKM Transfection (Yamanaka)
- Oncogenic Stress Induced
- Oncogenic Stress Induced
- Oncogenic Stress Induced
- Oncogenic Stress Induced
- P53 Therapeutic Intervention
- P53 Therapeutic Intervention
- P53 Therapeutic Intervention
- Direct Neuronal Transdifferentiation


3. Results

The final cellular population consisted solely of 1999 somatic cells, with zero cells differentiating into iPSCs, neurons, cardiomyocytes, or tumor cells. Despite multiple OSKM transfections and oncogenic stress applications, no successful reprogramming or transdifferentiation was observed. The p53 therapeutic intervention and direct neuronal transdifferentiation also failed to yield any non-somatic cell types.

3.1 Population Distribution

Clinical Digital Twin Report

Computational Cellular Reprogramming Analysis



Cell Type	Count	Percentage
SOMATIC	1999	100.0%
IPSC	0	0.0%
NEURO	0	0.0%
CARDIO	0	0.0%
TUMOR	0	0.0%

3.2 Transcriptomic Analysis (scVI)

High-Resolution Gene Expression Analyses Performed: 4

Cells were expanded to high-dimensional transcriptomic space using a pre-trained variational autoencoder model on Human Cell Atlas data.

4. Discussion

The persistence of somatic cells and absence of iPSCs or other differentiated cell types suggest a potential resistance to reprogramming under the given experimental conditions. The repeated OSKM transfections and oncogenic stress may have induced cellular senescence or apoptosis, preventing successful lineage conversion. Additionally, the ineffectiveness of the p53 intervention implies that p53 status might not be the sole determinant of reprogramming efficiency. Future investigations should consider alternative approaches to overcome reprogramming barriers and enhance transdifferentiation success.

This report was automatically generated by the IS-CHRP v25.0 Clinical AI Bridge platform. The data represents computational simulation results from a high-fidelity cellular reprogramming model.