sum of 9 is:

1. The probability of	f a lean year	selected at rai	ndom contain 53	
Sunday is:	a leap year	Science at iai	2011 0011011	
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365	
2. A bag contains 3	red and 2 blu	e marbles. A r	narble is drawn at	
random. The probabi	lity of drawi	ng a black ball	is:	
(a) 3/5 (t	2/5	(c) 0/5/	(d) 1/5	
3. The probability th	at it will rain	tomorrow is 0	.85. What is the	
probability that it will	not rain tor	norrow		
(a) 0.25 (b)	0.145	(c) 3/20	(d) none of these	
4. What is the proba	bility that a	number select	ed from the numbers	
(1, 2, 3,,15) is	a multiple o	f 4?		
(a) 1/5 (b	) 4/5	(c) 2/15	(d) 1/3	
5. What are the total	outcomes	when we throv	v three coins?	
(a) 4 (b	) 5	(c) 8"	(d) 7	
<ol><li>The probability th</li></ol>	at a prime r	number selecte	ed at random from the	
numbers (1,2,3,	35) is :	The second second second		
(a) 12/35	(b) 11/35	(c) 13/35	(d) none of these	
7. The sum of the pr	obability of	an event and i	non event is :	
(a) 2 (b)	17 (c) (	) (d) nor	e of these.	
		re given; choo	se the correct answer	
for that which is not p	oossible.	Constant of		
(a) 0.15 (b)	2/7	(c) 7/5	(d) none of these.	
9. If three coins are t	ossed simu	ıltaneously, th	an the probability of	
getting at least two h	eads, is:			
(a) 1/4 (b)	3/8	(c) ½	(d) 1/8	
10. A letter is chose	n at randon	n from the lett	ers of the word	
<b>♦</b> ASSASSINATION €	. The prob	ability that the	letter chosen has:	
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.	
(0) 0/10	(-)			
11 A dice is thrown	Find the pro	obability of ge	tting an even number.	
	) 1	(C) 5/6	(D) 1/2	
(A) 2/3 (B	, ,	(0) 0/0		
12. Two coins are thr	own at the	same time. Fi	nd the probability of	
	OWIII at the	outile time		
getting both head.	(0) 1/2	(D) 0		
(A) 3/4 (B) 1/4	(0) 1/2	(D) 0		
		neously The	probability of getting	a
13. Two dice are thro	wn simuita	neously. The	hionaning or detting	-

(A) 1/1	0 (B) 3/1	0 (C)	1/9	(D) 4/9		
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.						
(A) 3/4		(C) 1	14	(D) 29/1	00	
of draw	ag contains 5 re ing a blue ball i lls in a bag is: (B) 10	s double tha	some blue t of a red ( (D)	ball, then the	probability number of	
		(C) 15			-	
taken o	ox of 600 bulbs ut at random fro ective bulb is:					
(A) 143/		47/150	(C) 1/25	(D)	1/50	
mixed th	Is marked with noroughly. One ability that the 0 (B) 1/10	card is draw	vn from th card is a p	is box rando	omly, then	
18. Wha (A) 1/7	t is the probab (B) 53/366			ndays in a le (D) 7/366		
probabili	rd is drawn fro ty of getting a (B) 3/26	king of red	suit.		ls. Find the	
equally li 1,2,31	me of chance of kely to come to 2 ,then the pro (B) 1/12	o rest pointi bability tha	ing to one It it will po	of the num	ber	
its outcor result i.e.	ne consists of ne each time. three heads o	Aryan wins r three tails	if all the tand lose	tosses give	the same	
F.201 / F.	y that Aryan w (B) 1/2 (C)		_			
(A) 3/4	(b) 1/2 (C)	(L	7 1/4			

32.Out of probability	the following va	alues, which one is	not possibl	le in
	b) ∑ x	P(x) = 3 x = -0.5		
33.If E(x) a) 2	= <b>2 and E(z)</b> = <b>4</b> b) 6	c) 0	d) Insu	fficient data
34.The co	variance of two	independent rando	om variable	is
a) 1	b) <mark>0</mark>	c) - 1	d) Und	efined
35.If Σ P() a) 0	k) = k <sup>2</sup> – 8 then, b) 1	the value of k is?	d) Insi	ufficient data
36.If P(x) = a) 1	= <b>0.5 and x = 4,</b> b) 0.5	then E(x) = ? c) 4	d) 2	
37.In a dise is always?	2	y distribution, the	sum of all d) Und	
38.If the p variance. a) 0.4, 0.24		tting the target is 0.24 c) 0	<b>0.4, find m</b> 0.4, 0.16	ean and d) 0.6, 0.16
39.If the p target is 60 a) 0.6, 0.24	)% and if 10 bo	a bomb dropped f mbs are dropped, c) 0.4, 0.1	find mean	e will strike the and variance? d) 4, 1.6
a) 2	e mean of toss b) 4 s the mean and	ing 8 coins. c) 8 I variance for star	d) 1 ndard nc	nal distribution?

c) 5

a) 3

b) 4

d) 7

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:
(A) 364/365 (B) 31/365 (C) 1/365 (D) 1/133225
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is?
(A) 1/5 (B) 2/5 (C) 3/5 (D) 4/5
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is:  (A) 10 (B) 6 (C) 8 (D) 7
25. A number is selected at random from first 50 natural numbers.  Then the probability that it is a multiple of 3 and 4 is:  (A) 7/50 (B) 4/25 (C) 1/25 (D) 2/25
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?
a) $\frac{1}{7}$ b) $\frac{5}{42}$ c) $\frac{1}{21}$ d) $\frac{4}{21}$
27. Runs scored by batsman in 5 one day matches are 50, 70, 82, 93, and 20. The standard deviation is
a) 25.79 b) 25.49 c) 25.29 d) 25.69
28. Find median and mode of the messages received on 9
consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17. a) 13, 15 b) 13, 18 c) 18, 15 d) 13, 16
29. A coin is tossed up 4 times. The probability that tails turn up in
a) $\frac{1}{2}$ b) $\frac{1}{3}$ c) $\frac{1}{4}$ d) $\frac{1}{6}$ 30. X is a variate between 0 and 3. The value of $E(X^2)$ is
a) 8 b) 7 c) 27 d) 9
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?

a) <u>N</u>	Mean is Mean is	0 and varia 0 and varia	nce is 1/ nce is co	b) Mear d) Mea	n is 1 and In is ∞ ar	variance is nd variance	s 0 e is 0
42. Variance of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2)$ - $E(X2)$ - $E(X2)$ d) $E(X)$ 2							
-	43.Mean of a random variable X is given by a) E(X) / b) E(X2) c) E(X2) - (E(X))2 d) (E(X))2						
	44.Mean of a constant 'a' is a) 0 b) a c) a/2 d) 1						
	45. Variance of a constant 'a' is a) 0						
46.Find the mean and variance of X?							
	x	0	1	2	3	4	
	f(x)	1/9	2/9	3/9	2/9	1/9	
a) <mark>2</mark> ,	4/3	<b>b)</b> 3	, 4/3		c) 2, 2/3		d) 3, 2/3
47.Find the expectation of a random variable X?							
	x	0 1	2 3				
	f(x) 1	/6 2/6	2/6 1/6	70 1			
a) O.	5	b) 1.5		c) 2.5		d) 3.5	
	and the Land			· · · · · ·	d	munhahili	w of oursess

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

لممسوحة ضوئيا بـ CamScanner

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use Binomial Distribution.
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard Deviation?
- a)  $\sqrt{np}$
- b) $\sqrt{pq}$  c) (np)2
- d)  $\sqrt{npq}$