

UAE Medical and Behavioral Paper 1

Contents

1	Measurement of Weight Status	2
2	Participant Characteristics	3
2.1	Associations between Demographics and Percent of Overweight Cutoff	4
3	Medical Comorbidities	6
3.1	Number of Comorbidities	7
3.2	Distribution tests by Weight Status	7
3.3	Association between Comorbidities and Percent of Overweight Cutoff	8
4	Family History	10
4.1	Association with Yes/No Family History	10
4.2	Family Members with History of Obesity	11
4.3	Family Members with History of Eating Disorders	12
5	Sleep	14
5.1	Overall Averages	14
5.2	Association Between Sleep Sub-Scales and Percent of Overweight Cutoff	15
6	Strengths and Difficulties Questionnaire	19
6.1	Exploratory Sex x Percent of Overweight Models	20
6.2	Conduct Problems	21
6.3	Prosocial	23
7	Extra Tables by Sex	24
7.1	Participant Characteristics	24
7.2	Medical Comorbidities	25
7.3	Family History	26
7.4	Sleep	27
7.5	Strengths and Difficulties Questionnaire	28


1 Measurement of Weight Status

We decided to use the International Obesity Task Force (IOTF) designation of weight status for the sample. They use smoothed, sex-specific BMI curves meant to match the the BMI cutoffs for overweight (OW; 25 kg/m^2) and obesity (OB; 30 kg/m^2) at age 18 yrs.

Rather than BMI-zscore or BMI-percentile, we chose to use percent of overweight cutoff because recent studies shows it has a tighter association with measured adiposity:

$$\text{BMI \% of overweight} = \frac{\text{childBMI}}{\text{BMI at age- and sex- adjusted overweight cutoff}} * 100$$

<100 % - indicates child BMI is below the overweight cutoff for age and sex (i.e., has healthy weight) 100 % - indicates child BMI is the same as the overweight cutoff for age and sex >100 % - indicates child BMI is above the overweight cutoff for age and sex (i.e., has overweight or obesity)



Density plot of percent of overweight by sex. The shaded regions indicated those with healthy weight (blue), overweight (yellow), and obesity (red). The points show density of participants by sex (purple circles = female, orange triangles = males).

2 Participant Characteristics

Table 1: Demographic Characteristics by Weight Status

Characteristic	N	HW, N = 41	OW, N = 29	OB, N = 37
sex	107			
F		24 (59%)	14 (48%)	23 (62%)
M		17 (41%)	15 (52%)	14 (38%)
Age_yr	107	11.85 [8.02 - 17.37]	12.84 [8.15 - 17.54]	13.69 [7.31 - 17.84]
BMI	107	17.15 [12.71 - 22.72]	24.03 [18.70 - 28.86]	35.08 [21.87 - 55.52]
IOTF_pOWcutoff	107	80.86 [63.95 - 98.26]	109.66 [100.39 - 120.73]	155.80 [122.38 - 239.00]
Father_ed	102	12.79 [6.00 - 18.00]	13.60 [6.00 - 18.00]	12.20 [0.00 - 18.00]
Unknown		3	0	2
Mother_ed	99	13.38 [3.00 - 18.00]	13.93 [9.00 - 18.00]	12.40 [0.00 - 18.00]
Unknown		4	2	2
Month_AED	99			
<25,000 AED		10 (27%)	10 (34%)	11 (33%)
25,000 - 55,000 AED		21 (57%)	13 (45%)	19 (58%)
55,000 - 75,000 AED		2 (5.4%)	3 (10%)	1 (3.0%)
> 75,000 AED		4 (11%)	3 (10%)	2 (6.1%)
Unknown		4	0	4
DadNationality	101			
Emirati		40 (100%)	25 (96%)	33 (94%)
Omani		0 (0%)	1 (3.8%)	0 (0%)
Yemeni		0 (0%)	0 (0%)	2 (5.7%)
Unknown		1	3	2
MomNationality	104			
Emirati		38 (93%)	26 (93%)	32 (91%)
Omani		0 (0%)	1 (3.6%)	0 (0%)
Yemeni		0 (0%)	0 (0%)	1 (2.9%)
Moroccan		1 (2.4%)	0 (0%)	1 (2.9%)
Egyptian		2 (4.9%)	0 (0%)	1 (2.9%)
Bahrani		0 (0%)	1 (3.6%)	0 (0%)
Unknown		0	1	2

¹ n (%); Mean [Range]

2.1 Associations between Demographics and Percent of Overweight Cutoff

2.1.1 t-test for sex

Welch Two Sample t-test

```
data: IOTF_pOWcutoff by sex
t = -0.72449, df = 78.92, p-value = 0.4709
alternative hypothesis: true difference in means between group F and group M is not equal to 0
95 percent confidence interval:
 -20.558399  9.586342
sample estimates:
mean in group F mean in group M
      112.2213      117.7073
```

2.1.2 Correlation Matrix

Table 2: Correlations between percent of overweight cutoff and demographic characteristics

	Age_yr	Father_ed	Mother_ed	IOTF_pOWcutoff
Age_yr				
Father_ed	-0.02			
Mother_ed	-0.2*	0.51*		
IOTF_pOWcutoff	0.26*	-0.07	-0.15	

Table 3: P-values for the correlations between percent of overweight cutoff and demographic characteristics

	Age_yr	Father_ed	Mother_ed	IOTF_pOWcutoff
Age_yr				
Father_ed	0.806			
Mother_ed	0.049	0		
IOTF_pOWcutoff	0.006	0.491	0.131	

Only child age was associated with percent of overweight cutoff - older children tended to have higher percent of overweight cutoff indicating older children were more likely to have overweight or obesity. There was no association with father or mother education level, which differs from finding in the US. Hip to waist ratio was also not associated with percent of overweight cutoff.

2.1.3 One-Way ANOVA for Income Categories

Anova Table (Type III tests)

```
Response: IOTF_pOWcutoff
      Sum Sq Df F value Pr(>F)
(Intercept) 406511 1 282.6340 <2e-16 ***
Month_AED    1046 3  0.2424 0.8665
Residuals   136638 95
```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

There was no difference in percent of overweight by monthly income category.

2.1.4 Sensitivity Tests

Table 4: Linear Model: pOWcutoff - SES category + Maternal Education + Age + Sex

	b	se	t	p	
(Intercept)	71.310	27.957	2.551	0.012	
Month_AED25,000 - 55,000 AED	9.699	8.663	1.120	0.266	
Month_AED55,000 - 75,000 AED	19.268	16.747	1.150	0.253	
Month_AED> 75,000 AED	2.047	15.847	0.129	0.898	
Mother_ed	-1.654	1.254	-1.319	0.191	
Age_yr	4.239	1.464	2.894	0.005	***
sexM	10.293	7.621	1.351	0.180	

After controlling for family income, mother education, and child sex, child age was significantly associated with percent of overweight such that for each year older, the expected percent of overweight is predicted to increase 4.24%.

3 Medical Comorbidities

Table 5: Medical Comorbidities by Weight Status

Characteristic	N	HW, N = 41	OW, N = 29	OB, N = 37
nComorbid	107	2.05 [0.00 - 4.00]	2.24 [1.00 - 4.00]	2.19 [0.00 - 4.00]
VitDdeficiency	107			
Y		36 (88%)	29 (100%)	33 (89%)
N		5 (12%)	0 (0%)	4 (11%)
Anemia	40			
Iron Deficiency Anemia (ID)		9 (39%)	7 (70%)	5 (71%)
Thalassemia Minor (TM)		2 (8.7%)	1 (10%)	0 (0%)
G6PD Deficiency		1 (4.3%)	0 (0%)	0 (0%)
ID + TM		1 (4.3%)	1 (10%)	0 (0%)
ID + G6PD Deficiency		1 (4.3%)	0 (0%)	0 (0%)
Unspecified Anemia		9 (39%)	1 (10%)	2 (29%)
Unknown		18	19	30
Hyperlipidemia	12			
Hyperlipidemia		0 (NA%)	5 (100%)	6 (86%)
Hyperlipidemia - Mixed		0 (NA%)	0 (0%)	1 (14%)
Unknown		41	24	30
ThyroidConditions	25			
Abnormal Function		5 (71%)	4 (44%)	5 (56%)
Autoimmune Thyroiditis		1 (14%)	2 (22%)	2 (22%)
Autoimmune Hypothyroidism		0 (0%)	0 (0%)	0 (0%)
Unspecified Hypothyroidism		1 (14%)	2 (22%)	2 (22%)
Goiter		0 (0%)	1 (11%)	0 (0%)
Unknown		34	20	28
GlycemicStatus	28			
Impaired Fasting Glucose		4 (57%)	8 (73%)	7 (70%)
Impaired Glucose Tolerance Test		2 (29%)	2 (18%)	3 (30%)
Type-1 Diabetes		1 (14%)	1 (9.1%)	0 (0%)
Unknown		34	18	27
Acanthosis Nigricans	8	1 (100%)	0 (NA%)	7 (100%)
Unknown		40	29	30
Hypertension	3			
Essential Primary Hypertension		0 (NA%)	0 (NA%)	2 (67%)
High Blood Pressure		0 (NA%)	0 (NA%)	1 (33%)
Unknown		41	29	34
Metabolic Syndrome	2	1 (100%)	0 (NA%)	1 (100%)
Unknown		40	29	36
Growth.Stature	10			
Failure To Thrive (FT)		1 (14%)	0 (0%)	0 (0%)
Growth Hormone Deficiency		1 (14%)	0 (0%)	0 (0%)
Short Stature		3 (43%)	1 (100%)	2 (100%)
FT + ShortStature + Underweight		1 (14%)	0 (0%)	0 (0%)
Short Stature + Precocious Puberty		1 (14%)	0 (0%)	0 (0%)
Unknown		34	28	35
PCOS	4			
PCOS		1 (50%)	0 (NA%)	1 (50%)
Hirsutism		0 (0%)	0 (NA%)	1 (50%)
Hirsutism + Unspecified Ovarian Cysts		1 (50%)	0 (NA%)	0 (0%)
Unknown		39	29	35

¹ Mean [Range]; n (%)

3.1 Number of Comorbidities

3.1.1 mean (sd) number of comorbidities

```
[1] 2.149533
```

```
[1] 1.035263
```

3.1.2 Association with Number of Comorbidities - Poisson

Table 6: Anemia Status by Weight Status

	b	e ^b	se	e ^{2.5} CI	e ^{97.5} CI	z	p
(Intercept)	0.659	1.932	0.214	1.271	2.939	3.080	0.002
IOTF_pOWcutoff	0.001	1.001	0.002	0.997	1.004	0.527	0.598

There was no association between percent of overweight cutoff and number of comorbidities

3.1.3 sex differences in number of comorbidities

Welch Two Sample t-test

```
data: nComorbid by sex
```

```
t = 1.125, df = 101.31, p-value = 0.2632
```

```
alternative hypothesis: true difference in means between group F and group M is not equal to 0
```

```
95 percent confidence interval:
```

```
-0.1710783 0.6194034
```

```
sample estimates:
```

```
mean in group F mean in group M
```

```
2.245902      2.021739
```

```
F      M
```

```
1.0747982 0.9772781
```

3.2 Distribution tests by Weight Status

3.2.1 Vitamin D

Fisher's Exact Test for Count Data

```
data: xtabs(~VitDdeficiency + IOTF_3class, data = UAE_allDat)
```

```
p-value = 0.132
```

```
alternative hypothesis: two.sided
```

3.2.2 Anemia

Table 7: Anemia Status by Weight Status

	HW	OW	OB
N	18	19	30
Y	23	10	7

Pearson's Chi-squared test

```
data: xtabs(~as.factor(ifelse(is.na(UAE_allDat$Anemia), "N", "Y"))) + IOTF_3class, data = UAE_allDat
X-squared = 11.627, df = 2, p-value = 0.002986
```

3.2.3 Hyperlipidemia

Table 8: Hyperlipidemia Status by Weight Status

	HW	OW	OB
N	41	24	30
Y	0	5	7

Fisher's Exact Test for Count Data

```
data: xtabs(~as.factor(ifelse(is.na(UAE_allDat$Hyperlipidemia), "N", "Y"))) + IOTF_3class, data = UAE_allDat
p-value = 0.004283
alternative hypothesis: two.sided
```

3.2.4 Thyroid Conditions

Pearson's Chi-squared test

```
data: xtabs(~as.factor(ifelse(is.na(UAE_allDat$ThyroidConditions), "N", "Y"))) + IOTF_3class, data = UAE_allDat
X-squared = 1.8782, df = 2, p-value = 0.391
```

3.2.5 Glycemic Status

Pearson's Chi-squared test

```
data: xtabs(~as.factor(ifelse(is.na(UAE_allDat$GlycemicStatus), "N", "Y"))) + IOTF_3class, data = UAE_allDat
X-squared = 3.8464, df = 2, p-value = 0.1461
```

3.3 Association between Comorbidities and Percent of Overweight Cutoff

3.3.1 t-tests for Presence/Absence of Comorbidities

Table 9: t-tests for percent of overweight by absence vs presence of medical comorbidity

	AbsentMean	PresentMean	t	df	pvalue	sig
VitD Deficiency	114.56	114.84	-0.02	9.25	0.984	
Anemia	122.62	101.12	3.01	81.89	0.004	**
Thyroid Dysfunction	113.40	118.44	-0.59	39.42	0.559	
Glycemic Status	112.51	120.43	-0.86	39.25	0.393	

Table 10: Standard deviations for percent of overweight by absence vs presence of anemia

	x
N	35.677
Y	35.828

There were no difference in percent of overweight cutoff by presence/absence of Thyroid dysfunction or impaired glucose function. However, those with anemia tended to have lower percent of overweight. The mean with anemia was 101% indicating children were overweight on average. The mean for those without anemia were 123%, indicating the children were above the overweight cutoff.

4 Family History

Table 11: Family History by Weight Status

Characteristic	N	HW, N = 41	OW, N = 29	OB, N = 37
Fam_OB_YN	101			
yes		18 (49%)	21 (72%)	31 (89%)
no		19 (51%)	8 (28%)	4 (11%)
Unknown		4	0	2
nFam_Obesity	107	1.15 [0.00 - 6.00]	1.59 [0.00 - 4.00]	2.70 [0.00 - 7.00]
Mother	107	5 (12%)	2 (6.9%)	11 (30%)
Father	107	6 (15%)	4 (14%)	11 (30%)
Grandmother	107	11 (27%)	9 (31%)	13 (35%)
Grandfather	107	0 (0%)	2 (6.9%)	4 (11%)
Sister	107	5 (12%)	3 (10%)	8 (22%)
Brother	107	5 (12%)	3 (10%)	17 (46%)
Aunt	107	10 (24%)	17 (59%)	18 (49%)
Uncle	107	5 (12%)	6 (21%)	18 (49%)
Fam_ED_YN	96			
yes		2 (5.6%)	5 (19%)	4 (12%)
no		34 (94%)	22 (81%)	29 (88%)
Unknown		5	2	4
nFam_EatingDisorder	107	0.05 [0.00 - 1.00]	0.28 [0.00 - 2.00]	0.35 [0.00 - 7.00]
Mother	107	0 (0%)	0 (0%)	1 (2.7%)
Father	107	0 (0%)	0 (0%)	2 (5.4%)
Grandmother	107	1 (2.4%)	1 (3.4%)	2 (5.4%)
Grandfather	107	0 (0%)	1 (3.4%)	1 (2.7%)
Sister	107	1 (2.4%)	1 (3.4%)	0 (0%)
Brother	107	0 (0%)	1 (3.4%)	1 (2.7%)
Aunt	107	0 (0%)	3 (10%)	4 (11%)
Uncle	107	0 (0%)	1 (3.4%)	2 (5.4%)

¹ n (%); Mean [Range]

4.1 Association with Yes/No Family History

4.1.1 t-tests for Yes/No History of Obesity

Welch Two Sample t-test

data: IOTF_pOWcutoff by Fam_OB_YN

t = 4.828, df = 87.915, p-value = 5.776e-06

alternative hypothesis: true difference in means between group yes and group no is not equal to 0

95 percent confidence interval:

17.83736 42.79491

sample estimates:

mean in group yes mean in group no

125.08610 94.76996

Table 12: Standard deviations for percent of overweight by absence vs presence of family history of obesity

	x
yes	38.383
no	23.871

There was no differences in percent of overweight for those whose families had a history of eating disorder (reported by parent). There was a significant difference in percent of overweight between families that reported a history of obesity ('Yes') and those who did not ('No'). Those without a family history of obesity had a mean percent of overweight equal to 94%, indicating the children had healthy weight on average. Children with a family history of obesity had a mean percent of overweight equal to 124%, indicating the children had overweight or obesity.

4.1.2 distribution test for Yes/Now History of Eating Disorders across Weight Status

Fisher's Exact Test for Count Data

```
data: xtabs(~IOTF_3class + Fam_ED_YN, data = UAE_allDat)
p-value = 0.2633
alternative hypothesis: two.sided
```

4.2 Family Members with History of Obesity

Table 13: Overall Family Relationship Categories with History of Obeisty

Characteristic	N	N = 107
Mother	107	18 (17%)
Father	107	21 (20%)
Grandmother	107	33 (31%)
Grandfather	107	6 (5.6%)
Sister	107	16 (15%)
Brother	107	25 (23%)
Aunt	107	45 (42%)
Uncle	107	29 (27%)

¹ n (%)

4.2.1 mean (sd) number of family relationship categories

[1] 1.803738

[1] 1.750683

4.2.2 Association with Number of Relative Categories with History of Obesity - Poisson

Table 14: Poisson Model: Number of Family Relationship Categories Obeisty - pOWcutoff

	b	e ^b	se	e ^{2.5} CI	e ^{97.5} CI	z	p	
(Intercept)	-0.5512353	0.5762375	0.2254574	0.369320	0.8942968	-2.444964	0.0144866	*
IOTF_pOWcutoff	0.0093704	1.0094144	0.0016455	1.006104	1.0126204	5.694416	0.0000000	***


Percent of overweight cutoff was associated with reported number of relative categories with family history of obesity.

4.2.3 Sensitivity Tests

Table 15: Linear Model: Number of Family Relationship Categories Obesity - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	e ^b	se	e ^{2.5} CI	e ^{97.5} CI	z	p	
(Intercept)	-0.180	0.576	0.577	0.369	0.894	-0.313	0.754	
Month_AED25,000 - 55,000 AED	0.113	1.009	0.182	1.006	1.013	0.622	0.534	
Month_AED55,000 - 75,000 AED	0.380	0.576	0.307	0.369	0.894	1.239	0.215	
Month_AED> 75,000 AED	-0.127	1.009	0.372	1.006	1.013	-0.342	0.732	
Mother_ed	-0.029	0.576	0.024	0.369	0.894	-1.221	0.222	
Age_yr	0.011	1.009	0.031	1.006	1.013	0.335	0.738	
sexM	-0.111	0.576	0.159	0.369	0.894	-0.703	0.482	
IOTF_pOWcutoff	0.008	1.009	0.002	1.006	1.013	4.360	0.000	***

After controlling for family income, mother education, child age, and child sex, percent of overweight showed a trend-level association with number of family members with history of obesity. A child with 110% of overweight, compared to 100%, would have 1.08 times the odds of having an additional family member with a history of obesity.



4.3 Family Members with History of Eating Disorders

Table 16: Overall Family Relationship Categories with History of Eating Disorders

Characteristic	N	N = 107
Mother	107	1 (0.9%)
Father	107	2 (1.9%)
Grandmother	107	4 (3.7%)
Grandfather	107	2 (1.9%)
Sister	107	2 (1.9%)
Brother	107	2 (1.9%)
Aunt	107	7 (6.5%)
Uncle	107	3 (2.8%)

¹ n (%)

4.3.1 mean (sd) number of family relationship categories

[1] 0.2149533

[1] 0.8358482

5 Sleep

Table 17: Sleep by Weight Status

Characteristic	N	HW, N = 41	OW, N = 29	OB, N = 37
Bedtime_28hr	97	21.62 (1.46) [19.00 - 25.00]	22.66 (1.96) [20.00 - 27.50]	22.60 (2.04) [19.00 - 28.00]
Unknown		4	4	2
Waketime_24hr	100	6.53 (1.13) [5.00 - 10.50]	7.08 (2.10) [5.50 - 15.50]	7.09 (2.12) [5.00 - 14.00]
Unknown		2	4	1
Bedtime_Recommendation	97			
N		15 (41%)	15 (60%)	24 (69%)
Y		22 (59%)	10 (40%)	11 (31%)
Unknown		4	4	2
Sleep_Recommendation	99			
N		17 (44%)	13 (52%)	17 (49%)
Y		22 (56%)	12 (48%)	18 (51%)
Unknown		2	4	2
Bed_hr	99	8.68 (1.88) [2.00 - 11.50]	8.48 (1.70) [5.00 - 12.00]	8.21 (1.30) [6.00 - 11.00]
Unknown		2	4	2
CSHQ_BedtimeResit	94	7.77 (2.41) [5.00 - 13.00]	6.91 (1.83) [5.00 - 11.00]	7.19 (2.08) [5.00 - 13.00]
Unknown		6	6	1
CSHQ_SleepOnsetDelay	100	1.51 (0.68) [1.00 - 3.00]	1.62 (0.85) [1.00 - 3.00]	2.09 (0.92) [1.00 - 3.00]
Unknown		2	3	2
CSHQ_SleepDuration	99	4.55 (1.62) [3.00 - 8.00]	4.52 (1.85) [3.00 - 9.00]	4.89 (1.70) [3.00 - 9.00]
Unknown		3	4	1
CSHQ_SleepAnxiety	98	6.13 (2.51) [4.00 - 12.00]	5.76 (2.52) [4.00 - 12.00]	5.14 (2.10) [4.00 - 11.00]
Unknown		3	4	2
CSHQ_NightWaking_no16	96	2.89 (1.41) [2.00 - 6.00]	2.75 (0.99) [2.00 - 6.00]	3.09 (1.29) [2.00 - 6.00]
Unknown		3	5	3
CSHQ_Parasomnias	93	8.67 (2.33) [7.00 - 19.00]	8.58 (1.50) [7.00 - 12.00]	9.21 (2.61) [7.00 - 19.00]
Unknown		5	5	4
CSHQ_SleepDisorderBreathing	91	3.35 (1.11) [3.00 - 7.00]	3.65 (1.34) [3.00 - 8.00]	4.52 (1.88) [3.00 - 9.00]
Unknown		4	6	6
CSHQ_DaytimeSleepiness	91	13.14 (4.09) [6.00 - 22.00]	11.67 (3.31) [7.00 - 18.00]	12.52 (2.80) [6.00 - 19.00]
Unknown		5	5	6

¹ Mean (SD) [Range]; n (%)

Note: Bedtime_28hr was used to be able to look at bedtime continuously. Since the latest reported bedtime was 4am, the day was extended to 28 hours to so midnight = 24.00 and 4am = 28.00.

5.1 Overall Averages

Table 18: Overall average sleep

	mean	median	sd
Bed_hr	8.465	8.5	1.646
Bedtime_28hr	22.242	22.0	1.864
Waketime	6.869	6.5	1.794

5.2 Association Between Sleep Sub-Scales and Percent of Overweight Cutoff

5.2.1 Correlation Matrix - Sleep

Table 19: Correlations between sleep percent of overweight

	IOTF_pOWcutoff	Bed_hr	Bedtime_28hr	Waketime_24hr
IOTF_pOWcutoff				
Bed_hr	-0.09			
Bedtime_28hr	0.13	-0.48*		
Waketime_24hr	0.06	0.31*	0.63*	

Table 20: Correlations between sleep percent of overweight

	IOTF_pOWcutoff	Bed_hr	Bedtime_28hr	Waketime_24hr
IOTF_pOWcutoff				
Bed_hr	0.392			
Bedtime_28hr	0.192	0		
Waketime_24hr	0.584	0.002	0	

Percent of overweight was not associated with bedtime, waketime, or total sleep.

5.2.2 t-tests for recommendation categories - Sleep

Table 21: Percent of Overweight by Sleep by Recommendations

	No	Yes	t	df	pvalue	sig
Sleep Duration Rec	116.35	112.56	0.50	96.13	0.6154	
Bedtime Rec	120.90	108.16	1.68	85.20	0.0975	.

5.2.3 Correlation Matrix - Sleep Behavior

Table 22: Correlations between sleep subscales and percent of overweight

	IOTF_pOWcutoff	BedResit	OnsetDelay	Duration	Anxiety	NightWaking	Parasomnias	Total
IOTF_pOWcutoff								
BedResit	-0.13							
OnsetDelay	0.31*	0.13						
Duration	0.04	0.41*	0.4*					
Anxiety	-0.18	0.62*	-0.1	0.02				
NightWaking	0.05	0.21	0.05	0.29*	0.23*			
Parasomnias	0.04	0.17	0.02	0.04	0.12	0.41*		
DisorderBreathing	0.37*	0.19	0.03	0.1	0.13	0.4*	0.43*	
DaySleepiness	-0.04	0.28*	0.16	0.21	0.28*	0.05	0.18	0
Total	0.1	0.63*	0.29*	0.47*	0.52*	0.47*	0.52*	0

Table 23: Correlations between sleep subscales and percent of overweight

	IOTF_pOWcutoff	BedResit	OnsetDelay	Duration	Anxiety	NightWaking	Parasomnias	I
IOTF_pOWcutoff								
BedResit	0.207							
OnsetDelay	0.002	0.227						
Duration	0.679	0	0					
Anxiety	0.07	0	0.322	0.851				
NightWaking	0.61	0.051	0.621	0.005	0.025			
Parasomnias	0.692	0.108	0.833	0.693	0.256	0		
DisorderBreathing	0	0.074	0.785	0.344	0.222	0	0	
DaySleepiness	0.676	0.011	0.127	0.052	0.008	0.622	0.101	0
Total	0.401	0	0.013	0	0	0	0	0

Examining correlations between sleep sub-scales and percent of overweight cutoff reveals greater percent of overweight cutoff was associated with greater parent reported sleep onset delay and sleep disordered breathing.

5.2.4 Sensitivity Tests

Table 24: Linear Model: Sleep Onset Delay - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	se	t	p	
(Intercept)	0.076	0.683	0.111	0.912	
Month_AED25,000 - 55,000 AED	-0.031	0.205	-0.149	0.882	
Month_AED55,000 - 75,000 AED	-0.082	0.390	-0.210	0.834	
Month_AED> 75,000 AED	-0.122	0.409	-0.299	0.766	
Mother_ed	0.008	0.029	0.261	0.795	
Age_yr	0.073	0.036	2.010	0.048	*
sexM	0.128	0.184	0.697	0.488	
IOTF_pOWcutoff	0.006	0.003	2.126	0.037	*

5.2.4.1 Sleep Onset Delay After controlling for family income, mother education, child age, and child sex, percent of overweight was positively associated with sleep onset delay such that a child with 110% of overweight, compared to 100% of overweight, would be expected to have a sleep onset delay score that was 0.06 points higher (range of scores = 0 - 3). Additionally, age was positively associated with sleep onset delay such that each year older, the expected sleep disordered breathing score would be 0.12 points higher (range of scores = 0 - 7).





Table 25: Linear Model: Sleep Disordered Breathing - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	se	t	p	
(Intercept)	3.197	1.243	2.573	0.012	
Month_AED25,000 - 55,000 AED	-0.628	0.394	-1.591	0.116	
Month_AED55,000 - 75,000 AED	-0.885	0.715	-1.238	0.220	
Month_AED> 75,000 AED	0.202	0.790	0.255	0.799	
Mother_ed	-0.019	0.053	-0.362	0.718	
Age_yr	-0.038	0.072	-0.522	0.603	
sexM	-0.055	0.354	-0.154	0.878	
IOTF_pOWcutoff	0.016	0.005	3.255	0.002	**

5.2.4.2 Sleep Disordered Breathing After controlling for family income, mother education, child age, and child sex, percent of overweight was positively associated with sleep disordered breathing such that a child with 110% of overweight, compared to 100% of overweight, would be expected to have a sleep disordered breathing score that was 0.16 points higher(range of scores = 3 - 9).

Association between Percent of Overweight Cutoff and Sleep Disord



6 Strengths and Difficulties Questionnaire

Table 26: Strengths and Difficulties by Weight Status

Characteristic	N	HW, N = 41	OW, N = 29	OB, N = 37
SDQ_EmotionProb_raw	104	3.05 [0.00 - 8.00]	3.00 [0.00 - 7.00]	3.43 [0.00 - 9.00]
Unknown		2	1	0
SDQ_ConductProb_raw	102	2.00 [0.00 - 7.00]	1.68 [0.00 - 5.00]	2.24 [0.00 - 6.00]
Unknown		1	1	3
SDQ_HyperactiveProb_raw	103	3.65 [0.00 - 9.00]	3.07 [0.00 - 10.00]	3.46 [0.00 - 10.00]
Unknown		2	2	0
SDQ_PeerProb_raw	102	2.71 [0.00 - 7.00]	2.97 [0.00 - 5.00]	3.46 [1.00 - 6.00]
Unknown		3	0	2
SDQ_Prosocial_raw	102	8.47 [4.00 - 10.00]	8.82 [6.00 - 10.00]	8.11 [2.00 - 10.00]
Unknown		2	1	2
SDQ_TotalProb_raw	106	11.11 [0.00 - 29.00]	10.34 [4.00 - 20.00]	12.22 [3.00 - 23.00]
Unknown		1	0	0
SDQ_TotalProb_cat	93			
CloseToAverage		27 (73%)	17 (68%)	19 (61%)
High		5 (14%)	2 (8.0%)	5 (16%)
SlightlyRaised		3 (8.1%)	5 (20%)	4 (13%)
VeryHigh		2 (5.4%)	1 (4.0%)	3 (9.7%)
Unknown		4	4	6
SDQ_EmotionProb_cat	104			
CloseToAverage		24 (62%)	18 (64%)	20 (54%)
High		7 (18%)	5 (18%)	7 (19%)
SlightlyRaised		5 (13%)	3 (11%)	6 (16%)
VeryHigh		3 (7.7%)	2 (7.1%)	4 (11%)
Unknown		2	1	0
SDQ_ConductProb_cat	102			
CloseToAverage		26 (65%)	21 (75%)	22 (65%)
High		7 (18%)	4 (14%)	7 (21%)
SlightlyRaised		6 (15%)	3 (11%)	4 (12%)
VeryHigh		1 (2.5%)	0 (0%)	1 (2.9%)
Unknown		1	1	3
SDQ_HyperactivityProb_cat	103			
CloseToAverage		29 (74%)	24 (89%)	32 (86%)
High		1 (2.6%)	0 (0%)	0 (0%)
SlightlyRaised		8 (21%)	1 (3.7%)	3 (8.1%)
VeryHigh		1 (2.6%)	2 (7.4%)	2 (5.4%)
Unknown		2	2	0
SDQ_PeerProb_cat	102			
CloseToAverage		17 (45%)	12 (41%)	11 (31%)
High		10 (26%)	9 (31%)	11 (31%)
SlightlyRaised		7 (18%)	6 (21%)	7 (20%)
VeryHigh		4 (11%)	2 (6.9%)	6 (17%)
Unknown		3	0	2
SDQ_Prosocial_cat	102			
CloseToAverage		30 (77%)	22 (79%)	25 (71%)
Low		2 (5.1%)	2 (7.1%)	1 (2.9%)
SlightlyLowered		5 (13%)	4 (14%)	6 (17%)
VeryLow		2 (5.1%)	0 (0%)	3 (8.6%)
Unknown		2	1	2

¹ Mean [Range]; n (%)

There were no differences by weight status.

Table 27: Strengths and Difficulties

Characteristic	N	N = 107
SDQ_TotalProb_cat	93	
CloseToAverage		63 (68%)
High		12 (13%)
SlightlyRaised		12 (13%)
VeryHigh		6 (6.5%)
Unknown		14
SDQ_EmotionProb_cat	104	
CloseToAverage		62 (60%)
High		19 (18%)
SlightlyRaised		14 (13%)
VeryHigh		9 (8.7%)
Unknown		3
SDQ_ConductProb_cat	102	
CloseToAverage		69 (68%)
High		18 (18%)
SlightlyRaised		13 (13%)
VeryHigh		2 (2.0%)
Unknown		5
SDQ_HyperactivityProb_cat	103	
CloseToAverage		85 (83%)
High		1 (1.0%)
SlightlyRaised		12 (12%)
VeryHigh		5 (4.9%)
Unknown		4
SDQ_PeerProb_cat	102	
CloseToAverage		40 (39%)
High		30 (29%)
SlightlyRaised		20 (20%)
VeryHigh		12 (12%)
Unknown		5
SDQ_Prosocial_cat	102	
CloseToAverage		77 (75%)
Low		5 (4.9%)
SlightlyLowered		15 (15%)
VeryLow		5 (4.9%)
Unknown		5

¹ n (%)

There were no differences by weight status.

6.1 Exploratory Sex x Percent of Overweight Models

6.1.1 Emotional Problems

Often complains of headaches

Many worries

Often unhappy, downhearted

Nervous or clingy in new situations

Many fears, easily scared

Table 28: Logistic Model: Emotional Problems (Elevated vs Not) - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	\hat{e}^b	se	$\hat{e}^{2.5}$ CI	$\hat{e}^{97.5}$ CI	z	p
(Intercept)	0.554	1.740	1.740	0.057	5.520800e+01	0.318	0.750
Month_AED25,000 - 55,000 AED	0.013	1.013	0.531	0.359	2.932000e+00	0.024	0.981
Month_AED55,000 - 75,000 AED	0.369	1.446	1.049	0.155	1.134600e+01	0.351	0.725
Month_AED> 75,000 AED	-16.635	0.000	1381.974	NA	3.271604e+40	-0.012	0.990
Mother_ed	-0.006	0.994	0.076	0.856	1.158000e+00	-0.085	0.932
Age_yr	-0.102	0.903	0.098	0.740	1.092000e+00	-1.032	0.302
sexM	-0.096	0.909	0.537	0.309	2.599000e+00	-0.178	0.858
IOTF_pOWcutoff_c100	0.011	1.011	0.010	0.992	1.032000e+00	1.155	0.248
sexM:IOTF_pOWcutoff_c100	-0.005	0.995	0.013	0.969	1.020000e+00	-0.420	0.675

6.2 Conduct Problems

Often has temper tantrums or hot tempers

Generally obedient - Reverse Scored

Often fights with other children

Often lies or cheats

Steals from home, school or elsewhere

Table 29: Logistic Model: Conduct Problems (Elevated vs Not) - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	\hat{e}^b	se	$\hat{e}^{2.5}$ CI	$\hat{e}^{97.5}$ CI	z	p
(Intercept)	-1.688	0.185	1.672	0.006	4.778	-1.010	0.313
Month_AED25,000 - 55,000 AED	-0.352	0.703	0.526	0.249	1.989	-0.669	0.503
Month_AED55,000 - 75,000 AED	0.324	1.383	0.988	0.190	10.184	0.328	0.743
Month_AED> 75,000 AED	-0.387	0.679	0.933	0.099	4.124	-0.415	0.678
Mother_ed	-0.005	0.995	0.074	0.857	1.152	-0.069	0.945
Age_yr	0.137	1.147	0.095	0.954	1.389	1.443	0.149
sexM	-0.792	0.453	0.525	0.155	1.235	-1.510	0.131
IOTF_pOWcutoff_c100	-0.007	0.993	0.010	0.974	1.012	-0.720	0.471
sexM:IOTF_pOWcutoff_c100	0.011	1.011	0.012	0.987	1.036	0.875	0.382

6.2.1 Hyperactivity

Restless, overactive

Constantly fidgeting or squirming

Easily distracted, concentration wanders

Thinks things out before acting - Reverse Scored

Sees tasks through to the end - Reverse Scored

Table 30: Logistic Model: Hyperactivity Problems (Elevated vs Not) - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	e ^b	se	e ^{2.5} CI	e ^{97.5} CI	z	p
(Intercept)	0.989	2.689	2.182	0.037	2.194150e+02	0.453	0.650
Month_AED25,000 - 55,000 AED	-0.817	0.442	0.670	0.115	1.660000e+00	-1.220	0.222
Month_AED55,000 - 75,000 AED	-16.836	0.000	1584.870	NA	4.622569e+45	-0.011	0.992
Month_AED> 75,000 AED	0.095	1.100	1.083	0.110	8.845000e+00	0.088	0.930
Mother_ed	0.011	1.011	0.096	0.840	1.233000e+00	0.110	0.912
Age_yr	-0.155	0.856	0.126	0.659	1.089000e+00	-1.234	0.217
sexM	-0.767	0.464	0.694	0.105	1.700000e+00	-1.106	0.269
IOTF_pOWcutoff_c100	-0.002	0.998	0.012	0.973	1.022000e+00	-0.156	0.876
sexM:IOTF_pOWcutoff_c100	0.008	1.008	0.017	0.973	1.042000e+00	0.453	0.651

6.2.2 Peer Problems

Rather solitary, tends to play alone

Has at least one good friend - Reverse Scored

Generally liked by other children - Reverse Scored

Picked on or bullied

Gets on better with adults than with other children

Table 31: Logistic Model: Peer Problems (Elevated vs Not) - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	e ^b	se	e ^{2.5} CI	e ^{97.5} CI	z	p
(Intercept)	0.048	1.049	1.806	0.030	38.161	0.026	0.979
Month_AED25,000 - 55,000 AED	0.212	1.236	0.546	0.421	3.650	0.388	0.698
Month_AED55,000 - 75,000 AED	2.186	8.896	1.294	0.931	215.972	1.689	0.091
Month_AED> 75,000 AED	0.176	1.193	0.938	0.184	7.743	0.188	0.851
Mother_ed	-0.151	0.860	0.087	0.716	1.011	-1.726	0.084
Age_yr	0.152	1.164	0.101	0.959	1.431	1.497	0.134
sexM	0.329	1.390	0.520	0.509	3.971	0.633	0.527
IOTF_pOWcutoff_c100	0.024	1.024	0.012	1.002	1.050	2.047	0.041
sexM:IOTF_pOWcutoff_c100	-0.030	0.970	0.014	0.942	0.997	-2.098	0.036

When examining odds of experiencing elevated peer problems, controlling for family income, mother education, child age, and child sex, percent of overweight, there was a significant sex x percent of overweight interaction.

sex = F:

IOTF_pOWcutoff_c100	IOTF_pOWcutoff_c100.trend	SE	df	asympt.LCL	asympt.UCL
17	0.02370	0.01158	Inf	0.00101	0.0464

```
sex = M:
  IOTF_pOWcutoff_c100 IOTF_pOWcutoff_c100.trend      SE  df asymp.LCL asymp.UCL
                17                -0.00632 0.00867 Inf   -0.02332    0.0107
```

Results are averaged over the levels of: Month_AED
Confidence level used: 0.95

For females, a female child with 110% of overweight cutoff would have 1.27 times the odds of experiencing elevate peer problems than a female child with 100% of overweight cutoff. There was no association between percent of overweight cutoff and odds of peer problems in males.

6.3 Prosocial

Considerate of other people's feelings

Shares readily with other children

Helpful if someone is hurt

Kind to younger children

Often volunteers to help others

Table 32: Logistic Model: Prosocial Problems (Low vs Not) - SES category + Maternal Education + Age + Sex + pOWcutoff

	b	e^b	se	$e^{2.5}$ CI	$e^{97.5}$ CI	z	p
(Intercept)	-1.501	0.223	2.020	0.004	11.541	-0.743	0.457
Month_AED25,000 - 55,000 AED	0.018	1.018	0.647	0.292	3.825	0.027	0.978
Month_AED55,000 - 75,000 AED	-0.611	0.543	1.413	0.018	6.626	-0.433	0.665
Month_AED > 75,000 AED	1.766	5.849	1.048	0.772	50.723	1.686	0.092 .
Mother_ed	-0.037	0.964	0.085	0.810	1.139	-0.435	0.664
Age_yr	0.072	1.075	0.117	0.855	1.359	0.615	0.538
sexM	-1.479	0.228	0.772	0.041	0.908	-1.916	0.055 .
IOTF_pOWcutoff_c100	-0.004	0.996	0.011	0.973	1.017	-0.384	0.701
sexM:IOTF_pOWcutoff_c100	0.029	1.029	0.015	1.001	1.063	1.914	0.056 .

7 Extra Tables by Sex

7.1 Participant Characteristics

Table 33: Demographic Characteristics by Sex

Characteristic	N	F	M	t-test	chi/fisher
Age_yr	107	12.79 (2.84) [7.31 - 17.84]	12.70 (2.57) [8.04 - 17.54]	0.875	
BMI	107	24.85 (7.94) [12.71 - 47.60]	25.70 (10.34) [13.60 - 55.52]	0.646	
IOTF_pOWcutoff	107	112.22 (31.68) [67.29 - 193.18]	117.71 (43.37) [63.95 - 239.00]	0.471	
IOTF_WeightStatus	107				0.454
Thinness2		1 (1.6%)	2 (4.3%)		
Thinness1		5 (8.2%)	3 (6.5%)		
HW		18 (30%)	12 (26%)		
Overweight		14 (23%)	15 (33%)		
Obese		11 (18%)	3 (6.5%)		
MorbidlyObese		12 (20%)	11 (24%)		
Father_ed	102	13.08 (3.79) [0.00 - 18.00]	12.48 (3.14) [6.00 - 18.00]	0.384	
Unknown		3	2		
Mother_ed	99	13.14 (3.56) [3.00 - 18.00]	13.24 (3.04) [0.00 - 18.00]	0.884	
Unknown		4	4		
Month_AED	99				0.745
<25,000 AED		15 (27%)	16 (37%)		
25,000 - 55,000 AED		32 (57%)	21 (49%)		
55,000 - 75,000 AED		4 (7.1%)	2 (4.7%)		
> 75,000 AED		5 (8.9%)	4 (9.3%)		
Unknown		5	3		
DadNationality	101				1.00
Emirati		58 (97%)	40 (98%)		
Omani		1 (1.7%)	0 (0%)		
Yemeni		1 (1.7%)	1 (2.4%)		
Unknown		1	5		
MomNationality	104				0.526
Emirati		55 (92%)	41 (93%)		
Omani		1 (1.7%)	0 (0%)		
Yemeni		0 (0%)	1 (2.3%)		
Moroccan		2 (3.3%)	0 (0%)		
Egyptian		2 (3.3%)	1 (2.3%)		
Bahrani		0 (0%)	1 (2.3%)		
Unknown		1	2		

¹ Mean (SD) [Range]; n (%)

There were no differences by sex with the exception of females having a higher hip-to-waist ratio, which would be expected for this age range.

7.2 Medical Comorbidities

Table 34: Medical Comorbidities by Weight Status

Characteristic	N	F	M	t-test	chi/fisher
nComorbid	107	2.25 (1.07) [0.00 - 4.00]	2.02 (0.98) [0.00 - 4.00]	0.263	
VitDdeficiency	107				1.00
Y		56 (92%)	42 (91%)		
N		5 (8.2%)	4 (8.7%)		
Anemia	40				0.424
Iron Deficiency Anemia (ID)		13 (54%)	8 (50%)		
Thalassemia Minor (TM)		2 (8.3%)	1 (6.2%)		
G6PD Deficiency		0 (0%)	1 (6.2%)		
ID + TM		0 (0%)	2 (12%)		
ID + G6PD Deficiency		1 (4.2%)	0 (0%)		
Unspecified Anemia		8 (33%)	4 (25%)		
Unknown		37	30		
Hyperlipidemia	12				1.00
Hyperlipidemia		8 (89%)	3 (100%)		
Hyperlipidemia - Mixed		1 (11%)	0 (0%)		
Unknown		52	43		
ThyroidConditions	25				1.00
Abnormal Function		10 (56%)	4 (57%)		
Autoimmune Thyroiditis		3 (17%)	2 (29%)		
Autoimmune Hypothyroidism		0 (0%)	0 (0%)		
Unspecified Hypothyroidism		4 (22%)	1 (14%)		
Goiter		1 (5.6%)	0 (0%)		
Unknown		43	39		
GlycemicStatus	28				0.678
Impaired Fasting Glucose		9 (64%)	10 (71%)		
Impaired Glucose Tolerance Test		3 (21%)	4 (29%)		
Type-1 Diabetes		2 (14%)	0 (0%)		
Unknown		47	32		
Acanthosis Nigricans	8	6 (100%)	2 (100%)		
Unknown		55	44		
Hypertension	3				
Essential Primary Hypertension		1 (100%)	1 (50%)		
High Blood Pressure		0 (0%)	1 (50%)		
Unknown		60	44		
Metabolic Syndrome	2	1 (100%)	1 (100%)		
Unknown		60	45		
Growth.Stature	10				1.00
Failure To Thrive (FT)		0 (0%)	1 (17%)		
Growth Hormone Deficiency		0 (0%)	1 (17%)		
Short Stature		3 (75%)	3 (50%)		
FT + ShortStature + Underweight		1 (25%)	0 (0%)		
Short Stature + Precocious Puberty		0 (0%)	1 (17%)		
Unknown		57	40		
PCOS	4				
PCOS		2 (50%)	0 (NA%)		
Hirsutism		1 (25%)	0 (NA%)		
Hirsutism + Unspecified Ovarian Cysts		1 (25%)	0 (NA%)		
Unknown		57	46		

¹ Mean (SD) [Range]; n (%)

Presence of different co-morbidities did not differ by sex, nor did the number of co-morbidities

7.3 Family History

Table 35: Family History by Sex

Characteristic	N	F	M	t-test	chi/fisher
Fam_OB_YN	101				0.895
yes		41 (71%)	29 (67%)		
no		17 (29%)	14 (33%)		
Unknown		3	3		
nFam_Obesity	107	1.80 (1.80) [0.00 - 7.00]	1.80 (1.71) [0.00 - 5.00]	1.00	
Mother	107	11 (18%)	7 (15%)		
Father	107	9 (15%)	12 (26%)		
Grandmother	107	19 (31%)	14 (30%)		
Grandfather	107	4 (6.6%)	2 (4.3%)		
Sister	107	10 (16%)	6 (13%)		
Brother	107	15 (25%)	10 (22%)		
Aunt	107	26 (43%)	19 (41%)		
Uncle	107	16 (26%)	13 (28%)		
Fam_ED_YN	96				0.517
yes		5 (8.9%)	6 (15%)		
no		51 (91%)	34 (85%)		
Unknown		5	6		
nFam_EatingDisorder	107	0.21 (0.97) [0.00 - 7.00]	0.22 (0.63) [0.00 - 3.00]	0.98	
Mother	107	1 (1.6%)	0 (0%)		
Father	107	1 (1.6%)	1 (2.2%)		
Grandmother	107	2 (3.3%)	2 (4.3%)		
Grandfather	107	2 (3.3%)	0 (0%)		
Sister	107	2 (3.3%)	0 (0%)		
Brother	107	1 (1.6%)	1 (2.2%)		
Aunt	107	3 (4.9%)	4 (8.7%)		
Uncle	107	1 (1.6%)	2 (4.3%)		

¹ n (%); Mean (SD) [Range]

There were no differences by sex.

7.4 Sleep

Table 36: Sleep by Sex

Characteristic	N	F	M	t-test	chi/fisher
Bedtime_28hr	97	22.48 (1.98) [19.00 - 28.00]	21.91 (1.66) [19.00 - 25.50]	0.129	
Unknown		5	5		
Waketime_24hr	100	7.22 (2.07) [5.00 - 15.50]	6.39 (1.19) [5.00 - 11.50]	0.014	
Unknown		3	4		
Bedtime_Recommendation	97				0.169
N		35 (62%)	19 (46%)		
Y		21 (38%)	22 (54%)		
Unknown		5	5		
Sleep_Recommendation	99				0.558
N		29 (51%)	18 (43%)		
Y		28 (49%)	24 (57%)		
Unknown		4	4		
Bed_hr	99	8.55 (1.54) [6.00 - 12.00]	8.35 (1.80) [2.00 - 11.50]	0.548	
Unknown		4	4		
CSHQ_BedtimeResit	94	7.72 (2.10) [5.00 - 13.00]	6.85 (2.17) [5.00 - 13.00]	0.056	
Unknown		8	5		
CSHQ_SleepOnsetDelay	100	1.69 (0.86) [1.00 - 3.00]	1.81 (0.83) [1.00 - 3.00]	0.486	
Unknown		3	4		
CSHQ_SleepDuration	99	4.86 (1.62) [3.00 - 9.00]	4.40 (1.80) [3.00 - 9.00]	0.198	
Unknown		4	4		
CSHQ_SleepAnxiety	98	6.02 (2.58) [4.00 - 12.00]	5.24 (2.05) [4.00 - 11.00]	0.099	
Unknown		5	4		
CSHQ_NightWaking_no16	96	3.11 (1.36) [2.00 - 6.00]	2.68 (1.11) [2.00 - 6.00]	0.094	
Unknown		6	5		
CSHQ_Parasomnias	93	8.75 (2.26) [7.00 - 19.00]	8.97 (2.26) [7.00 - 19.00]	0.634	
Unknown		6	8		
CSHQ_SleepDisorderBreathing	91	3.75 (1.51) [3.00 - 9.00]	3.92 (1.60) [3.00 - 8.00]	0.601	
Unknown		9	7		
CSHQ_DaytimeSleepiness	91	12.64 (3.35) [6.00 - 18.00]	12.39 (3.74) [6.00 - 22.00]	0.747	
Unknown		8	8		

¹ Mean (SD) [Range]; n (%)

There were no differences by sex.

7.5 Strengths and Difficulties Questionnaire

Table 37: Strengths and Difficulty by Sex

Characteristic	N	F	M	t-test	chi/fisher
SDQ_EmotionProb_raw	104	3.54 (2.17) [0.00 - 9.00]	2.69 (2.22) [0.00 - 8.00]	0.053	
Unknown		2	1		
SDQ_ConductProb_raw	102	2.00 (1.66) [0.00 - 6.00]	1.98 (1.71) [0.00 - 7.00]	0.945	
Unknown		2	3		
SDQ_HyperactiveProb_raw	103	3.12 (2.31) [0.00 - 10.00]	3.85 (2.37) [0.00 - 10.00]	0.120	
Unknown		2	2		
SDQ_PeerProb_raw	102	2.85 (1.42) [0.00 - 5.00]	3.30 (1.49) [0.00 - 7.00]	0.124	
Unknown		2	3		
SDQ_Prosocial_raw	102	8.38 (1.46) [4.00 - 10.00]	8.53 (1.93) [2.00 - 10.00]	0.659	
Unknown		3	2		
SDQ_TotalProb_raw	106	11.13 (5.05) [0.00 - 23.00]	11.50 (5.65) [3.00 - 29.00]	0.729	
Unknown		0	1		
SDQ_TotalProb_cat	93				0.411
CloseToAverage		37 (71%)	26 (63%)		
High		8 (15%)	4 (9.8%)		
SlightlyRaised		5 (9.6%)	7 (17%)		
VeryHigh		2 (3.8%)	4 (9.8%)		
Unknown		9	5		
SDQ_EmotionProb_cat	104				0.347
CloseToAverage		31 (53%)	31 (69%)		
High		13 (22%)	6 (13%)		
SlightlyRaised		10 (17%)	4 (8.9%)		
VeryHigh		5 (8.5%)	4 (8.9%)		
Unknown		2	1		
SDQ_ConductProb_cat	102				0.98
CloseToAverage		40 (68%)	29 (67%)		
High		11 (19%)	7 (16%)		
SlightlyRaised		7 (12%)	6 (14%)		
VeryHigh		1 (1.7%)	1 (2.3%)		
Unknown		2	3		
SDQ_HyperactivityProb_cat	103				0.704
CloseToAverage		48 (81%)	37 (84%)		
High		0 (0%)	1 (2.3%)		
SlightlyRaised		8 (14%)	4 (9.1%)		
VeryHigh		3 (5.1%)	2 (4.5%)		
Unknown		2	2		
SDQ_PeerProb_cat	102				0.312
CloseToAverage		23 (39%)	17 (40%)		
High		19 (32%)	11 (26%)		
SlightlyRaised		13 (22%)	7 (16%)		
VeryHigh		4 (6.8%)	8 (19%)		
Unknown		2	3		
SDQ_Prosocial_cat	102				0.482
CloseToAverage		42 (72%)	35 (80%)		
Low		3 (5.2%)	2 (4.5%)		
SlightlyLowered		11 (19%)	4 (9.1%)		
VeryLow		2 (3.4%)	3 (6.8%)		
Unknown		3	2		

¹ Mean (SD) [Range]; n (%)

There were no differences by sex.

\end{document}