


# Lecture Notes to a course on Lie Algebras

Winter 2018, Technion IIT

*Lectures by Amos Nevo*  
*Typed by Elad Tzorani*



Last updated January 20, 2019

# Contents

<b>Preface</b>	<b>iii</b>
Technicalities . . . . .	iii
Course Literature . . . . .	iii
 <b>I Lie Algebras</b>	 <b>1</b>
<b>1 Preliminaries</b>	<b>2</b>
1.1 Basic definitions . . . . .	2
1.2 Structure constants . . . . .	4
1.3 Linear representations . . . . .	5
1.4 Sub-algebras and ideals . . . . .	6
<b>2 Structure of Lie algebras</b>	<b>9</b>
2.1 Nilpotent Lie algebras . . . . .	9
2.1.1 Flags . . . . .	12
2.2 Solvable Lie algebras . . . . .	13
<b>3 Jordan-Chevalley decomposition</b>	<b>17</b>
3.1 The Chinese remainder theorem . . . . .	17
3.2 Decomposition of vector spaces . . . . .	17
<b>4 Cartan's criterion for semi-simplicity</b>	<b>21</b>
4.1 Preliminary results . . . . .	21
4.2 Cartan's criterion . . . . .	22
<b>5 Killing form</b>	<b>23</b>
5.1 Some properties of the Killing form . . . . .	23
<b>6 Derivations of simple algebras</b>	<b>27</b>
<b>7 Toral subalgebras</b>	<b>29</b>
7.1 Orthogonality of root systems . . . . .	31
7.2 Rationality properties of the root system . . . . .	34
7.3 Geometry of Euclidean root systems . . . . .	39
7.4 Pairs of roots in a root system . . . . .	39
7.5 The action of the reflection $\sigma_\alpha$ on the $\alpha$ -chain through $\beta$ . . . . .	40
<b>8 The classification of simple Lie algebras over an algebraically closed field of characteristic 0</b>	<b>43</b>
8.1 Coxeter graphs and Dynkin diagrams . . . . .	43
8.2 Classification of connected Coxeter graphs . . . . .	44

# Preface

## Technicalities

These aren't formal notes related to the course and henceforward there is *absolutely no guarantee* that the recorded material is in correspondence with the course expectations, or that these notes lack any mistakes.

In fact, there probably are mistakes in the notes! I would highly appreciate if any comments or corrections were sent to me via email at [tzorani.elad@gmail.com](mailto:tzorani.elad@gmail.com).

Elad Tzorani.

## Course Literature

The recommended course literature is as follows.

**Humphreys, James E.:** Introduction to Lie algebras and representation theory.

**Jacobson, Nathan:** Lie algebras. New York, 1962.



**Part I**

**Lie Algebras**

# Chapter 1

## Preliminaries

The course will be entirely algebraic, with possibly few examples from analysis.

This will allow us to discuss issues regarding the algebraic properties of Lie algebras. We might be interested in infinite-dimensional Lie algebras, but in this course we discuss only finite-dimensional algebras. In this course one of our main goals is a classification theorem for simple Lie algebras. We assume knowledge in linear algebras and specifically bilinear forms.

### 1.1 Basic definitions

Let  $\mathbb{F}$  be a field, and  $V$  a finite-dimensional vector-space over  $\mathbb{F}$ .

**Definition 1.1.1.**  $V$  is a *generalised  $\mathbb{F}$ -algebra* if it comes with a map  $m: V \times V \rightarrow V$  which is bilinear.

$$\begin{aligned}m(v_1 + v_2, w) &= m(v_1, w) + m(v_2, w) \\m(v, w_1 + w_2) &= m(v, w_1) + m(v, w_2) \\m(av, bw) &= abm(v, w)\end{aligned}$$

**Example.** Let  $V$  be an associative algebra. Here  $m$  is an associative operation which is left and right distributive on addition in  $v$ . Equivalently: If we denote  $m(v, w) = v \odot w$  then

$$\begin{aligned}(v \odot w) \odot u &= v \odot (w \odot u) \\v \odot (u + w) &= v \odot u + v \odot w \\(u + w) \odot v &= u \odot v + w \odot v\end{aligned}$$

**Remark 1.1.2.** Here associativity means the following.

$$m(v, m(w, u)) = m(m(v, w), u)$$

**Examples.** 1. Every field  $k$  is an  $\mathbb{F}$ -algebra over any subfield  $\mathbb{F}$ .

2.  $M_n(k)$  is an  $\mathbb{F}$ -algebra.

3.  $P_n$ , polynomials over  $k$  of degree smaller or equal to  $n$ , is an  $\mathbb{F}$ -algebra.

**Definition 1.1.3.** A Lie algebra  $L$  over  $\mathbb{F}$  is an  $\mathbb{F}$ -algebra, so  $\exists m: L \times L \rightarrow L$ , which generally need not be associative, but instead satisfies the following *Jacobi identity*,

$$m(X, m(Y, Z)) + m(Z, m(X, Y)) + m(Y, m(Z, X)) = 0$$

and additionally, antisymmetry of the multiplication

$$m(X, Y) = -m(Y, X).$$

If  $\text{char}\mathbb{F} = 2$  we require  $m(X, X) = 0$ .

**Notation 1.1.4.** The "multiplication" in  $L$  is called *bracket*, and denoted  $m(X, Y) = [X, Y]$  ( $X$  bracket  $Y$ ).

**Remark 1.1.5.** In these terms we write the Jacobi identity as follows.

$$[X, [Y, Z]] + [Z, [X, Y]] + [Y, [Z, X]] = 0$$

**Definition 1.1.6.** A **Lie algebra**  $L$  is a vector space over  $\mathbb{F}$  with a bilinear map  $[\cdot, \cdot] : L \times L \rightarrow L$ , which is anti-symmetric and satisfies the Jacobi identity.

**Definition 1.1.7.** Given a Lie algebra  $L$ , a vector subspace  $L_0 \subseteq L$  is called a **Lie sub-algebra** if it is closed under brackets. I.e.

$$X, Y \in L_0 \implies [X, Y] \in L_0.$$

**Examples.** 1. *Abelian Lie algebras:* The bracket is the zero form.

$$\forall X, Y \in L: [X, Y] = 0$$

**Example.**  $\mathbb{F}$  is itself a Lie algebra as well as any  $\mathbb{F}$ -vector space  $V$  under the bracket

$$\forall u, v \in V: [u, v] = 0.$$

**Example.** Let  $A$  be any associative  $\mathbb{F}$ -algebra, and define on  $A$  *another* bilinear operation, namely

$$[a, b] = ab - ba.$$

This is called **the commutator of  $a$  and  $b$** . Then  $[\cdot, \cdot] : A \times A \rightarrow A$ .

**Exercise.** This bracket satisfies the Jacobi identity, and is anti-symmetric.

Given a solution to this exercise,  $(A, [\cdot, \cdot])$  is a Lie algebra.

In particular,  $M_n(k)$  is a Lie algebra under the bracket  $[A, B] = AB - BA$ . This algebra is *very important* and is denoted  $\mathfrak{gl}_n(k)$ .

**Exercise.** Consider the subspace

$$\{A \in \mathfrak{gl}_n(k) \mid \text{tr} A = 0\} \subseteq \mathfrak{gl}_n(k).$$

Is the subspace a Lie algebra? Yes! Since for any  $A, B \in \mathfrak{gl}_n(k)$  we have that  $\text{tr}(AB) = \text{tr}(BA)$ , we get that  $\text{tr}[A, B] = 0$ . The sub-Lie-algebra of zero-trace matrices is denoted  $\mathfrak{sl}_n(k)$ .

**Exercise (Lie algebras associated with bilinear forms).** Let  $V$  be a vector space over  $\mathbb{F}$ , and  $B : V \times V \rightarrow \mathbb{F}$  be a bilinear form. Assume  $B$  is anti-symmetric. Define

$$L_B = \{X \in \text{End}(V) \mid B(Xv, w) = -B(v, Xw)\}.$$

Check that  $L_B$  is a vector subspace of  $\text{End}(V)$ . Consider the bracket operation on  $\text{End}(V)$ , defined by  $[T, S] = TS - ST$ . Is  $L_B$  closed under brackets?

**Solution.** We compute as follows.

$$\begin{aligned} B([X, Y]v, w) &= B((XY - YX)v, w) \\ &= B(XYv, w) - B(YXv, w) \\ &= -B(Yv, Xw) + B(Xv, Yw) \\ &= B(v, YXw) - B(v, XYw) \\ &= B(v, (YX - XY)w) \\ &= -B(v, [X, Y]w) \end{aligned}$$

In conclusion,  $L_B$  is a sub-Lie-algebra of  $\text{End}(V)$ , the Lie algebra associated with the form  $B$ .

**Exercise.** Let  $S$  be a symmetric bilinear form, and let

$$L_S = \{X \in \text{End}(V) \mid S(Xv, w) = -S(v, Xw)\}.$$

Then again,  $L_S$  is a Lie sub-algebra.

**Examples (Sub-algebras of  $\mathfrak{gl}_n(\mathbb{F})$ ).** 1.

$$\mathfrak{t}(n, \mathbb{F}) = \left\{ \begin{pmatrix} a_{11} & & a_{i,j} \\ & \ddots & \\ 0 & & a_{nn} \end{pmatrix} \mid a_{ij} \in \mathbb{F} \right\}$$

is closed under the bracket operation, for if  $A, B \in \mathfrak{t}(n, \mathbb{F})$  then  $AB \in \mathfrak{t}(n, \mathbb{F})$  and so  $AB - BA \in \mathfrak{t}(n, \mathbb{F})$ .

2.

$$\mathfrak{n}(n, \mathbb{F}) = \left\{ \begin{pmatrix} 0 & & a_{i,j} \\ & \ddots & \\ 0 & & 0 \end{pmatrix} \mid a_{ij} \in \mathbb{F} \right\}$$

is a Lie sub-algebra of  $\mathfrak{t}(n, \mathbb{F})$ .

3.

$$\mathfrak{d}(n, \mathbb{F}) = \left\{ \begin{pmatrix} a_1 & & 0 \\ & \ddots & \\ 0 & & a_n \end{pmatrix} \mid a_i \in \mathbb{F} \right\}$$

an abelian sub-algebra.

## 1.2 Structure constants

Let  $L$  be a Lie algebra and let  $X_1, \dots, X_n$  be a basis of  $L$ . Then the bracket operation is completely determined by the structure constants with respect to the basis.

$$[X_i, X_j] = \sum_{k=1}^n c_k^{i,j} X_k$$

The **structure constants**  $c_k^{i,j}$  contain full information on the bracket operation of course. These satisfy two properties associated with anti-symmetry and the Jacobi identity of the brackets. The property associated to anti-symmetry is  $c_k^{i,j} = -c_k^{j,i}$ . The other property (associated to the Jacobi identity) is left as an **Exercise**.

**Example.**

$$\mathfrak{gl}_n(\mathbb{F}) = \text{span} \{E_{i,j} \mid 1 \leq i, j \leq n\}$$

In the basis  $E_{ij}$  the structure constants are very simple. We have the following.

$$[E_{i,j}, E_{k,l}] = \delta_{j,k} E_{i,l} - \delta_{l,i} E_{k,j}$$

Hence all the structures constants are 1 or  $-1$ .

**Definition 1.2.1.** Let  $L_1, L_2$  be Lie algebras. A **Lie algebra homomorphism** between  $L_1$  and  $L_2$  is a linear map  $T: L_1 \rightarrow L_2$  satisfying

$$T[X, Y] = [TX, TY].$$

**Definition 1.2.2.** Let  $L$  be a Lie algebra. A sub-space  $I \subseteq L$  is called a **Lie-ideal** of  $L$  if for all  $X \in L$  and  $Y \in I$ , we have that  $[X, Y] \in I$ . This is written also by

$$[L, I] = \text{span} \{[X, Y] \mid X \in L, Y \in I\} \subseteq I.$$

**Definition 1.2.3.** Let  $L$  be a Lie algebra and  $L_0 \subseteq L$  be a sub-space. The **Lie normaliser** of  $L_0$  is

$$N(L_0) = \{X \in L \mid [X, L_0] \subseteq L_0\}.$$

The **Lie centraliser** of  $L_0$  is

$$Z(L_0) = \{X \in L \mid [X, L_0] = 0\}.$$



**Definition 1.2.4.** Let  $L$  be a Lie algebra. If  $[X, Y] = 0$  one says that  $X$  and  $Y$  commute. We sometimes refer to the bracket as the commutator.

**Example.** Two sub-spaces  $L_1, L_2 \subseteq L$  of a Lie algebra commute if their commutators are zero. I.e.

$$[L_1, L_2] = 0.$$

**Remark 1.2.5.** Although we have linearity of the bracket, we do need to take the span in the above example. If we take  $X, X' \in L_1$  and  $Y, Y' \in L_2$  we can't always express  $[X, Y] + [X', Y']$  as a bracket of two elements, although it certainly is in the span.

## 1.3 Linear representations

**Definition 1.3.1.** A **linear representation** of a Lie algebra  $L$  over  $\mathbb{F}$  is a Lie-algebra homomorphism  $T: L \rightarrow \text{End}(V) \cong \mathfrak{gl}_n(\mathbb{F})$  where  $V$  is an  $n$ -dimensional vector space over  $\mathbb{F}$ .

**Remark 1.3.2.** The bracket operation on  $\text{End}(V)$  is the usual one, namely  $[A, B] = AB - BA$ .

Let us define another large collection of Lie algebras. First, let  $A$  be a generalised  $\mathbb{F}$ -algebra, and denote  $m(a, b) = a \odot b$ .

**Definition 1.3.3.** A **derivation** of the generalised algebra  $A$  is a linear map  $\delta: A \rightarrow A$  satisfying the following property.

$$\delta(a \odot b) = \delta(a) \odot b + a \odot \delta(b)$$

**Definition 1.3.4.**

$$\text{Der}(A) := \{\delta \in \text{End}(A) \mid \delta \text{ is a derivation.}\}$$

**Remark 1.3.5.**  $\text{Der}(A)$  is clearly a linear sub-space of  $\text{End}(A)$ . Now, if  $\delta_1$  and  $\delta_2$  are derivations,  $\delta_1 \circ \delta_2$  is *not* a derivation, usually. But,  $[\delta_1, \delta_2] = \delta_1 \circ \delta_2 - \delta_2 \circ \delta_1$  is in fact a derivation.

**Conclusion.**  $\text{Der}(A)$ , with the bracket inherited from  $\text{End}(A)$  is a Lie algebra.

*Proof.* We compute the following.

$$\begin{aligned} [\delta_1, \delta_2](a \odot b) &= (\delta_1 \circ \delta_2 - \delta_2 \circ \delta_1)(a \odot b) \\ &= \delta_1 \circ \delta_2(a \odot b) - \delta_2 \circ \delta_1(a \odot b) \\ &= \delta_1(\delta_2(a) \odot b + a \odot \delta_2(b)) - \delta_2(\delta_1(a) \odot b + a \odot \delta_1(b)) \\ &= \delta_1\delta_2(a) \odot b + \delta_2(a) \odot \delta_1(b) + \delta_1(a) \odot \delta_2(b) + a \odot \delta_1\delta_2(b) \\ &\quad - (\delta_2\delta_1(a) \odot b + \delta_1(a) \odot \delta_2(b) + \delta_2(a) \odot \delta_1(b) + a \odot \delta_2\delta_1(b)) \\ &= (\delta_1\delta_2 - \delta_2\delta_1)(a) \odot b + a \odot (\delta_1\delta_2 - \delta_2\delta_1)(b) \end{aligned}$$

■

**Example.** 1. If  $A$  is an associative algebra, then  $\text{Der}(A)$  is a Lie algebra,  $\text{Der}(A) \subseteq \text{End}(A)$ .  $\text{Der}(A)$  is a sub-Lie-algebra of  $\text{End}(A)$  under bracket of linear transformations.

2. A Lie algebra is a generalised algebra and so  $\text{Der}(L)$  is another Lie algebra.

**Fact 1.3.6 (important).** There is a very natural collection of derivations of any Lie algebras. For each  $x \in L$ , let us define a linear transformation denoted  $\text{ad}(x): L \rightarrow L$  (this stands for "adjoint") via  $\text{ad}(x)(y) = [x, y]$ . (This is linear from the bi-linearity of the bracket) In fact,  $\text{ad}(x)$  is a derivation of  $L$ . Namely,

$$\text{ad}(x)([y, z]) = [\text{ad}(x)y, z] + [y, \text{ad}(x)z].$$

Indeed,

$$\begin{aligned} \text{ad}(x)([y, z]) &= [x, [y, z]] \\ &= [\text{ad}(x)y, z] + [y, \text{ad}(x)z] \\ &= [[x, y], z] + [y, [x, z]] \end{aligned}$$

which is an identity as a consequence of the Jacobi identity.

**Conclusion.** The set  $\{\text{ad}(x) \mid x \in L\} \subseteq \text{Der}(L)$  is a sub-algebra. We have the map  $x \mapsto \text{ad}(x)$  which is obviously linear (from bi-linearity of the bracket). So,  $\text{ad}(L) := \{\text{ad}(x) \mid x \in L\}$  is a linear sub-space. In fact it is a Lie sub-algebra of  $\text{Der}(L)$ .

*Proof.* We have to show that  $[\text{ad}(x), \text{ad}(y)] = \text{ad}(x)\text{ad}(y) - \text{ad}(y)\text{ad}(x)$  is in the space  $\text{ad}(L)$ . But, actually  $[\text{ad}(x), \text{ad}(y)] = \text{ad}[x, y]$ , as the following proposition states.

**Proposition 1.3.7.**  $\text{ad}: L \rightarrow \text{Der}(L)$  is a Lie algebra homomorphism.

*Proof.* Let us compute.

$$\begin{aligned} [\text{ad}(x), \text{ad}(y)](z) &= \text{ad}(x)\text{ad}(y)(z) - \text{ad}(y)\text{ad}(x)(z) \\ &= [x, [y, z]] - [y, [x, z]] \\ &\stackrel{\star}{=} [[x, y], z] \\ &= \text{ad}[x, y](z) \end{aligned}$$

where the  $\star$  is given from the Jacobi identity. ■

In conclusion,  $\text{Der}(L)$  is a Lie sub-algebra of  $\text{End}(L)$  under bracket, and  $\text{ad}: L \rightarrow \text{Der}(L) \subseteq \text{End}(L)$  is a linear representation of the Lie algebra  $L$  with the image being  $\text{ad}(L) = \{\text{ad}(x) \mid x \in L\}$ . ■

**Example.** Given  $L_0 \subseteq L$  a sub-space. Then  $N(L_0) = \{x \mid [x, L_0] \subseteq L_0\}$  is the set of elements  $x$  such that the linear transformation  $\text{ad}(x)$  leaves the subspace  $L_0$  invariant.  $N_L(L_0)$  is a Lie sub-algebra, and if  $L_0$  is a Lie sub-algebra, then  $L_0$  is an ideal of  $N_L(L_0)$ .

**Example.** The condition  $[X, Y] = 0$  means  $Y \in \ker(\text{ad}(x))$  or equivalently  $x \in \ker(\text{ad}(y))$ . Therefore

$$\begin{aligned} Z(L_0) &= \{x \in L \mid [x, L_0] = 0\} \\ &= \{x \in L \mid L_0 \subseteq \ker(\text{ad}(x))\}. \end{aligned}$$

$Z(L_0)$  is a Lie sub-algebra of  $L$ , the Lie sub-algebra of elements commuting with every  $x \in L_0$ .

**Remark 1.3.8.** If  $L_0 \subseteq L$  is a Lie sub-algebra, then  $N(L_0)$  is the largest sub-algebra such that  $L_0$  is an ideal in it.

**Remark 1.3.9.**  $Z_L(L)$  is the center of  $L$ , and an ideal. Indeed, if  $z \in Z(L)$ , and  $x \in L$ , then  $\text{ad}[x, z] = \text{ad}x \text{ad}z - \text{ad}z \text{ad}x$  and  $L \subseteq \ker \text{ad}z$ , so  $L \subseteq \ker \text{ad}[x, z]$ , so  $[x, z] \in Z(L)$  and  $Z(L)$  is an ideal.

## 1.4 Sub-algebras and ideals

**Remark 1.4.1.** 1. If  $L_1$  and  $L_2$  are Lie sub-algebras, then  $L_1 + L_2$  generally is *not*!

2. Suppose  $I = L_1$  is an ideal and  $L_2$  a sub-algebra. Then  $I + L_2$  is a sub-algebra.

3. If  $L_1 = I$  and  $L_2 = J$  are ideals, then the Lie sub-algebra  $I + J$  is an ideal. Indeed  $[x, i] \in I$  and  $[x, j] \in J$  for all  $j$ , so  $[x, I + J] \subseteq I + J$ .

**Definition 1.4.2.** The *commutator* of two sub-algebras  $L_1, L_2$  is defined to be

$$\text{Span} \{[X, Y] \mid X \in L_1, Y \in L_2\}.$$

**Remark 1.4.3.** The commutator of two sub-algebras is *not* in general a sub-algebra. Generally  $[[X, Y], [X', Y']]$  isn't in  $[L_1, L_2]$  if  $X, X' \in L_1$  and  $Y, Y' \in L_2$ . Let

$$\sum_{i=1}^n [X_i, Y_i] \in [L_1, L_2]$$

and

$$\sum_{j=1}^m [X'_j, Y'_j] \in [L_1, L_2].$$

Then

$$\left[ \sum_{i=1}^n [X_i, Y_i] \sum_{j=1}^m [X'_j, Y'_j] \right] = \sum_{\substack{i=1 \\ j=1}}^n [ [X_i, Y_i], [X'_j, Y'_j] ]. \quad (1.1)$$

1. If  $L_1 = I$  is an ideal, then  $[I, L_2] \subseteq I$ , is a sub-space of  $I$ .
2. If  $L_1 = I$  and  $L_2 = J$  are ideals, then  $[I, J] \subseteq I \cap J$ , and it is an ideal of  $L$ . Equation 1.1 shows that  $[I, J]$  is indeed a sub-algebra. Now, let  $[i, j] \in [I, J]$ , and let  $x \in L$ . We should show that  $[x, [i, j]] \in [I, J]$  which is sufficient for the span. Now

$$[x, [i, j]] \stackrel{\text{Jacobi identity}}{=} [[x, i], j] + [i, [x, j]] = [i', j] + [i, j'] \in [I, J]$$

as required.

**Conclusion.**  $I + J$  and  $[I, J]$  are ideals if  $I$  and  $J$  are.

**Remark 1.4.4.** In general  $[I, J] \subseteq I \cap J$ , but the inclusion may be strict.

**Examples.** 1. Take  $L$  an abelian Lie algebra and  $I, J$  any two sub-spaces which are both sub-algebras, and ideals. Then  $[I, J] = 0$ , but  $I \cap J$  may be large.

2. Take  $L$  a Lie algebra of upper-triangular matrices, and  $I = J$  the ideal of strict upper-triangular matrices. Then  $[I, I]$  contains matrices that have zero entries in the diagonal above the main diagonal, hence  $[I, I] \subsetneq I \cap J = I$ .

**Definition 1.4.5.** If  $[I, J] = 0$ , we say that  $I$  and  $J$  *commute*.

**Remark 1.4.6.**  $L$  is an ideal of itself, so  $[L, L] = \text{Span}\{[X, Y] \mid X, Y \in L\}$  is also an ideal, *the commutator ideal of  $L$* .

**Definition 1.4.7.**  $L$  is *abelian* if  $[L, L] = 0$ .

**Definition 1.4.8.**  $L$  is *perfect* if  $[L, L] = L$ .

**Definition 1.4.9.**  $L$  is called a *simple Lie-algebra* if  $\dim L > 1$  and  $L$  has no non-trivial ideals.

**Exercise.** A simple Lie algebra is in particular perfect.

**Proposition 1.4.10.** If  $\varphi: L \rightarrow L'$  is a Lie-algebra homomorphism, then  $\ker \varphi$  is an ideal.

**Definition 1.4.11.** For any ideal  $I \triangleleft L$ , the factor vector space  $L/I = \{\ell + I \mid \ell \in L\}$  has a structure of a Lie algebra, given by the following.

$$[x + I, y + I]_{L/I} := [x, y]_L + I$$

**Remark 1.4.12.** The above is well defined since

$$[x + i, y + i'] = [x, y] + [i, y] + [x, i'] + [i, i'] \equiv [x, y] \pmod{I}.$$

The identities for Lie algebras follow immediately from those on  $L$ .

**Theorem 1.4.13 (1<sup>st</sup> homomorphism theorem).**

$$\begin{aligned} \pi: L &\rightarrow L/I \\ x &\mapsto x + I \end{aligned}$$

is a surjective Lie-algebra homomorphism, and  $\ker \pi = I$ .

**Theorem 1.4.14 (2<sup>nd</sup> homomorphism theorem).** If  $I$  and  $J$  are ideals of  $L$ , and  $I \subset J$ , then the map

$$\begin{aligned} \varphi: L/I &\rightarrow L/J \\ x + I &\mapsto x + J \end{aligned}$$

is a well-defined Lie-algebra epimorphism. We have from the first homomorphism theorem that

$$L/I / \ker \varphi \cong L/J$$

and

$$\ker \varphi = J/I$$

therefore

$$L/I / J/I \cong L/J.$$

**Theorem 1.4.15** (*3<sup>rd</sup> homomorphism theorem*). *Given any two ideals  $I, J$ , their intersection  $I \cap J$  is an ideal of  $L$  and we have a map*

$$\begin{aligned} \psi: I &\rightarrow I + J/J \\ i &\mapsto i + J \end{aligned}.$$

*This is a Lie-algebra homomorphism which is obviously surjective, with kernel  $I \cap J$ , hence*

$$I/I \cap J \cong I + J/J$$

*with Lie-algebra homomorphism induced by  $\psi$ .*

**Remark 1.4.16.** If  $L_0$  is an arbitrary Lie sub-algebra of  $L$ , and  $J \triangleleft L$ , then  $J \cap L_0 \triangleleft L_0$  and  $J \triangleleft L_0 + J$ , and the Lie algebras  $L_0 + J/J$  and  $L_0/L_0 \cap J$  are isomorphic under the canonical map  $\psi$ .

# Chapter 2

## Structure of Lie algebras

### 2.1 Nilpotent Lie algebras

**Definition 2.1.1.** The commutator ideal  $[L, L]$  is denoted  $L^{(1)}$ . Similarly we denote  $L^{(n)} = [L^{(n-1)}, L]$ , which is an ideal of  $L$ .

**Remark 2.1.2.** The above gives a descending chain

$$L = L^{(0)} \supseteq L^{(1)} \supseteq L^{(2)} \supseteq \dots$$

and since  $\dim L < \infty$ , this sequence has to stabilise. It is however possible that  $[L, L] = 0$ , if  $L$  is abelian, or that  $[L, L] = L$ , if  $L$  is perfect.

**Definition 2.1.3.** If  $L^{(n)} = 0$  for some  $n$ ,  $L$  is called a **nilpotent Lie algebra**. If  $L^{(n)} = 0$  and  $L^{(n-1)} \neq 0$ , we call  $n - 1$  the **index of nilpotency**.

**Note 2.1.4.** In some books  $n$  itself is called the index of nilpotency.

**Definition 2.1.5.** The sequence of ideals  $L^{(n)}$  is called **the descending central series** of  $L$ .

**Remark 2.1.6.**  $L^{(k)} \triangleleft L$  and hence  $L^{(l)} \triangleleft L^{(k-1)}$ . Also  $L^{(k-1)} / L^{(k)}$  is an *abelian* algebra since  $L^{(k)} = [L^{(k-1)}, L] \supseteq [L^{(k-1)}, L^{(k-1)}]$  and in general an ideal  $I \triangleleft M$  is such that  $M/I$  is abelian if and only if  $I \supseteq [M, M]$ .

**Proposition 2.1.7.** Let  $\varphi: L_1 \rightarrow L_2$  be an epimorphism of Lie algebras. Then  $\varphi(L_1^{(n)}) = L_2^{(n)}$ .

**Exercise.** Prove the above proposition. For  $n = 1$ , we have  $\varphi([L_1, L_1]) \subseteq [L_2, L_2]$ , but in fact equality holds (**Exercise!**). Similarly prove for any  $n \in \mathbb{N}$ .

**Proposition 2.1.8.** Let  $L$  be a nilpotent Lie algebra.

1. Every Lie sub-algebra and every factor Lie algebra are also nilpotent.
2. For  $M$  a Lie algebra, if  $M / Z(M)$  is nilpotent, so is  $M$ .
3.  $Z(L) \neq 0$ .

*Proof.* 1. **Sub-algebras:** If  $L_0 \subset L$  is a Lie sub-algebra, then clearly  $L_0^{(k)} \subseteq L^{(k)}$ . So if  $L^{(n)} = 0$  then  $L_0^{(n)} = 0$  and the index of nilpotency of  $L_0$  is bounded by that of  $L$ .

**Factor algebras:** Let  $\bar{L} = \varphi(L) = L/I$  be an epimorphic image of  $L$ . Then  $L^{(k)} = \varphi(L^{(k)})$ , so if  $L^{(k)} = 0$ , then  $\bar{L}^{(k)} = 0$ . We similarly have a bound on the nilpotency index of the factor algebra.

2. Suppose  $\bar{L} = L/Z$  is nilpotent. Then  $\bar{L}^{(n)} = \bar{0}$  for some  $n$ . So

$$\varphi(L^{(n)}) = \bar{L}^{(n)} = \bar{0}.$$

Then

$$\varphi(L^{(n)}) = \bar{L}^{(n)} = 0$$

and therefore  $L^{(n)} \subseteq Z(L) = \ker \varphi$ . Therefore  $[L^{(n)}, L] \in [Z(L), L] = 0$ . So  $L^{(n+1)} = 0$ , and so the index of nilpotency may increase by 1.

3. By definition,

$$L^{(0)} \supseteq L^{(1)} \supseteq \dots \supseteq L^{(n-1)} \supsetneq L^{(n)} = 0$$

for some  $n \in \mathbb{N}$ . Now  $[L^{(n-1)}, L] = L^{(n)} = 0$ , so certainly  $L^{(n-1)} \subseteq Z(L)$  and  $Z(L) \neq 0$ . ■

**Exercise.**

$$\mathfrak{n}(n, \mathbb{F}) = \left\{ \begin{pmatrix} 0 & & a_{i,j} \\ & \ddots & \\ 0 & & 0 \end{pmatrix} \mid a_{ij} \in \mathbb{F} \right\}$$

is a nilpotent Lie sub-algebra of  $M_n(\mathbb{F})$ .

**Example.** In  $\mathfrak{n}(2, \mathbb{F})$ , the commutator of any two elements is zero.

$$\left[ \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & y \\ 0 & 0 \end{pmatrix} \right] = 0$$

Therefore  $\mathfrak{n}(3, \mathbb{F})$  is a one-dimensional abelian algebra. For  $\mathfrak{n}(3, \mathbb{F})$ , the commutator of an element  $\begin{pmatrix} 0 & 0 & x \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

with any other element is zero. However,

$$\left[ \begin{pmatrix} 0 & u & w \\ 0 & 0 & v \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & x & y \\ 0 & 0 & z \\ 0 & 0 & 0 \end{pmatrix} \right] = \begin{pmatrix} 0 & 0 & uz - vx \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Then  $[L, L] \neq 0$ . However,  $L^{(2)} = [[L, L], L] = 0$ . Hence  $\mathfrak{n}(3, \mathbb{F})$  is nilpotent of index 1.<sup>1</sup> Hence  $L/[L, L]$  is abelian of dimension 2, and  $[L, L]$  abelian of dimension 1, and it is central (contained in the center). Being of dimension 1, we conclude  $[L, L] = Z(L)$ .

$\mathfrak{n}(3, \mathbb{F})$  is isomorphic to the **first Heisenberg algebra** denoted  $\mathfrak{H}_1$ .

**Proposition 2.1.9.** For every  $n \geq 2$  and field  $\mathbb{F}$ ,  $\mathfrak{n}(n, \mathbb{F})$  is a nilpotent Lie algebra of nilpotency index  $n - 2$ .

**Definition 2.1.10.** An element  $x \in L$  is called **ad-nilpotent** if  $\text{ad } x$  is a nilpotent linear transformation on  $L$ . Namely,  $\exists k \in \mathbb{N}$ :  $(\text{ad } x)^k = 0$ .

**Remark 2.1.11.** In general, in a Lie algebra  $L$  which is nilpotent of index at most  $n - 1$ ,  $L^{(n)} = 0$ , or equivalently

$$[[\dots [[x_1, x_2], x_3], x_4] \dots, x_n], x_{n+1}] = 0$$

for all  $x_1, \dots, x_{n+1}$ . Equivalently the product (in any order) of the linear transformations  $\text{ad } x_2, \dots, \text{ad } x_{n+1}$  is zero.

**Theorem 2.1.12 (Engel).** Let  $L$  be a Lie algebra such that every element of  $L$  is ad-nilpotent. Then  $L$  is a nilpotent Lie algebra.

For the proof we shall develop some properties of nilpotent linear Lie algebras, namely Lie sub-algebras of  $\text{End}(V)$ .

**Proposition 2.1.13.** Let  $X \in \text{End}(V)$  be a nilpotent linear transformation on  $V$ . Then  $\text{ad}(X)$  is a nilpotent linear transformation on  $\text{End}(V)$ , in particular  $\text{ad}(X) \in \text{End}(\text{End}(V))$ .

---

<sup>1</sup>Similarly once can show that  $\mathfrak{n}(n, \mathbb{F})$  is nilpotent of index  $n - 2$ .

*Proof.* Define for each  $X \in \text{End}(V)$  two linear maps on  $\text{End}(V)$ :

$$\begin{aligned}\lambda_X(Y) &= XY \\ \rho_X(Y) &= YX\end{aligned}$$

Clearly if  $X^k = 0$  then  $\rho_X^k = \lambda_X^k = 0$ . Furthermore,  $\rho_X$  commutes with  $\lambda_X$  (as linear maps on  $\text{End}(V)$ ). I.e.  $[\lambda_X, \rho_X] = 0$ . This is obvious because  $(XY)X = X(YX)$ . In general, in any associative algebra, (or any ring) the sum or the difference of two commuting nilpotent elements is also a nilpotent element.

We have

$$(\lambda_X - \rho_X)(Y) = XY - YX = [X, Y] = \text{ad}(X)(Y)$$

so it suffices to prove the last claim, since this implies  $\text{ad } X$  is nilpotent.

By the binomial formula,

$$(a - b)^N = \sum_{j=0}^N \binom{N}{j} a^j (-b)^{N-j}.$$

If  $a^k = b^k = 0$ , then for large  $N$  s.t.  $\min\{j, N - j\} \geq k$ , the sum vanishes. ■

**Remark 2.1.14.**  $\text{ad } X: L \rightarrow L$  is a nilpotent linear transformation with index of nilpotency being  $n - 1$ .

**Remark 2.1.15.** We saw that  $X \in \text{End}(V)$  is nilpotent,  $\text{ad } X \in \text{End}(\text{End}(V))$  is nilpotent. The converse is *not* true. For example take  $X = I + \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$  which is *not* nilpotent, but is  $\text{ad}$ -nilpotent.

**Theorem 2.1.16.** Let  $L \subseteq \mathfrak{gl}_n(V)$  all of whose elements are nilpotent linear transformations. Then there exists  $v \neq 0$  such that  $\forall X \in L: Xv = 0$ . Namely, a sub-algebra of  $\mathfrak{gl}_n(V)$  consisting of nilpotent elements has a non-trivial joint kernel.

*Proof.* Let us prove the theorem by induction on  $\dim L$ .

**Induction Basis:** The theorem is clearly true if  $\dim L = 1$ . Then  $L = \mathbb{F}x$  and  $x$  is nilpotent, so there's  $v \neq 0$  such that  $xv = 0$ .

**Induction Step:** (I) Assume the statement of the theorem for all linear Lie algebra of dimension less than  $n \geq 2$ . Let  $L$  have dimension  $n \geq n$  and let  $L_0 \subseteq L$  be a sub-algebra of strictly smaller dimension. (e.g. the span of a single matrix) Consider the linear maps  $\text{ad } x$  where  $x \in L_0$ . We have  $L_0 \subseteq L \subseteq \text{End}(V)$ . Now  $\text{ad } x$  leaves both the linear sub-spaces  $L_0$  and  $L$  invariant. In fact  $L$  is  $\text{ad } y$  invariant for any  $y \in L$ . (since  $L$  is closed under brackets) So  $\text{ad } x(L) \subseteq L$  and  $\text{ad } x(L_0) \subseteq L_0$ . Therefore  $\text{ad } X$  also acts on  $L/L_0$ <sup>2</sup> via

$$\overline{\text{ad } x}(y + L_0) = \text{ad } x(y) + L_0.$$

Now,

$$\dim \{\text{ad } x \mid x \in L_0\} \stackrel{\star}{\leq} \dim L_0 < L$$

where  $\star$  is true because  $\text{ad}$  is linear on  $L_0$ , and cannot expand the dimension. But,  $\overline{\text{ad}}(L_0)$  is in fact a linear Lie algebra consisting of linear transformations of  $\mathcal{U} := L/L_0$ , because we saw that  $\text{ad}$  is in fact a Lie-algebra homomorphism. Now, each  $\overline{\text{ad } x}$  with  $x \in L_0$ , is a nilpotent linear transformation on the factor  $L/L_0$ , since  $x$  and hence  $\text{ad } x$  are nilpotent linear maps. Furthermore,  $\dim \overline{\text{ad}}(L_0) < \dim L$ , so by the induction hypothesis,  $\overline{\text{ad}}(L_0)$  has a non-trivial vector in the joint kernel. I.e.  $\exists y + L_0 \neq L_0$  such that  $\overline{\text{ad}}(x)(y + L_0) = 0 + L_0$  for all  $x \in L_0$ . Namely,  $[x, y] + L_0 = 0 + L_0$  for all  $x \in L_0$ , or equivalently  $[x, y] \in L_0$  for all  $x \in L_0$ , so  $y$  normalises the sub-algebra  $L_0$ . So,  $\text{span}(L_0, y)$  is a Lie sub-algebra of  $L$ <sup>3</sup>, containing  $L_0$  strictly.<sup>4</sup>

**Remark 2.1.17.**  
 $N_1 := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$   
and  
 $N_2 := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$   
are both nilpotent matrices, but  
 $A := [N_1, N_2] = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$  is *not*.  
These matrices are a linear basis of  $\mathfrak{sl}_2 := \{z \in \mathfrak{gl}_2 \mid \text{tr } z = 0\}$ .

(II) Let now  $L_0$  be a sub-algebra of  $L$  such that  $L_0 \subsetneq L$  and it's maximal with this property. Applying the previous argument to  $L_0$ , we have  $N_L(L_0) \supsetneq L_0$  and therefore  $N_L(L_0) = L$ . So such an  $L_0$  is an ideal of  $L$ .

(III) Consider  $L/L_0$ , which is a Lie algebra. If  $x_0 + L_0 \neq L_0$  then  $\mathbb{F}(x_0 + L_0)$  is a Lie sub-algebra of  $L/L_0$ . Its inverse image in  $L$  (under the canonical Lie-algebra homomorphism  $L \rightarrow L/L_0$ ) is a Lie sub-algebra of  $L$ , containing  $L_0$ . But, having chosen  $L_0$  to be maximal, and because  $x_0 \notin L_0$ , we have  $\mathbb{F}x_0 + L_0 \supsetneq L_0$ . So  $L = \mathbb{F}x_0 + L_0$ , namely  $L_0$  is an ideal of co-dimension 1. So our sub-algebra  $L_0$  which has  $L_0 \subsetneq L$ , and maximal with this property, turns out to be an ideal of co-dimension 1.

<sup>2</sup>This isn't necessarily a Lie algebra.

<sup>3</sup>We know that the normaliser is a sub-algebra.

<sup>4</sup>Since  $y \notin L_0$ .

- (IV) Now consider the action of  $\text{ad}(L_0)$  on  $V$ . Now  $\dim L_0 < \dim L$  and so by the induction hypothesis there's  $v_0 \in V$  such that  $v_0 \neq 0$  and  $xv_0 = 0$  for all  $x \in L_0$ . We must find  $v \neq 0$  such that  $xv = 0$  for all  $x \in L$ . Let now

$$W = \{w \in V \mid \forall x \in L_0: xw = 0\}$$

be the common kernel of non-zero elements in  $L_0$ . We claim that  $W \subseteq V$  is invariant under the transformations in  $L$ . This finishes the proof, because it follows that  $x_0$  leaves  $W$  invariant, and since  $x_0$  is nilpotent, it must have a non-zero vector  $v \in W$  such that  $x_0v = 0$ . This  $v$  satisfies that  $x_0v = 0$  and  $xv = 0$  for all  $x \in L_0$ , and therefore  $xv = 0$  for all  $x \in L$ .

- (V) We have to show that indeed  $W$  is invariant under  $L$ . Let  $y \in L$  and let  $w \in W$ . We should show that  $yw \in W$ . So, we must show that for all  $x \in L_0$ , we have that  $x(yw) = 0$ . We shall prove this. We have

$$x(yw) = y(xw) + [x, y](w).$$

Now  $xw = 0$  since  $x \in L_0$  and  $w \in W$ , and  $[x, y]w = 0$  since  $[x, y] \in L_0$  and  $w \in W$  (for  $L_0$  is an ideal). Therefore  $x(yw) = 0$  as required. ■

We remind Engel's theorem, for which we proved the above.

**Theorem 2.1.18 (Engel).** *Let  $L$  be a Lie algebra such that every element of  $L$  is ad-nilpotent. Then  $L$  is a nilpotent Lie algebra.*

*Proof.* Consider  $\text{ad}: L \rightarrow \text{End}(L)$ .  $\text{ad}(L)$  is a linear Lie algebra consisting of linear maps on  $L$ . By assumption,  $\text{ad}(L)$  consists of nilpotent linear maps and by the previous theorem, there is  $z \in L \setminus \{0\}$  which is the common kernel of all  $\text{ad } x$  with  $x \in L$ . Namely,

$$\exists z \neq 0 \forall x \in L: [x, z] = \text{ad } x(z) = 0.$$

So,  $z \in Z(L)$  and consider  $L/Z(L) = \bar{L}$  which is a Lie algebra of dimension strictly less than  $\dim L$ . But,  $\bar{L}$  is also ad-nilpotent since

$$\overline{\text{ad}}(x): \bar{L} \rightarrow \bar{L}$$

is a transformation obtained from  $\text{ad } x$  by passing to a factor space. So by induction on the dimension,  $L/Z$  is nilpotent, and so  $L$  is nilpotent since we saw that if  $L/Z$  is nilpotent ( $Z$  the center) then  $L$  is nilpotent. This proves that  $L$  is nilpotent.

### 2.1.1 Flags

Let  $V$  be a vector space over  $\mathbb{F}$ . A **full flag** in  $V$  is a sequence of linear subspaces

$$V_0 = 0 \subsetneq V_1 \subsetneq V_2 \subsetneq \dots \subsetneq V_n = V$$

such that  $\dim V_i = i$  for all  $0 \leq i \leq n$ .

A **partial flag** is any sequence

$$W_1 \subsetneq W_2 \subsetneq \dots \subsetneq W_k$$

of nested subspaces. ■

**Definition 2.1.19.** Given a full flag, a linear transformation  $T: V \rightarrow V$  is said to **stabilise the flag** if  $TV_i \subseteq V_i$  for  $0 \leq i \leq n$ .

Let us choose  $e_1 \in V_1$ ,  $e_1, e_2 \in V_2$  a basis, etc. such that  $e_1, \dots, e_i$  is a basis of  $V_i$ . The matrix representing  $T$  in this basis is upper-triangular (because  $TV_i \subseteq V_i$ ). Conversely, a linear transformation  $S$  represented in this basis by an upper triangular matrix, stabilises the flag.

Similarly, given for example a partial flag  $V_0 \subsetneq W_1 \subsetneq W_2 \subseteq V$ , with  $\text{rank } W_i = k_i$ , we can choose a basis of  $W_1$ , complement it to a basis of  $W_2$ , then to a basis of  $V$ . A transformation  $U$  stabilises the partial flag if and only if it's represented by a block upper-triangular matrix with blocks of sizes  $k_1, k_2 - k_1, n - k_2$ . In our linear theorem we that every linear Lie algebra  $L$  consisting of nilpotent linear maps, has a non-zero vector in the common kernel. It follows that  $L$  stabilises a full flag, and in a basis adapted to this flag (as we chose before) all linear transformations in our algebra have a common upper-triangularisation, with zeroes on the main diagonal. We want to show that indeed  $L$  stabilises a full flag.



**Claim 2.1.20.**  $L$  stabilises a full flag.

*Proof.* There's  $v \in V$  non-zero such that for all  $x \in L$ ,  $xv = 0$ . Let  $V_1 = \mathbb{F}v$  and consider  $V/V_1$ . Now  $V_1$  is invariant under all  $x \in L$ , so  $xV_1 \subseteq V_1$ , since  $xv = 0$ . So  $x$  defines a transformation  $\bar{x}: V/V_1 \rightarrow V/V_1$ . This collection  $\{\bar{x} \mid x \in L\}$  is a nilpotent linear Lie algebra. Therefore,  $\bar{L}$  has a vector  $v_2$  such that  $x(v_2 + V_1) = 0 + V_1$ , and where  $v_2 + v_1 \neq 0 + V_1$ . So, if  $V_2 = \text{span}\{v_1, v_2\}$  then  $xV_2 \subseteq V_2$ . Furthermore,  $xV_1 = 0$  and  $xv_2 \in V_1$ . More generally, by induction,  $\bar{L}$  stabilises a full flag in  $V/V_1$ , and its inverse image in  $V$ , together with  $V_1$  is a full flag in  $V$ , which is invariant under all  $x \in L$ . Also, in the basis associated to this flag, the representing matrix has 0 on the diagonal. So every linear nilpotent Lie algebra stabilises a flag, with representing matrices as described.

Now,  $\mathfrak{n}(n, \mathbb{F}) \subseteq \mathfrak{t}(n, \mathbb{F})$  is a nilpotent Lie algebra. ■

**Conclusion.** Every linear nilpotent Lie algebra has a basis in which it is represented by a sub-algebra of  $\mathfrak{n}(n, \mathbb{F})$ .

**Corollary 2.1.21.** Let  $L$  be a nilpotent algebra.  $L$  must have an invariant flag. This flag gives a sequence of ideals

$$0 = I_0 \subsetneq I_1 \subsetneq I_2 \subsetneq \dots \subsetneq I_n$$

where each  $I_j$  is an ideal of  $L$  and they have dimension  $\dim I_j = j$ .

## 2.2 Solvable Lie algebras

**Definition 2.2.1 (Derived sequence of ideals).** Let  $L$  be a Lie algebra. Denote  $D_1(L) = L^{(1)} = [L, L]$  and similarly  $D_k(L) = [D_{k-1}(L), D_{k-1}(L)]$  for all  $k$ .  $(D_k)_{k \in \mathbb{N}_+}$  is the **derived sequence of ideals** for  $L$ .

**Definition 2.2.2.**  $L$  is **solvable** if  $D_k(L) = 0$  for some  $k$ .

**Remark 2.2.3.** Every nilpotent Lie algebra is solvable.  $L^{(k)} = 0$  implies  $D_k(L) = 0$ .

**Definition 2.2.4.** If  $D_k(L) = 0$  and  $D_{k-1}(L) \neq 0$  where  $k \in \mathbb{N}_+$ , we say  $L$  is **solvable of index**  $k - 1$ .

**Example.** The simplest solvable non-nilpotent algebra is the 2-dimensional algebra of  $2 \times 2$  matrices generated by  $X = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  and  $Y = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ . Here  $[X, Y] = XY - YX = Y$ . We have  $D_1(L) = [L, L] = \text{Span}(Y)$ . We have  $D_2(L) = 0$  since  $[L, L]$  is abelian. But,  $[[L, L], L] = [L, L]$ , so  $L$  is not nilpotent.

**Example.** For every  $n$  and  $\mathbb{F}$ ,  $\mathfrak{t}(n, \mathbb{F})$  is a solvable algebra.  $D_1(\mathfrak{t}(n, \mathbb{F})) \subseteq \mathfrak{n}(n, \mathbb{F})$  and  $\mathfrak{n}(n, \mathbb{F})$  is nilpotent, so  $\mathfrak{t}(n, \mathbb{F})$  is solvable.

**Proposition 2.2.5 (properties of solvable algebras).** 1. Every sub-algebra and quotient algebra of  $L$  is also solvable.

2. If  $I$  is an ideal in  $L$  and both  $I$  and  $L/I$  are solvable, then  $L$  is solvable.

3. If  $I$  and  $J$  are solvable ideals, then  $I + J$  is also solvable.

*Proof.* First, if  $\varphi: L \rightarrow L'$  is an epimorphism of Lie algebras, then  $\varphi([L, L]) = [L', L']$ , and in general

$$\varphi(D_k(L)) = D_k(\varphi(L)) = D_k(L').$$

1. Clearly, if  $L_0 \subseteq L$  then  $D_k(L_0) \subseteq D_k(L)$ , so  $D_k(L)$  implies  $D_k(L_0)$ , and  $L_0$  is solvable of index at most that of  $L$ .

Similarly, if  $L' = L/I$  is a quotient algebra, and  $\varphi: L \rightarrow L/I$  is the canonical epimorphism, then  $D_k(L') = \varphi(D_k(L))$  and  $D_k(L) = 0$  implies  $L'$  is solvable of index at most that of  $L$ .

2. Suppose that  $\bar{L} = L/I$  is solvable. Then  $D_k(\bar{L}) = \bar{0}$  and equivalently  $D_k(L) \subseteq I$ . Now, if  $I$  is a solvable ideal, then  $D_l(I) = 0$  for some  $l$ . Then  $D_{k+l}(L) \subseteq D_l(I) = 0$ . So,  $L$  is solvable of index at most  $l + k$ .

3. If  $I, J$  are solvable ideals, consider  $I + J/J \cong I/(I \cap J)$ . Since  $I$  is solvable, so is  $I/(I \cap J)$ . So then  $I + J/J$  is solvable. Since  $J$  is solvable, we get by (2). ■

**Proposition 2.2.6.** Every Lie algebra  $L$  has a unique maximal solvable ideal, containing all other solvable ideals.

*Proof.* Let  $R$  be a solvable ideal, maximal with this property. If  $I$  is any solvable ideal,  $R + I \supseteq R$  is solvable. Hence from maximality  $R + I = R$ , and hence  $I \subseteq R$ . ■

**Remark 2.2.7.** We obtained that  $R$  is the sum of all solvable ideals.

**Definition 2.2.8.**  $R$  is called the **solvable radical** of  $L$ , denoted  $\mathfrak{R} = \text{Rad}(L)$ .

**Question 2.2.9.** We say that if  $L/I$  and  $I$  are solvable, then  $L$  is solvable. Is it true that if  $L/I$  and  $I$  are nilpotent then  $L$  is nilpotent?

The answer is no. Take  $L = \left\{ X = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, Y = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \right\}$ . Then  $I = [L, L] \leq L$  is  $\text{Span}(Y)$ . Both  $I$  and  $L/I$  are abelian, hence nilpotent. However,  $L$  is not!

**Definition 2.2.10.** A Lie algebra  $L$  is called **semi-simple** if its radical vanishes.

**Exercise.** Let  $L$  be a Lie algebra. Prove that  $L/\text{Rad}(L)$  is semi-simple. Namely,  $\text{Rad}\left(L/\text{Rad}(L)\right) = 0$ .

**Theorem 2.2.11 (Lie's theorem on solvable algebras).** Let  $\mathbb{F}$  be an algebraically-closed field such that  $\text{char}(\mathbb{F}) = 0$ . Let  $L \subseteq \text{End}(V)$  be a solvable Lie algebra. Let  $V$  be a vector space over  $\mathbb{F}$ .

1. There's  $v$  non-zero which is a joint eigenvector of all  $x \in L$ .
2.  $L$  stabilises a full flag
$$0 = V_0 \subseteq V_1 \subseteq V_2 \subseteq \dots \subseteq V_n = V.$$
3. In a basis adapted to the flag,  $e_1, \dots, e_n$  such that  $\text{Span}\{e_1, \dots, e_i\} = V_i$ , all linear transformations  $X \in L$  are represented by upper triangular matrices.

**Example.**  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \in M_2(\mathbb{R})$  has eigenvalues  $\pm i$ . Take  $L = \mathbb{R} \cdot \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ , which is abelian and hence solvable. This has no upper-triangularisation because the eigenvalues are in  $\mathbb{R}$ . The conclusion of Lie's theorem doesn't hold for Lie algebras over  $\mathbb{R}$ .

*Proof.* We prove the theorem by induction on  $\dim(L)$ . The theorem has a similar conclusion to Engel's theorem and the proof follows similar lines. It suffices to prove 1 as the rest follows by modding out eigenvectors.

**Basis:** If  $\dim L = 1$ , then  $L = \mathbb{F} \cdot X$  and the theorem holds since every linear transformation is conjugate to an upper-triangular one over an algebraically-closed field.

**Step:** Since  $L$  is solvable,  $[L, L] \subsetneq L$ , but  $L/[L, L]$  is abelian, and in an abelian algebra, every sub-space is an ideal. Choose any subspace of codimension 1 in  $L/[L, L]$  and take its pre-image. This is an ideal  $K$  of  $K \leq L$  of codimension 1. By induction, since  $K$  is solvable, we have a joint eigenvector  $v$  of  $K$ . For  $Y \in K$  denote  $\lambda(Y)$  the eigenvalue of  $v$  under  $Y$ . Clearly

$$(\alpha Y_1 + \beta Y_2)v = \alpha Y_1 v + \beta Y_2 v = \alpha \lambda(Y_1)v + \beta \lambda(Y_2)v = \lambda(\alpha Y_1 + \beta Y_2)v$$

so  $\lambda: K \rightarrow \mathbb{F}$  is a linear functional. Let us define the **lambda-characteristic sub-space**  $W_\lambda \subseteq V$  by

$$W_\lambda = \{v \in V \mid \forall Y \in K: Yv = \lambda(Y)v\}$$

This is the sub-space consisting of all joint eigenvectors of  $K$  with joint eigenvalue  $\lambda$ .

**Note 2.2.12.** If we show that  $W_\lambda$  is  $L$ -invariant, the proof is complete, since  $L = \mathbb{F}X + K$  for some  $X$ , and in particular,  $W_\lambda$  is  $X$ -invariant, and  $X$  has an eigenvector  $u \in W_\lambda$  (since  $\mathbb{F} = \overline{\mathbb{F}}$ ). So,  $u$  is a joint eigenvector of  $L$ .

**Lemma 2.2.13.**  $W_\lambda$  is  $L$ -invariant.

*Proof.* Write  $L = K + \mathbb{F}X_0$  with some  $X_0$ .

- I) We need to show that for  $w \in W_\lambda$  and  $X \in L$  we have  $Xw \in W_\lambda$ . So we need to show that  $Y(Xw) = \lambda(Y)Xw$  for all  $Y \in K$ , by definition of  $W_\lambda$ . Now,  $Y(Xw) = XYw - [X, Y]w$ . Recalling that  $K$  is an ideal in  $L$ , we have  $[X, Y] \in K$  for all  $X \in L, Y \in K$ . So

$$\begin{aligned} Y(Xw) &= XYw - [X, Y]w \\ &= \lambda(Y)Xw - \lambda([X, Y])w \end{aligned}$$

since  $w \in W_\lambda$  and  $[X, Y] \in K$ . So we have to prove that

$$\forall X \in L \forall Y \in K: \lambda([X, Y]) = 0. \quad (2.1)$$

for all  $X \in L$  and  $Y \in K$ .

- II) To show (2.1), fix  $X \in L$  and fix  $w \in W$ . Consider the sequence

$$w, Xw, X^2w, \dots, X^{n-1}w, X^n w$$

where  $n$  is the least positive integers such that the sequence is linearly dependant. So, if we define

$$U_i = \text{Span} \{w, Xw, \dots, X^{i-1}w\}$$

then  $\dim U_i = i$  for  $1 \leq i \leq n$ . Also  $U_n = U_{n+1} = U_{n+2} = \dots$

- III) **Claim 2.2.14.** Each  $U_i$  for  $1 \leq i \leq n$  is invariant under  $K$ . Namely  $YU_i \subseteq U_i$  for all  $Y \in K$ .

*Proof.* We prove this claim inductively. First Let's see that  $U_1$  is  $K$ -invariant.

- (i)  $U_1$  is  $K$ -invariant for  $Yw = \lambda(Y)w$  for all  $Y \in K$ .
- (ii)  $U_2$  is  $K$ -invariant. Write  $U_2 = \mathbb{F}w + \mathbb{F}Xw$ . We've seen

$$YXw = \lambda(Y)Xw - \lambda([X, Y])w \in \mathbb{F}Xw + \mathbb{F}w.$$

So,  $K$  leaves  $U_2$  invariant, but in fact we know more:

$$YXw \equiv \lambda(Y)Xw \pmod{U_1}$$

since  $YXw = \lambda(Y)Xw + cw$ . So

$$YXw - \lambda(Y)Xw \in U_1.$$

- (iii) We claim that in general,

$$\forall 1 \leq i \leq n-1 \forall Y \in K: YX^i w \equiv \lambda(Y)X^i w \pmod{U_i}. \quad (2.2)$$

To see that, compute again.

$$\begin{aligned} YX^i w &= YX(X^{i-1}w) \\ &= XYX^{i-1}w - [X, Y]X^{i-1}w \end{aligned}$$

- By the induction hypothesis,  $YX^{i-1}w = \lambda(Y)X^{i-1}w + w'$  where  $w' \in U_{i-1}$ . So

$$X(YX^{i-1}w) = \lambda(Y)X^i w + Xw'.$$

But by definition,  $XU_{i-1} \subseteq U_i$ . Hence

$$XYX^{i-1}w = \lambda(Y)X^i w + w''$$

where  $w'' \in U_i$ .

- The second summand

$$[X, Y]X^{i-1}w = \lambda([X, Y])X^{i-1}w + w'''$$

where  $w''' \in U_{i-1}$  by the induction hypothesis. This means

$$[X, Y]X^{i-1}w \in U_i + U_{i-1} \subseteq U_i.$$

The net conclusion is that

$$YX^i w = \lambda(Y)X^i w + w''''$$

with  $w'''' \in U_i$ . So  $YX^i w \equiv \lambda(Y)X^i w \pmod{U_i}$  for all  $1 \leq i \leq n$ . ■

IV) We have proved (2.2). Formulated otherwise is says that in the basis of  $U_n = \{w, Xw, \dots, X^{n-1}w\}$  given by the sequence, the representing matrix of *every*  $Y \in K$  is upper triangular (that statement follows immediately from the fact the we proved  $KU_i \subseteq U_i$ ) and in fact, the diagonal has only the entry  $\lambda(Y)$ . So,  $\text{tr } Y|_{U_n} = n\lambda(Y)$  for *every*  $Y \in K$ . In particular, this is true for elements  $Y \in K$  which are of the form  $[X, Y]$  with  $Y \in K$ . So  $\text{tr } [X, Y]|_{U_n} = n\lambda([X, Y])$ .

We expect the trace of  $[X, Y]$  to vanish, and that is true here since both  $X$  and  $Y$  preserve  $U_n$ . The fact that  $U_n$  is  $X$ -invariant is obvious, and we saw that  $U_n$  is also invariant under *every*  $Y \in K$ . So

$$[X, Y]|_{U_n} = [X|_{U_n}, Y|_{U_n}]$$

and it follows that

$$\text{tr } [X, Y]|_{U_n} = 0 = n\lambda([X, Y]).$$

Now<sup>5</sup>  $\lambda([X, Y]) = 0$  for all  $Y \in K$  and  $X \in L$ . So we are done. ■

**Remark 2.2.15.** For every vector space  $V$  over a field  $\mathbb{F}$ , we can consider the spaces of flags over  $V$ .

**Example.** Consider the space of all lines in  $V$ . Namely

$$\text{Gr}_1(V) := \{\ell \subseteq V \mid \dim \ell = 1\}$$

(Grassmann 1, also known as the projective space over  $V$ ). Similarly we can take

$$\text{Gr}_k = \{W \subseteq V \mid \dim W = k\}$$

*the Grassmann variety of  $k$ -vector-spaces in  $V$ .* We can look more generally at any configuration

$$\text{Gr}_{k_1, \dots, k_m} := \{\ell_1 \subsetneq \ell_2 \subsetneq \dots \subsetneq \ell_m \mid \dim \ell_i = k_i, \ell_i \subseteq V\}.$$

Preservence of this flag corresponds to an existence of a basis such that the matrices have a certain upper-block-triangular form.

**Corollary 2.2.16.** *Let  $L$  be a solvable algebra over  $\mathbb{F}$ , where  $\mathbb{F} = \bar{\mathbb{F}}$  and  $\text{char } \mathbb{F} = 0$ . There is a full flag of ideals in  $L$ , namely*

$$0 \subsetneq L_1 \subsetneq L_2 \subsetneq \dots \subsetneq L_{n-1} \subsetneq L$$

where  $n = \dim L$ .

*Proof.* Consider  $\text{ad}: L \rightarrow \text{End}(L)$ .  $\text{ad}(L)$  is a solvable Lie algebra, so it stabilises a full flag by Lie's theorem (2). The corresponding sub-spaces are ideals: they satisfy  $\text{ad}(L)(L_i) \subseteq L_i$  and  $[L, L_i] \subseteq L_i$ , so  $L_i \triangleleft L$ . ■

**Corollary 2.2.17.** *If  $L$  is a solvable Lie algebra (with  $\mathbb{F}$  as above) then the commutator ideal  $[L, L]$  is nilpotent.*

*Proof.* Consider again the adjoint representation. We show that every element  $X \in [L, L]$  is ad-nilpotent as a linear transformation on  $L$ .<sup>6</sup>

So,  $\text{ad}(L)$  is a linear Lie algebra, solvable, and has a basis in which all linear transformations in  $\text{ad}(L)$  are upper-triangular. But, the (usual Lie) commutator of two upper-triangular matrices is a nilpotent matrix (as a strictly upper-triangular matrix). Hence

$$[\text{ad}(L), \text{ad}(L)] \subseteq \{\text{upper triangular matrices with 0 on the diagonal}\}.$$

Since we have  $[\text{ad}(L), \text{ad}(L)] = \text{ad}[L, L]$ , (since  $\text{ad}$  is a Lie-algebra homomorphism) so every  $X \in [L, L]$  is ad-nilpotent. ■

<sup>5</sup>For that we use  $\text{char}(\mathbb{F}) = 0$  and the proof wouldn't work otherwise

<sup>6</sup>It suffices to show it is ad-nilpotent when acting on  $[L, L]$ .

# Chapter 3

## Jordan-Chevalley decomposition

### 3.1 The Chinese remainder theorem

**Theorem 3.1.1 (Chinese remainder theorem).** Let  $R$  be a commutative unital ring, and let  $I, J$  be two ideals in  $R$  such that  $I + J = R$ . Then, given any  $a, b \in R$  there exists  $X \in R$  such that  $X \equiv a \pmod{I}$  and  $X \equiv b \pmod{J}$ .

*Proof.* Consider  $\pi: R \twoheadrightarrow R/I$  the canonical homomorphism. Since  $R = I + J$  clearly  $\pi: I \rightarrow R/I$  is also surjective. So for all  $a \in R$ ,  $\pi: I + a \rightarrow R/I$  is also surjective. So there is  $x \in I + a$  such that  $\pi(x) = b + J$ . So for any chosen  $b \in R$  we have  $x$  such that  $x \equiv a \pmod{I}$  and  $x \equiv b \pmod{J}$ . ■

**Theorem 3.1.2 (Chinese remainder theorem (more general)).** More generally, let  $I_1, \dots, I_n$  be ideals in  $R$  such that  $I_i + \bigcap_{j \neq i} I_j = R$  for any  $i \in [n]$ . Then, given  $a_1, \dots, a_n$  arbitrary, there is  $x \in R$  such that  $x \equiv a_i \pmod{I_i}$  for all  $i \in [n]$ .

*Proof.* By the Chinese remainder theorem<sup>1</sup>, for each  $i$  we can choose  $x_i$  such that  $x_i \equiv 1 \pmod{I_i}$  and  $x_i \equiv 0 \pmod{I_j}$  for  $j \neq i$ . Finally  $x = \sum_{i=1}^n x_i a_i$  satisfies  $x \equiv a_i \pmod{I_i}$  for all  $i \in [n]$ . ■

**Example.** Look in particular at the polynomial ring  $\mathbb{F}[x]$ . That is a Euclidean ring, hence a PID. So, every ideal  $I \triangleleft \mathbb{F}[x]$  is of the form  $p\mathbb{F}[x]$ . What does it mean that  $I + J = \mathbb{F}[x]$ ? It means that if  $J = q\mathbb{F}[x]$ , that  $p\mathbb{F}[x] + q\mathbb{F}[x] = \mathbb{F}[x]$ , so  $p$  and  $q$  are coprime. I.e. for some  $u(x), v(x)$  we have  $p(x)u(x) + q(x)v(x) = 1$ . Conversely, if  $p, q$  are co-prime polynomials, then there are such  $u(x)$  and  $v(x)$  such that  $p(x)u(x) + q(x)v(x) = 1$ . So  $p\mathbb{F}[x] + q\mathbb{F}[x] = \mathbb{F}[x]$ .

**Remark 3.1.3.** If  $p_1, \dots, p_n$  are pairwise co-prime, then

$$\bigcap_{j \neq i} p_j \mathbb{F}[x] = \left( \prod_{j \neq i} p_j \right) \mathbb{F}[x].$$

**Conclusion.** The Chinese remainder theorem, applied to  $\mathbb{F}[x]$ , implies that given pairwise co-prime polynomials  $p_1, \dots, p_n$ , and arbitrary  $a_1, \dots, a_n$ , there is a polynomial  $p$  such that  $p \equiv a_i \pmod{p_i \mathbb{F}[x]}$  for all  $i \in [n]$ .

### 3.2 Decomposition of vector spaces

**Proposition 3.2.1.** Let  $T$  be a linear transformation on a vector space over  $\mathbb{F}$  (arbitrary). Let  $f_T$  be the characteristic polynomial, and write  $f_T = p_1 p_2$  where  $p_1, p_2$  are co-prime. Then  $V$  decomposes to the direct sum of two  $T$ -invariant subspaces  $V = V_1 \oplus V_2$ , and more precisely  $V_1 = \ker p_1(T)$  and  $V_2 = \ker p_2(T)$ .

*Proof.* Start by writing  $u_1 p_1 + u_2 p_2 = 1$  for some polynomials  $u_i$ . Consider the ring homomorphism  $\mathbb{F}[x] \rightarrow \mathbb{F}[T]$  given by  $x \mapsto T$  and deduce that

$$I = u_1(T) p_1(T) + u_2(T) p_2(T).$$

Writing that again for all  $v \in V$ , we get

$$v = u_1(T) p_1(T) v + u_2(T) p_2(T) v. \quad (3.1)$$

(i) First,  $\ker p_1(T) \cap \ker p_2(T) = 0$ , by (3.1).

---

<sup>1</sup>to which we shall henceforward sometimes refer to as CRT

- (ii)  $\ker p_1(T) + \ker p_2(T) = V$ , since  $u_1(T)p_1(T)v \in \ker p_2(T)$  and  $u_2(T)p_2(T)v \in \ker p_1(T)$  as follows from  $f_T = p_1p_2$  and  $f_T(T) = p_1(T)p_2(T) = 0$  by Cayley-Hamilton.

So every vector  $v \in V$  is a sum of a vector in  $\ker p_1(T)$  and a vector in  $\ker p_2(T)$ .  $v \in \ker p_1(T)$  implies  $p_1(T)(Tv) = 0$ , so  $Tv \in \ker p_1(T)$ , and the kernel is an invariant sub-space. Similarly for  $\ker p_2(T)$ . ■

**Proposition 3.2.2.** *Let  $T$  be a linear transformation and assume its different eigenvalues  $a_1, \dots, a_n$  are all in  $\mathbb{F}$ . Write  $f_T(x) = \prod_{i=1}^n (x - a_i)^{m_i}$ . Then  $V$  decomposes to a direct sum  $V = \bigoplus_{i=1}^n V_i$  of  $T$ -invariant sub-spaces where  $V_i = \ker p_i(T)$  and  $p_i(T) = (T - a_i)^{m_i}$ .*

*Proof.* This follows immediately from the previous proposition, applied to  $(x - a_i)^{m_i}$  and  $\prod_{j \neq i} (x - a_j)^{m_j}$ . ■

**Theorem 3.2.3 (Jordan-Chevalley).** *Let  $T$  be a linear transformation over  $\mathbb{F}$  and assume that all of its eigenvalues are in  $\mathbb{F}$ . There exist two linear transformations  $T_s, T_n$  such that the following hold.*

- (i)  $T = T_s + T_n$
- (ii)  $T_n$  is nilpotent, and  $T_s$  is diagonalisable.
- (iii)  $T_s$  and  $T_n$  commute.
- (iv)  $T_s$  and  $T_n$  commute with  $T$  and with any other transformation that commutes with  $T$ .
- (v)  $T_s$  and  $T_n$  are given as polynomials in  $T$  without constant terms.
- (vi) If  $A \subseteq B$  are two sub-spaces and  $TB \subseteq A$ , then  $T_s B$  and  $T_n B \subseteq A$ .
- (vii) The first three properties determine the decomposition uniquely.

*Proof.* (I) Write  $f(x) = \prod_{i=1}^n (x - a_i)^{m_i}$  with  $a_i \neq a_j$  for  $i \neq j$ . Then  $V = \bigoplus_{i=1}^n V_i$  with  $V_i = \ker p_i(T)$  where  $p_i(T) = (T - a_i)^{m_i}$  as we saw. There exists a polynomial  $p(x)$  such that  $p(x) \equiv a_i \pmod{p_i}$  for  $i \in [n]$  and  $p(x) \equiv 0 \pmod{x}$ . This follows from CRT as follows: If some  $a_i = 0$ , then the condition  $p(x) \equiv 0 \pmod{x}$  is satisfied, and otherwise  $x$  is co-prime to each  $p_i$ , so that we can solve and find  $p(x)$  as stated.

(II) Define  $q(x) = x - p(x)$ , so  $p(x) + q(x) = x$ , so  $p(T) + q(T) = T$ . Define  $T_s = p(T)$  and  $T_n = q(T)$ . Clearly  $T_s + T_n = T$ ,  $T_s + T_n$  commute with  $T$  and with any other transformation that commute with  $T$ . In addition,  $p, q$  have no constant terms, by construction.

(III) We now restrict  $T_s$  and  $T_n$  to  $V_i$ , which is invariant under  $T$ , hence invariant under  $p(T)$  and  $q(T)$ . Now  $p(x) \equiv a_i \pmod{p_i}$ . That is  $p(x) - a_i = u_i(x)(x - a_i)^{m_i}$  for some polynomial  $u_i(x)$ . Here  $p_i(x) = (x - a_i)^{m_i}$ . But,  $V_i = \ker p_i(T) = \ker (T - a_i)^{m_i}$ , by definition. So obviously, for  $v_i \in V_i$  we have

$$(p(T) - a_i)v_i = u_i(T)p_i(T)v_i = 0.$$

So  $p(T)$  acts as the scalar  $a_i$  on  $V_i$ ! So  $T_s$  is a diagonalisable transformation with the same eigenvalues as  $T$ , namely  $a_1, \dots, a_n$ , each obtained  $\dim V_i$  times.<sup>2</sup> We claim that the restriction of  $q(T) = T_n$  to each  $V_i$  is nilpotent! Indeed, if  $v_i \in V_i$ , then

$$q(T)v_i = T_nv_i = (T - p(T))v_i = Tv - a_iv_i = (T - a_i)v_i.$$

Since  $v_i \in \ker p_i(T)$ , it follows that

$$T_n^{m_i}v_i = (T - a_i)^{m_i}v_i = 0.$$

So,  $T_n$  is nilpotent in each  $V_i$ , hence nilpotent.

So,  $T = T_s + T_n$  where  $T_s$  is diagonalisable,  $T_n$  is nilpotent, and they commute with each other and with every transformation commuting with  $T$ , and are given by polynomials in  $T$  without constant terms.

- (IV) *Action on sub-spaces:* If  $A \subseteq B$  and  $TB \subseteq A$ , then  $T^2B \subseteq TA \subseteq TB \subseteq A$ , so it follows that any polynomial in  $T$  without constant terms satisfies  $f(T)B \subseteq A$ .

---

<sup>2</sup>The characteristic polynomial of  $p(T)$  restricted to  $V_i$  is  $p_i(x)$ . This characteristic polynomial is co-prime to the characteristic polynomial of  $p(T)$  on  $V_j$ . So,  $f_{p_i(T)} \mid (x - a_i)^{m_i}$ , but the product of all these (partial) characteristic polynomials (on the invariant subspaces  $V_i$ ) is  $f_p(T)$ . This implies that  $\dim V_i = m_i$ .

(V) *Uniqueness*: Suppose that  $T = Q_s + Q_n$  so that  $Q_s$  is diagonalisable,  $Q_n$  is nilpotent, and  $Q_s = Q_n = Q_s = Q_n$ . We show  $Q_s = T_s$  and  $Q_n = T_n$ . But  $T = T_s + T_n = Q_s + Q_n$ . The fact that  $Q_s, Q_n$  commute implies that they commute with  $T = Q_s + Q_n$ . Hence,  $T_s$  and  $T_n$  commute with  $Q_s$  and  $Q_n$ , since they commute with every transformations commuting with  $T$ .<sup>3</sup> Consider  $T_s - Q_s = Q_n - T_n$ .  $T_n$  commutes with  $Q_n$  and the sum of two commuting nilpotent transformations is nilpotent by the binomial theorem. We now claim that  $T_s - Q_s$  is diagonalisable, and then, since all of its eigenvalues are zero (as a nilpotent transformation) it must be the zero transformation, so  $T_s = Q_s$  and  $T_n = Q_n$ . Indeed,  $T_s$  and  $Q_s$  are commuting diagonalisable transformations, so they have a common diagonalisation (when all the eigenvalues are in the field, because of diagonalisability). So, their sum or difference is also diagonalisable. ■

**Exercise.** Let  $\mathcal{F} \subset \mathfrak{gl}(V)$  be any set of commuting diagonalisable matrices. Then there is a basis of common eigenvectors to all transformations in  $\mathcal{F}$ . One can use induction on the dimension.

**Remark 3.2.4.** Write  $f_T(x) = \prod_{i=1}^n (x - a_i)^{m_i} = \prod_{i=1}^n p_i(x)$ .  $T|_{V_i}$  has a characteristic polynomial  $(x - a_i)^{m_i} = p_i(x)$ . Indeed,  $V_i = \ker p_i(T)$  by definition. So,  $(T - a_i)^{m_i} = p_i(T)$  acts as 0 on  $V_i$ . So the characteristic polynomial of  $T|_{V_i}$  has only  $a_i$  as a root. So it is  $(x - a_i)^{k_i}$ . But, this characteristic polynomial is co-prime to the characteristic polynomial of  $T$  on  $V_j$  when  $j \neq i$ . The product of all these partial characteristic polynomials on  $V_i$  is simply  $f_T$ . So,  $m_i = k_i$  and the characteristic polynomial of  $T|_{V_i}$  is  $(x - a_i)^{m_i} = p_i(x)$ . Now,  $p(T)$  leaves  $V_i$  invariant and acts on this  $m_i$ -dimensional space as a scalar. So, it has characteristic polynomial  $(x - a_i)^{m_i}$ .

**Remark 3.2.5.**  $f_{T_s}(x) = f_T(x)$ , but  $m_{T_s}(x) = \prod_{i=1}^n (x - \lambda_i)$  where  $\lambda_1, \dots, \lambda_n$  are the distinct eigenvalues of  $T$ .

**Definition 3.2.6.** A linear transformation is called **semi-simple** if all the roots of its minimal polynomial have multiplicity 1.

**Fact 3.2.7.** If the roots of  $f_T(x)$  are in  $\mathbb{F}$ , then  $T$  is semi-simple if and only if it's diagonalisable.

**Proposition 3.2.8.** Let  $T: V \rightarrow V$  be linear.

1. If  $S$  is diagonalisable, then so is  $\text{ad } S: \text{End } V \rightarrow \text{End } V$ .
2. If  $S$  is nilpotent, then  $\text{ad } S$  is nilpotent on  $\text{End } V$ .
3. If  $T = T_s + T_n$  is a Jordan-Chevalley decomposition for  $T$ , then  $\text{ad } T = \text{ad } T_s + \text{ad } T_n$  is the Jordan-Chevalley decomposition of  $\text{ad } T$ .

*Proof.* 1. Let  $T$  be diagonalisable over  $\mathbb{F}$  with eigenvectors  $v_1, \dots, v_n$  and eigenvalues  $\lambda_1, \dots, \lambda_n$ .

We first show that if  $D = \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}$  is diagonal, then  $\text{ad } D$  is diagonalisable on  $M_n(\mathbb{F})$ . In fact,

let  $\mathfrak{a}$  be the algebra of diagonal matrices, and let  $E_{i,j}$  be the matrices where  $(E_{i,j})_{k,\ell} = \delta_{i,k}\delta_{j,\ell}$ . Then  $\mathfrak{a} \subseteq \ker \text{ad } D$  since  $\mathfrak{a}$  is a commutative Lie algebra, and  $D \in \mathfrak{a}$ . Each  $E_{i,j}$  is an eigenvector of  $\text{ad } D$  and  $\text{ad } D(E_{i,j}) = DE_{i,j} - E_{i,j}D = (\lambda_i - \lambda_j)E_{i,j}$ . So,  $E_{i,i}$  and  $E_{i,j}$  form a basis of eigenvectors of  $\text{ad } D$ .<sup>4</sup> Let  $T$  be now a general linear map. Write  $D = PTP^{-1}$  and then  $\text{ad}(PTP^{-1})E_{i,j} = (\lambda_i - \lambda_j)E_{i,j}$ . We write  $T = P^{-1}DP$  and so

$$\begin{aligned} \text{ad } T(P^{-1}E_{i,j}P) &= T P^{-1}E_{i,j}P - P^{-1}E_{i,j}PT \\ &= P^{-1}DE_{i,j}P - P^{-1}E_{i,j}DP \\ &= P^{-1}(DE_{i,j} - E_{i,j}D)P \\ &= (\lambda_i - \lambda_j)P^{-1}E_{i,j}P \end{aligned}$$

therefore  $P^{-1}E_{i,j}P$  is an eigenvector of  $\text{ad } T$  with eigenvalue  $\lambda_i - \lambda_j$ .

<sup>3</sup>We constructed  $T_n$  and  $T_s$ , and they satisfy all the properties,  $Q_n$  and  $Q_s$  are currently more general.

<sup>4</sup>If the eigenvalues are distinct, we obtain that the kernel is generated by the  $E_{i,i}$ .

2. Consider  $\text{ad } S(X) = SX - XS = \lambda_S(X) - \rho_S(X)$  where  $\lambda_S(X) = SX$  and  $\rho_S(X) = XS$  and  $\lambda_S, \rho_S$  are two *commuting* nilpotent transformations on  $\text{End } V$  and so  $\lambda_S - \rho_S$  is also nilpotent by the binomial theorem.
3. By our characterisation,  $\text{ad } T_s$  is diagonalisable, and  $T_n$  is nilpotent. So,  $\text{ad } [T_s, T_n] = 0 = [\text{ad } T_s, \text{ad } T_n]$ . So  $\text{ad } T_s, \text{ad } T_n$  commute, so they are the Jordan-Chevalley decomposition. ■



# Chapter 4

## Cartan's criterion for semi-simplicity

### 4.1 Preliminary results

**Proposition 4.1.1.** *Let  $U \subseteq W \subseteq \text{End } V$  be linear subspaces. Define*

$$M = \{X \in \text{End } V \mid [X, W] \subseteq U\} = \{X \in \text{End } V \mid \text{ad } X(W) \subseteq U\}.$$

*Assume that  $\text{char } \mathbb{F} = 0$  and  $\mathbb{F} = \bar{\mathbb{F}}$ .*

*Let  $X \in M$ , if  $\text{tr } XY = 0$  for all  $Y \in M$  then  $X$  is nilpotent.*

*Proof.*  $X = X_s + X_n$ , so we show that  $X_s = 0$ . Let  $\lambda_1, \dots, \lambda_n$  be the eigenvalues of  $X_s$ . Consider the  $\mathbb{Q}$ -vector space defined by the linear span of the eigenvalues.

$$E := \text{span}_{\mathbb{Q}} \{\lambda_1, \dots, \lambda_n\}$$

Then  $\dim_{\mathbb{Q}} E < \infty$ . We want to show that  $\dim_{\mathbb{Q}} E = 0$  and then  $\lambda_i = 0$  for all  $i$ , and  $T_s = 0$ .

To show that  $E = 0$ , it is enough to show that  $E^* = \text{Hom}_{\mathbb{Q}}(E, \mathbb{Q}) = 0$ . Let  $f: E \rightarrow \mathbb{Q}$  be a  $\mathbb{Q}$ -linear functional, we want to show that  $f(\lambda_i) = 0$  for all  $i \in [n]$ .

To do that, let  $Y$  be the linear transformation such that in the basis  $B$  of eigenvectors of  $X_s$ , it (Y) is with values  $f(\lambda_i)$  on the diagonal. I.e.  $Yv_i = f(\lambda_i)v_i$  for all  $i \in [n]$ . So, the eigenvalues of  $\text{ad } Y$  are  $f(\lambda_i) - f(\lambda_j)$  for  $i, j \in [n]$  as we saw. The eigenvalues of  $\text{ad } X_s$  are  $\lambda_i - \lambda_j$  and there exists a polynomial  $p$  with no constant term such that  $p(\lambda_i - \lambda_j) = f(\lambda_i) - f(\lambda_j)$ .<sup>1</sup> Note that because  $f$  is linear, where  $\lambda_i - \lambda_j = \lambda_k - \lambda_\ell$  then  $f(\lambda_i) - f(\lambda_j) = f(\lambda_k) - f(\lambda_\ell)$ . We can arrange  $p$  to have no constant term. If  $\lambda_i - \lambda_j = 0$  then  $f(\lambda_i) - f(\lambda_j) = 0$  so  $p(0) = 0$ . Otherwise, 0 is distinct from  $\lambda_i - \lambda_j$  for  $i \neq j$ . So we can add it in. Given  $p$ , consider  $p(\text{ad } X_s)$ . It is diagonalisable since  $\text{ad } X_s$  is, and its eigenvalues coincide with those of  $\text{ad } Y$  with the same multiplicities. So,  $p(\text{ad } X_s) = \text{ad } Y$ . Now,  $\text{ad } X_s$  is a polynomial *without* constant term in  $\text{ad } X$ .<sup>2</sup>

We conclude that  $\text{ad } Y$  is a polynomial without constant term in  $\text{ad } X$ . Recall, that our assumptions are that

$$M = \{X \in \text{End } V \mid [X, W] \subseteq U\} = \{X \in \text{End } V \mid \text{ad } X(W) \subseteq U\}.$$

But, we chose  $X \in M$ , so we conclude that  $Y$  also satisfies that  $\text{ad } Y(W) \subseteq U$ .<sup>3</sup> So  $Y \in M$ . Our basic assumption was that  $\text{tr } XY = 0$  for all  $Y \in M$ .

We claim that  $\text{tr } XY = \sum_{i \in [n]} \lambda_i f(\lambda_i)$ . Let's compute in the basis  $B$  of eigenvectors of  $X_s$  (and also  $Y$ ).

$$XYv_i = f(\lambda_i)Xv_i = f(\lambda_i)(X_s + X_n)v_i = \lambda_i f(\lambda_i)v_i + f(\lambda_i)X_nv_i.$$

$X_n$  is nilpotent, and so the claim follows.

So we conclude that  $\sum_{i=1}^n \lambda_i f(\lambda_i) = \text{tr } XY = 0$ . Now,  $f(\lambda_i)$  are rational numbers because  $f: E \rightarrow \mathbb{Q}$  is a  $\mathbb{Q}$ -linear functional we obtain

$$\begin{aligned} f\left(\sum_{i \in [n]} \lambda_i f(\lambda_i)\right) &= \sum_{i \in [n]} f(\lambda_i f(\lambda_i)) \\ &= \sum_{i \in [n]} f(\lambda_i) f(\lambda_i) \\ &= \sum_{i \in [n]} f(\lambda_i)^2 \\ &= 0. \end{aligned}$$

<sup>1</sup> via Lagrange's polynomial of interpolation

<sup>2</sup> since  $\text{ad } X_s$  is the semi-simple part of  $\text{ad } X$ , and the semi-simple part is a polynomial in the transformation

<sup>3</sup> as a polynomial without constant term in  $\text{ad } X$ , which has this property

Finally  $f(\lambda_i) = 0$  for all  $i \in [n]$ . Then  $f = 0$  so  $E^* = \{0\}$  so  $E = \{0\}$  so  $X_s = 0$  so  $X = X_n$  is nilpotent. ■

**Theorem 4.1.2 (Lagrange's polynomial of interpolation).** Let  $a_1, \dots, a_m$  be distinct in any field  $\mathbb{F}$  and let  $b_1, \dots, b_m \in \mathbb{F}$ . There is a unique polynomial  $p(x) \in \mathbb{F}[x]$  such that  $p(a_i) = b_i$  for all  $i \in [m]$ .  $p$  is unique among polynomials of degree at most  $m - 1$ .

**Proposition 4.1.3.** Let  $L \subseteq \text{End } V$  be a Lie algebra. Assume  $\mathbb{F} = \bar{\mathbb{F}}$  and  $\text{char } \mathbb{F} = 0$ . If  $\text{tr } XY = 0$  for every  $X \in [L, L]$  and every  $Y \in L$  then  $L$  is a solvable Lie algebra.

*Proof.* We use the previous proposition. First, for all  $Y \in L$ ,  $\text{ad}(Y)L \subseteq [L, L]$  by definition. Consider  $[L, L] \subseteq \text{End } V$ ,  $U = [L, L]$ ,  $W = L$  and let  $M = \{Z \in \text{End } V \mid [Z, L] \subseteq [L, L]\}$ . Then  $L \subseteq M$ . We assume  $\text{tr } XY = 0$  for all  $Y \in L$ , but to use the previous proposition, we need to show that  $\text{tr } XZ = 0$  for all  $Z \in M$ . If we show that, then by the previous proposition,  $X$  is nilpotent, so  $[L, L]$  consists only of nilpotent linear transformations, and by the result preceding Engel's theorem<sup>4</sup>,  $[L, L]$  is nilpotent and so  $L$  is solvable.

To show that  $\text{tr } XZ = 0$  write  $X = [U, V]$  since  $X \in [L, L]$ . Now,

$$\begin{aligned} \text{tr}([U, V], Z) &= \text{tr}((UV - VU)Z) \\ &= \text{tr}(UVZ - VUZ) \\ &= \text{tr}(UVZ) - \text{tr}(V(UZ)) \\ &= \text{tr}(UVZ) - \text{tr}(UZV) \\ &= \text{tr}(U(VZ - ZV)) \\ &= \text{tr}(U[V, Z]) \end{aligned}$$

and  $Z \in M$  so  $[V, Z] = [Z, V] \in [L, L]$ . Also,  $U \in L$ , hence

$$\text{tr}(U[V, Z]) = \dots = \text{tr}(U[V, Z]) = \text{tr}([V, Z]U) = 0$$

by assumption. So, every  $X \in [L, L]$  is nilpotent. Then  $[L, L]$  is nilpotent, and then  $L$  is solvable. ■

**Remark 4.1.4.** We just saw that for a linear Lie algebra,  $L \subseteq \text{End } V$ ,  $\text{tr}([L, L]L) = \{0\}$  implies  $L$  is solvable.

## 4.2 Cartan's criterion

**Theorem 4.2.1.** Let  $L$  be any Lie algebra over a field  $\mathbb{F}$  as above. If  $\text{tr } \text{ad } x \text{ad } y = 0$  for all  $x \in [L, L]$  and  $y \in L$ , then  $L$  is solvable.

*Proof.* Consider  $\text{ad } L \subseteq \text{End } L$  and  $\text{ad } [L, L] = [\text{ad } L, \text{ad } L]$ , and so  $\text{ad } L$  is solvable by the proposition for the linear case. So  $\text{ad } (L) = L / \ker \text{ad} = L / Z(L)$  is solvable, and so  $L$  is solvable.<sup>5</sup> ■

<sup>4</sup>A Lie algebra of nilpotent transformations is nilpotent

<sup>5</sup>As an exercise, if  $L / Z(L)$  is solvable, so is  $L$ .

# Chapter 5

## Killing form

Let  $L$  be a Lie algebra and define a symmetric bilinear form

$$B_L(X, Y) := \text{tr}(\text{ad } X \text{ ad } Y).$$

This is called the ***Killing form***.

***Proposition 5.0.1 (invariance of the Killing form).***

$$B([x, y], z) = B(x, [y, z])$$

*Proof.* For any three linear transformations  $R, T, S$  ( $\text{ad } X, \text{ad } Y, \text{ad } Z$ , respectively) we shall compute  $\text{tr}([T, S]R)$  and  $\text{tr}(T[S, R])$  and show equality.

$$\begin{aligned} \text{tr}([T, S]R) &= \text{tr}((TS - ST)R) \\ &= \text{tr}(TSR) - \text{tr}(RST) \\ &= \text{tr}(TSR) - \text{tr}(TRS) \\ &= \text{tr}(T(SR - RS)) = \text{tr}(T[S, R]) \end{aligned}$$

hence there's equality. ■

***Proposition 5.0.2 (more invariance of the Killing form).***

$$B(\text{ad } Y(X), Z) + B(X \text{ ad } Y(Z)) = 0$$

*Proof.*  $B([X, Y], Z) = B(-\text{ad } Y(X), Z)$  and use the previous proposition. ■

More generally, let  $\pi: L \rightarrow \text{End } V$  be *any* linear representation of  $L$  (namely, a Lie-algebra homomorphism into  $\text{End } V$ ). Define

$$B_\pi(x, y) = \text{tr}(\pi(x)\pi(y)).$$

The proves of the above propositions stay the same, therefore

$$B_\pi([x, y], z) = B_\pi(x, [y, z]).$$

$B_\pi$  is a symmetric bilinear form satisfying the symmetry condition.

**Conclusion.** The Cartan criterion can be formulated to say that  $B_\pi(L, [L, L]) = 0$ , thus  $L$  is solvable.

### 5.1 Some properties of the Killing form

***Proposition 5.1.1.*** *Let  $I \triangleleft L$  be an ideal of  $L$ . Then the Killing form  $K_I$  of  $I$ , as a Lie algebra on its own is  $K_L|_{I \times I}$ , namely the restriction of the Killing form of  $L$  to  $I \times I$ .*

*Proof. Lemma 5.1.2.* *Let  $T: V \rightarrow V$  be a linear transformation. Assume that  $TV \subseteq W$ . We have a linear transformation  $T|_W: W \rightarrow W$ . Then  $\text{tr } T = \text{tr } T|_W$ .*

*Proof.* Indeed, if  $w_1, \dots, w_k$  is a basis of  $W$ , and  $w_{k+1}, \dots, w_n$  is a completion to a basis of  $V$ , then  $T$  is of the form  $\begin{pmatrix} T|_W & * \\ 0 & 0 \end{pmatrix}$ . ■

Now, let  $x \in I$  and  $y \in L$ , and consider the linear transformation  $\text{ad } x \text{ ad } y: L \rightarrow L \rightarrow I$ . So by the lemma,  $\text{tr ad } x \text{ ad } y$  on  $L$  is equal to  $\text{tr ad } x \text{ ad } y|_I$ . So,  $K_L(x, y) = \text{tr}(\text{ad ad } y|_I)$ . But now take  $x \in I$  and *also*  $y \in I$ . Then

$$K_I(x, y) = \text{tr}(\text{ad } x)|_I \cdot (\text{ad } y)|_I = \text{tr}((\text{ad } x \text{ ad } y)|_I).$$

$K_L(x, y) = K_I(x, y)$  if  $x, y \in I$ . ■

**Definition 5.1.3.** The *radical* of a symmetric bilinear form  $B$  is defined by the following.

$$\text{Rad}(B) = \{y \in V \mid B(x, y) = 0 \forall x \in V\}$$

**Remark 5.1.4.**  $\text{Rad}(B) \neq 0$  if and only if the form is degenerate.

**Proposition 5.1.5.** *The radical of the killing form is an ideal. In fact, this is true for any symmetric bilinear form which satisfies the symmetry condition, namely  $B([x, y], z) = B(x, [y, z])$ .*

*Proof.* Let  $x \in \text{Rad}(B)$  and  $y \in L$ . We should show that  $[x, y] \in \text{Rad}(L)$ . But, for all  $z \in L$ ,  $B([x, y], z) = B(x, [y, z]) = 0$  since  $x \in \text{Rad}(B)$ .

Hence  $[x, y]$  is  $B$ -orthogonal to all  $z \in L$ , hence  $[x, y] \in \text{Rad}(B)$ . ■

**Conclusion.** Consider the ideal  $I = \text{Rad}(K_L)$ . Then  $K_I = K_L|_{I \times I}$  as we proved. This restriction is obviously identically zero.

So,  $I = \text{Rad}(L)$  has  $K_I(I, [I, I]) = 0$  and hence by the Cartan criterion is solvable.

Namely, the radical of the Killing form is a solvable ideal.

**Conclusion.** We defined  $\text{Rad}(L)$  as the unique maximal solvable ideal<sup>1</sup> and so we conclude from the above that  $\text{Rad}(K_L) \subseteq \text{Rad}(L)$ .

**Conclusion.** If  $L$  is a semi-simple Lie algebra, then by definition  $\text{Rad}(L) = 0$  and so  $\text{Rad}(K_L) = 0$ , so the Killing form is non-degenerate.

**Theorem 5.1.6.** *A Lie algebra  $L$  is a semi-simple if and only if the Killing form  $K_L$  is non-degenerate. Namely,  $\text{Rad}(L) = 0$  if and only if  $\text{Rad}(K_L) = 0$ .*

*Proof.* We already saw that  $\text{Rad}(K_L) \subseteq \text{Rad}(L)$ , so we know that  $\text{Rad}(L) = 0$  implies  $\text{Rad}(K_L)$ .

Now assume that  $K_L$  is a non-degenerate form, and we have to show that  $L$  has no non-trivial solvable ideals. To do this, we start with a lemma.

**Lemma 5.1.7.** *If  $L$  has a non-trivial solvable ideal, then  $L$  has a non-trivial abelian ideal.*

*Proof.* First, if  $M$  is any solvable algebra, then for some  $k$ ,  $D_k(M) = [D_{k-1}(M), D_{k-1}(M)] = 0$  with  $D_{k-1}(M) \neq 0$ . So, by definition,  $D_{k-1}(M)$  is a non-trivial abelian ideal of  $M$ .

Now, let  $L$  be a general Lie algebra, and  $I$  a general ideal of  $L$ . Then  $D_k(I)$  are obviously ideals of  $I$ . In fact, they are also ideals of  $L$ !

**Claim 5.1.8.**  $D_k(I) \triangleleft L$ .

**Remark 5.1.9.** *Not every ideal of  $I$  is an ideal of  $L$ !*

*Proof.* To show that  $D_k(I)$  is an ideal of  $L$ , it is necessary and sufficient to show that it is an invariant sub-space under each of the linear transformations  $\text{ad } x: L \rightarrow L$  for all  $x \in L$ .

Now,  $\text{ad } x$  leaves  $I$  invariant, since  $I$  is an ideal, and  $\text{ad } x|_I$  is a derivation of  $I$ . It satisfies for all  $y, z \in L$  that

$$\text{ad } x[y, z] = [\text{ad } x(y), z] + [y, \text{ad } x(z)]$$

by the Jacobi identity. So,  $\text{ad } x: L \rightarrow L$  is a derivation, it is a map  $\delta: L \rightarrow L$  satisfying

$$\delta[y, z] = [\delta(y), z] + [y, \delta(z)].$$

Hence  $\text{ad } x|_I: I \rightarrow I$  is a derivation of  $I$ .

We conclude the proof<sup>2</sup> by saying that  $D_1(I) = [I, I]$  is invariant under *all* derivations of  $I$ :

$$\delta([I, I]) \subseteq [\delta(I), I] \subseteq [I, I]$$

Similarly,  $D_k(I)$  are invariant under *all* derivations (proof by induction). So, each  $D_k(I)$  is invariant under all  $\text{ad } x$  for  $x \in L$ . ■

<sup>1</sup>which contains every other solvable ideal

<sup>2</sup>of the claim

This claim proves the lemma, since if  $I$  is a solvable ideal, then  $D_{k-1}(I) \neq 0$  and  $D_k(I) = 0$  for some  $k$ , so  $D_{k-1}(I)$  is an abelian ideal of  $I$  which is an ideal of  $L$ . ■

Going back to the proof of the theorem, we have  $L$  which has a non-degenerate Killing form and we show it has no non-trivial abelian ideals. By the lemma, it has then a trivial solvable radical.

Let  $I \triangleleft L$  be abelian, namely  $[I, I] = 0$ . Let  $x \in I$  and  $y \in L$ . Then  $\text{ad } x \text{ ad } y: L \rightarrow I$ . We claim that  $(\text{ad } x \text{ ad } y)^2: L \rightarrow [I, I] = 0$ . If we prove this,  $\text{ad } x \text{ ad } y$  is a nilpotent linear transformation. So  $\text{tr ad } x \text{ ad } y = K_L(x, y) = 0$  for all  $y \in L$ . Because  $K_L$  is non-degenerate, this gives  $x = 0$ , hence  $I = 0$ .

To finish the proof, let  $z \in L$  and let  $w = \text{ad } x \text{ ad } y(z) \in I$ . Now,

$$\text{ad } x \text{ ad } y(w) = (\text{ad } x \text{ ad } y)^2(z) = [x, [y, w]] \in [I, I] = 0$$

since  $[y, w] \in I$  (because  $w \in I$ ). ■

**Theorem 5.1.10 (The decomposition of semi-simple algebras to simple ideals).** *Let  $L$  be a semi-simple Lie algebra. Then there are simple ideals  $I_i \triangleleft L$ , for  $i \in [k]$ , such that  $L = \bigoplus_{i \in [k]} I_i$ .  $I_i$  are uniquely determined up to order, and every simple ideal of  $L$  coincides with one of them.*

We remind that if  $A, B$  are Lie algebras, then

$$A \oplus B := \{(a, b) \mid a \in A, b \in B\}$$

is a Lie algebra under

$$[(a, b), (a', b')] = ([a, a'], [b, b'])$$

and then  $[(A, 0), (0, B)] = 0$  so the ideals  $(A, 0)$  and  $(0, B)$  commute.

**Remark 5.1.11.** For a semi-simple Lie algebra  $L$ , then  $K_{I_i} = (K_L)_{I_i \times I_i}$ .

*Proof.* If  $L$  is simple, the statement is obvious, so assume  $L$  is not simple.

Let  $J$  be an ideal of  $L$ , and assume that  $J$  is proper, and a minimal ideal. So  $\dim J < \dim L$  and  $J$  does not contain a non-trivial ideal of  $L$ . Consider now

$$J^\perp := \{y \in L \mid K(J, y) = 0\} = \{y \in L \mid K(x, y) = 0 \forall x \in J\}.$$

We claim that  $J^\perp$  is also an ideal. If  $x \in J$ ,  $y \in L$  and  $z \in J^\perp$  then  $[y, z] \in J^\perp$  since

$$K([z, y], x) = K(z, [y, x]) = 0$$

for  $z \in J^\perp$  and  $[y, x] \in J$  as  $x \in J$ . Therefore  $[z, y] \in J^\perp$ . So,  $I := J \cap J^\perp$  is also an ideal. So,  $J \cap J^\perp$  is also an ideal, but  $K_I = (K_L)_{I \times I}$  is zero identically by definition.

So,  $I$  is a solvable ideal by the Cartan criterion. Since we assumed that  $\text{Rad}(L) = 0$ , we get  $I = 0$  so  $J \cap J^\perp = 0$ . Therefore  $L = J \oplus J^\perp$ .

Now both  $J$  and  $J^\perp$  have trivial radical because  $L$  has trivial radical by the following sentence.  $[J, J^\perp] \subseteq J \cap J^\perp = 0$  and  $[J, J^\perp]$  commute which means an ideal of  $J$  or  $J^\perp$  is an ideal of  $L$ .

In fact, we have shown that when the Killing form is non-degenerate, every ideal  $J$  has a direct complement  $J^\perp$  which is also an ideal. Both of these also have non-degenerate Killing forms, since they have trivial radicals. By induction on the dimension, both  $J$  and  $J^\perp$  are a direct sum of simple ideals, which are also ideals of  $L$ . Therefore  $L$  is a direct sum of simple ideals

$$L = \bigoplus_{i \in [k]} I_i$$

and  $[I_i, I_j] = 0$  for  $i \neq j$ .

Now, let  $J$  be a simple ideal of  $L$ , we need to show  $J = I_{i_0}$  for some  $i_0$ . Clearly

$$\text{brs } J, L = \left[ J, \bigoplus_{i \in [k]} I_i \right] = \bigoplus_{i \in [k]} [J, I_i].$$

$[J, I_i]$  is an ideal of  $J^3$ , so  $[J, I_i] = 0$  or  $[J, I_i] = J$ .

$[J, L] \neq 0$  for otherwise  $Z(L) \supseteq J$ , but  $Z(L)$  is an abelian ideal. So,  $[J, I_{i_0}] = J$  for precisely one  $i_0$ .<sup>4</sup>

We want to show  $J = I_{i_0}$ . We already know  $J \subseteq I_{i_0}$  because  $J = [J, I_{i_0}] \subseteq I_{i_0}$ .  $I_{i_0}$  is simple, and  $J$  is an ideal, so  $J = I_{i_0}$ . ■

<sup>3</sup> $[J, [J, I_i]] \subseteq [J, I_i]$  since  $[J, I_i] \subseteq I_i$ .

<sup>4</sup>Otherwise,  $J$  has non-trivial ideals.

**Conclusion.** Let  $L$  be a semi-simple algebra. So  $L = \bigoplus_{i \in [k]} I_i$  for uniquely determined simple ideals. Then

1. Every ideal  $I \triangleleft L$  is a direct sum of some of the  $I_i$ . This is true since  $I$  itself is a semi-simple algebra, and every simple ideal of  $I$  is a simple ideal of  $L$ .

*Proof.*  $K_L$  is non-degenerate, so  $I \oplus I^\perp = L$ , again  $I \cap I^\perp = 0$  so  $[I, I^\perp] \subseteq I \cap I^\perp = 0$ ,  $I, I^\perp$  commute. So every ideal of  $I$  is an ideal of  $L$ . So, every simple ideal of  $I$  is a simple ideal of  $L$ . ■

2. Every factor  $L/I$ , where  $I \triangleleft L$  is also isomorphic to a direct sum of simple ideals isomorphic to some of the  $I_i$ .

This is an immediate consequence of 1.

## Chapter 6

# Derivations of simple algebras

Remind our assumptions  $\text{char}(\mathbb{F}) = 0$  and  $\mathbb{F} = \bar{\mathbb{F}}$ .

**Theorem 6.0.1.** *For every  $\delta \in \text{Der}(L)$ ,  $\text{Der}(L)$  contains also  $\delta_n, \delta_s$ . In fact, this is true for every  $\mathbb{F}$ -algebra  $L$ : The Lie algebra  $\text{Der}(L)$  contains the diagonalizable and nilpotent Jordan-Chevalley components of its elements.*

**Proposition 6.0.2.** *Define*

$$U_i = \left\{ v \in V \mid \exists k \in \mathbb{N}: (T - a_i)^k v = 0 \right\}.$$

*Then  $U_i = W_i$ .*

*Proof.*  $U_i$  is a  $T$ -invariant subspace. So,  $T|_{U_i}$  has a characteristic polynomial, and a minimal polynomial. Obviously, for some large  $N$ ,  $(T - a_i)^N \equiv 0$  on  $U_i$ . The minimal polynomial of  $T|_{U_i}$  divides  $(x - a_i)^N$ . On the other hand,  $f_T(x) := \prod_{i \in [r]} (x - a_i)^{m_i}$  acts as zero of  $V$  and on  $U_i$ . So, the minimal polynomial of  $T|_{U_i}$  divides  $f_T$ . So that minimal polynomial is a power of  $(x - a_i)$ , and the power is bounded by  $m_i$ . In particular,  $(x - a_i)^{m_i} v = 0$  for all  $v \in U_i$ . ■

**Definition 6.0.3.** Let  $\delta: L \rightarrow L$  be a derivation of an arbitrary  $\mathbb{F}$ -algebra, we define

$$L_a := \left\{ x \in L \mid \exists k: (\delta - a)^k(x) = 0 \right\}.$$

**Remark 6.0.4.**  $L_a$  is a sub-space of  $L$ , which is non-zero when  $a = a_i$  for the transformation  $T = \delta$ .

**Lemma 6.0.5.** *For every  $a$  and  $b$ , we have*

$$L_a \cdot L_b \subseteq L_{a+b}.$$

*Proof.* This follows from the generalised Leibniz formula.<sup>1</sup>

$$(\delta - (a + b)I)^n(x \circ y) = \sum_{j=0}^n \binom{n}{j} (\delta - aI)^{n-j}(x) (\delta - bI)^j(y)$$

For sufficiently large  $n$ , at least one of  $(\delta - aI)^{n-j}(x)$  or  $(\delta - bI)^j(y)$  vanishes for  $x \in L_a, y \in L_b$ , for every  $j$ . Then

$$(\delta - (a + b)I)^n(x \circ y) = 0$$

and so  $x \circ y \in L_{a+b}$ . ■

*Proof (theorem 6.0.1).* Let  $\delta \in \text{Der } L$  and write  $\delta = \delta_s + \delta_n$  in  $\text{End}(L)$ . To show  $\delta_n, \delta_s \in \text{Der } L$ , it is enough to show that  $\delta_s \in \text{Der } L$ .

To show that, we use that fact that it acts as a scalar in each of the subspaces  $L_a \neq 0$ . Namely, the scalar  $a$ . We use lemma 6.0.5 as follows. Fix  $x \in L_a$  and  $y \in L_b$  and compute

$$\delta_s(x \circ y) = (a + b)(x \circ y)$$

since  $x \circ y \in L_{a+b}$ . But,

$$\delta_s(x) \circ y + x \circ \delta_s(y) = ax \circ y + x \circ by = (a + b)(x \circ y).$$

So,

$$\delta_s(x \circ y) = \delta_s(x) \circ y + x \circ \delta_s(y),$$

and  $\delta_s$  is a derivation. ■

---

<sup>1</sup>The proof for the formula is done by induction, although it is surprisingly long.

**Theorem 6.0.6.** Let  $L$  be a semi-simple algebra. Then  $\text{ad}(L) = \text{Der}(L)$ .

Equivalently, the derivations of a semi-simple algebra are **inner derivations**, namely, given by  $y \mapsto [x, y] = \text{ad } x(y)$ .

**Lemma 6.0.7.** Let  $L$  be a Lie algebra and let  $\delta: L \rightarrow L$  be a derivation. Remind that each  $x \in L$  defines a linear map  $\text{ad } x$  which is also a derivation.

Then  $[\delta, \text{ad } x] = \text{ad } \delta(x)$ .

*Proof.* In the exercise sheets. ■

**Corollary 6.0.8.**  $\text{ad}(L) \subseteq \text{Der}(L)$  is an ideal of the algebra of derivations.

*Proof (theorem 6.0.6).* Let  $M := \text{ad } L$ , be an ideal of  $\text{Der } L$ . Furthermore,  $L \cong \text{ad } L = M$ , since  $\ker \text{ad}$  is the center  $Z(L)$  and a semi-simple algebra has a trivial center. So the Killing form of  $L$  is non-degenerate. So the Killing form of  $M$ , being its restriction, is non-degenerate.<sup>2</sup>

Now consider  $I = M^\perp$  the orthogonal of the Killing form of  $\text{Der } L$ . Now  $I \cap M = \{0\}$  since the Killing form of  $M$  (which is the restriction from  $\text{Der } L$ ) is non-degenerate. In addition, the orthogonal of an ideal under the Killing form is also an ideal, as we saw. So  $I$  is an ideal, and  $\text{Der } L = I \oplus M$ .

Finally,

$$[I, M] \subseteq I \cap M = 0$$

since they are ideals. Hence  $I$  and  $M$  commute, so  $[\delta, \text{ad } x] = 0$  if  $\delta \in I$ , and

$$[\delta, \text{ad } x] = 0 = \text{ad } \delta(x)$$

as we saw. Now,  $\text{ad}$  is injective on  $L$ , so  $\delta(x) = 0$  for all  $x \in L$ , and so  $\delta = 0$ . Hence  $I = 0$  and hence  $M = \text{Der } L$ . ■

**Theorem 6.0.9 (Abstract Jordan Chevalley decomposition of semi-simple Lie algebras).** For  $x \in L$  we can write  $x = x_s + x_n$ , with the properties

$$[x_s, x_n] = [x, x_s] = [x, x_n] = 0$$

and  $\text{ad } x_s$  is diagonalisable on  $L$ ,  $\text{ad } x_n$  is nilpotent on  $L$ , and the decomposition is unique.

*Proof.* Write  $\text{ad } x = T_n + T_s$  a Jordan-Chevalley decomposition in  $\text{End } L$ . From  $\text{Der } L = \text{ad } L \cong L$  so  $T_n, T_s$  come from elements  $x_n, x_s$  with  $x = x_s + x_n$ . The properties follow from those on  $\text{End } L$ . ■

**Exercise.** Assume  $L \subseteq \mathfrak{gl}(V)$  is semi-simple.

Show that the abstract Jordan-Chevalley decomposition of  $L$  agree with that of  $L$  viewed as linear transformations on  $V$ .

---

<sup>2</sup>We've seen that the restriction of the Killing form to an ideal, is the Killing form of the ideal.



# Chapter 7

## Toral subalgebras

Assume  $S$  is semi-simple,  $\text{char}(\mathbb{F}) = 0$  and  $\mathbb{F} = \bar{\mathbb{F}}$ . If every  $x \in L$  is ad-nilpotent, then  $L$  is a nilpotent Lie algebra. So for some  $x \in L$ , we have  $x_s \neq 0$ <sup>1</sup> where  $x = x_s + x_n$  and  $\text{ad } x_s$  is diagonalisable. So,  $x_n$  is ad-nilpotent and  $x_s$  is ad-diagonalisable.  $\mathbb{F} \cdot x_s$  is a Lie sub-algebra consisting of ad-diagonalisable elements, and any such a sub-algebra is called a **toral** sub-algebra.

**Theorem 7.0.1.** *A toral sub-algebra is abelian.*

*Proof.* Let  $T$  be toral, and  $x \in T$ . We want to show that  $\text{ad}_T(x) = 0$ , i.e.  $\text{ad } x: T \rightarrow T$  is the zero map. So we want to show  $[x, y] = 0$  for all  $y \in T$ .

Assume towards a contradiction that this isn't the case. Then since  $\text{ad } x$  is diagonalisable, there's  $y \in T$  non-zero which is an eigenvector of  $\text{ad}_T(x)$  with non-zero eigenvalue.

$\text{ad } x$  is diagonalisable on  $L$ , hence on the invariant subspace  $T$ , and if all its eigenvalues are 0 it is the zero map. So

$$[x, y] = ay \neq y$$

with  $a \neq 0$ . We should derive a contradiction.

Consider

$$\text{ad}_T(y, x) = [y, x] = -ay.$$

Then  $\text{ad}_T(y)(x)$  is a vector which is an eigenvector of  $\text{ad}_T(y)$  with eigenvalue 0.

But,  $\text{ad}_T(y)$  is diagonalisable in  $T$ , so *every* element of  $T$  is a linear combination of eigenvectors of  $\text{ad}_T(y)$ . So

$$x = \sum_i f_i x_i$$

with  $x_i$  being eigenvectors of  $\text{ad}_T(y)$  and  $f_i \neq 0$ .

Apply  $\text{ad}_T(y)$  to this equation, we get

$$-ay = \text{ad}_T(y)(x) = \sum_j f_j x_j$$

where the sum is ranging of  $j$  such that  $x_j$  is an eigenvector of  $\text{ad}_T(y)$  with non-zero eigenvalue and  $f'_j \neq 0$ .

But, we have this way a representation of the non-zero vector  $-ay$  as a linear combination of eigenvectors of  $\text{ad}_T(y)$  with non-zero eigenvalue. Because  $y$  itself is an eigenvector with eigenvalue 0 of  $\text{ad}_T(y)$ , this is a contradiction. ■

Let  $L$  be a semi-simple Lie algebra and let  $0 \neq H \leq L$  be a maximal toral sub-algebra. Then each  $h \in H$  has the property that  $\text{ad } h: L \rightarrow L$  is diagonalisable, and since  $H$  is abelian,  $L$  can be decomposed to joint eigen-spaces of all  $h \in H$ .

$$L_\alpha = \{x \in L \mid [h, x] = \alpha(h)x\}$$

Here,  $\alpha: H \rightarrow \mathbb{F}$  is a linear functional. If  $x$  is a joint eigenvector, we get the following where  $h_1, h_2 \in H$ .

$$[h_1, x] = \alpha(h_1)x, \quad [h_2, x] = \alpha(h_2)x, \quad [h_1 + h_2, x] = \alpha(h_1 + h_2)x$$

We can write

$$L = C_L(H) \oplus \bigoplus_{\alpha \in \Phi} L_\alpha$$

---

<sup>1</sup> $L$  is semi-simple, so it isn't nilpotent.

where

$$C_L(H) = L_0 = \{x \in L \mid \forall h \in H: [h, x] = 0\}$$

is the *centraliser of  $H$  in  $L$*  and where

$$\Phi = \{\alpha \in H^* = \text{Hom } H, \mathbb{F} \mid L_\alpha \neq 0\} \setminus \{0\}$$

is the set of roots of  $L$  under the toral sub-algebra  $H$ , *the root system of  $H$  with respect to  $L$* . We also write

$$\Phi = \Phi(L, H).$$

We list a few properties of the root system  $\Phi(L, H)$  in general.

Property 1.

$$[L_\alpha, L_\beta] \subseteq L_{\alpha+\beta}$$

*Proof.* Let  $x \in L_\alpha, y \in L_\beta$ . To see  $[x, y] \in L_{\alpha+\beta}$ , compute using the Jacobi identity:

$$\begin{aligned} \text{ad } h[x, y] &= [\text{ad } h(x), y] + [x, \text{ad } h(y)] \\ &= \alpha(h)[x, y] + \beta(h)[x, y] \\ &= (\alpha + \beta)(h)[x, y] \end{aligned}$$

Hence  $[x, y] \in L_{\alpha+\beta}$ . ■

Property 2. If  $\alpha + \beta \neq 0$ , then  $L_\alpha \perp L_\beta$  under the Killing form on  $L$ .

*Proof.* By the symmetry property of  $K$ ,

$$K([h, x], y) = \alpha(h)K(x, y) = -K(x, [h, y]) = -\beta(h)K(x, y).$$

Therefore

$$(\alpha + \beta)(h)K(x, y) = 0,$$

but if  $\alpha + \beta(h) \neq 0$  for some  $h \in H$ , then  $K(x, y) = 0$ , so  $L_\alpha \perp L_\beta$ . ■

**Proposition 7.0.2.** *If  $L$  is a semi-simple algebra, then the restriction of  $K$  to  $C_L(H)$  is non-degenerate.*

**Remark 7.0.3.**  $C_L(H)$  is *not* an ideal of  $L$ .

*Proof.* Let  $x \in C_L(H)$ . Then since for  $\alpha \neq 0$ ,  $\alpha + 0 \neq 0$ ,  $L_0 \perp L_\alpha$  for all  $\alpha \in \Phi$ .

So, if  $x \in L_0$  is orthogonal to  $L_0$  under  $K$ ,  $K(x, L_0) = 0$ , then  $K(x, L) = 0$  by linearity.

But in a semi-simple algebra,  $K$  is non-degenerate, so  $x = 0$ . Hence  $K|_{L_0}$  is non-degenerate. ■

**Theorem 7.0.4.** *If  $L$  is a semi-simple Lie algebra and  $H$  is a maximal toral sub-algebra, then  $C_L(H) = H$ .*

*Proof.* Let  $C = C_L(H)$ .

Step 1.  $C$  contains the Jordan-Chevalley components of its elements. This follows since the Jordan-Chevalley components are polynomials without constant term in the underlying element. Now, an element of  $C$  maps (under  $\text{ad}$ )  $H$  to 0, and hence the same is true for the Jordan-Chevalley components. So the Jordan Chevalley components are in  $C_L(H) = C$ .

Step 2. Every  $\text{ad}$ -diagonalisable element  $w \in C$  is in fact already in  $H$ . Indeed,  $H + F \cdot w$  is also toral, since  $w$  commutes with every  $h \in H$ , and so by maximality of  $H$ ,  $w \in H$ .

Step 3. The restriction of  $K$  to  $H$  is non-degenerate (not just that of  $K$  to  $C$ ). Suppose  $K(h, H) = 0$  for some  $h$ . Now let  $n \in C$  be  $\text{ad}$ -nilpotent. Then,  $\text{ad } h \text{ ad } n$  is nilpotent since

$$[\text{ad } h, \text{ad } n] = \text{ad } [h, n] = 0$$

and  $\text{ad } n$  is nilpotent. Therefore  $\text{tr}(\text{ad } h \text{ ad } n) = 0$ , so  $K(h, n) = 0$ . Therefore,  $h$  is  $K$ -orthogonal to every diagonalisable element in  $C^2$ , and to every  $\text{ad}$ -nilpotent element in  $C$ , and so  $K(h, C) = 0$ .

$K$  is non-degenerate on  $C$ , so  $h = 0$ . Hence  $K$  is non-degenerate on  $H$ .

---

<sup>2</sup>since such an element is already in  $H$  as we saw, and we assume  $K(h, H) = 0$

Step 4.  $C$  is a nilpotent algebra, since every element in  $C$  is ad-nilpotent and by Engel's theorem. Indeed if  $x \in C$  then  $x = x_n + x_s$ , and  $\text{ad } x = \text{ad } x_n + \text{ad } x_s$ , where  $\text{ad } x_s$  is diagonalisable, hence  $x_s \in H$ . Now

$$\text{ad } x = \text{ad } x_n + \text{ad } x_s|_C$$

and since  $x_s \in H$ ,  $\text{ad } x_s$  is the zero operator on  $C$ .

So, on  $C$   $\text{ad } x$  is equal to an ad-nilpotent operator, so  $x$  is ad-nilpotent on  $C$ .

Step 5.  $H \cap [C, C] = 0$  since by the symmetry property of the Killing form

$$K(H, [C, C]) = K([H, C], C) = K(0, C)$$

which is in turn true for  $C = C_L(H)$  means  $[C, H] = 0$ .

Since  $K$  is non-degenerate on  $H$ ,  $H \cap [C, C] = 0$ <sup>3</sup>.

Step 6. Assume towards a contradiction  $[C, C] \neq 0$ .

**Exercise.** In a nilpotent Lie algebra, every non-zero ideal intersects the center non-trivially.

Let  $z \in Z(C) \cap [C, C] \setminus \{0\}$ .  $z$  is *not* ad-diagonalisable, because in that case it would be in  $H$ , and as we just saw  $H \cap [C, C] = 0$ .

So,  $z$  has a non-trivial ad-nilpotent part, denoted  $z_n$ . Since  $z$  takes  $C$  to 0 under  $\text{ad}$ , so do  $z_s, z_n$ <sup>4</sup>. So,  $z_n$  is itself in  $Z(C)$ . But then  $[z_n, c] = 0$  for all  $c$ , and

$$\text{tr ad } z_n \text{ ad } c = 0$$

■

since  $\text{ad } z_n$  is nilpotent, and commutes with  $\text{ad } C$ . So,  $K(z_n, c) = 0$  for all  $c$ , and so  $K$  is degenerate on  $C$ , unless  $z_n$  is zero. This contradiction implies  $[C, C] = 0$  and  $C$  is abelian.

Step 7. We should show  $C_L(H) = C = H$ . Do this as an exercise.

**Corollary 7.0.5.** *The killing form of  $L$  restricted to  $H$  is a non-degenerate symmetric form.*

## 7.1 Orthogonality of root systems

Assume  $L$  is semi-simple over an algebraically-closed field  $\mathbb{F}$  of zero characteristic. As we saw,  $K$  is non-degenerate on  $H$ , and so we can transfer to  $H^* = \text{Hom}(H, \mathbb{F})$  by duality: For each  $\alpha \in H^*$  there is a unique  $h_\alpha$  such that  $K(h, h_\alpha) = \alpha(h)$ .<sup>5</sup> We can define

$$K(\alpha, \beta) := K(h_\alpha, h_\beta)$$

and obtain also  $K(\alpha, \beta) = \alpha(h_\beta) = \beta(h_\alpha)$ .

**Theorem 7.1.1 (orthogonality properties of root systems).** 1.  $\Phi$  spans  $H^*$ .

2.  $-\Phi = \Phi$

3. If  $\alpha \in \Phi$  and  $x \in L_\alpha, y \in L_{-\alpha}$  then

$$[x, y] = K(x, y) h_\alpha.$$

4.

$$\dim [L_\alpha, L_{-\alpha}] = 1$$

and this is spanned by  $h_\alpha$  for  $\alpha \in \Phi$ .

5.

$$\alpha(h_\alpha) = K(\alpha, \alpha)$$

and in particular this is non-zero.

<sup>3</sup>otherwise a non-zero vector in it is the radical of the form on  $H$ .  $h \in H \cap [C, C]$  implies  $K(h, H) = 0$  and this implies  $h = 0$ .

<sup>4</sup>by the Jordan-Chevalley properties

<sup>5</sup> $w \mapsto K(\cdot, w)$  for  $w \in H$  is a linear isomorphism  $H \rightarrow H^*$

6. If  $x \in L_\alpha \setminus \{0\}$ , there exists  $y \in L_{-\alpha} \setminus \{0\}$  such that  $x, y, \hat{h}_\alpha := [x, y]$  span a three-dimensional Lie sub-algebra of  $L$ , which is isomorphism to  $\mathfrak{sl}_2(\mathbb{F})$  under the following isomorphism.

$$\begin{aligned} x &\mapsto \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ y &\mapsto \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \\ \hat{h}_\alpha &\mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \end{aligned}$$

7.  $\hat{h}_\alpha$  is given by

$$\hat{h}_\alpha = \frac{2h_\alpha}{K(\alpha, \alpha)}$$

and

$$\hat{h}_{-\alpha} = -\hat{h}_\alpha.$$

*Proof.* 1. Let

$$H_0 := \text{span} \{h_\alpha \mid \alpha \in \Phi\}.$$

By duality,  $\Phi$  spans  $H^*$  iff  $H_0$  spans  $H$ .

If  $H_0$  does not span  $H$ , there is a functional

$$\gamma: H \rightarrow \mathbb{F}$$

non-zero and such that  $\gamma(H_0) = 0$ . So,  $\gamma(h_\alpha) = 0$  for all  $\alpha \in \Phi$ , so  $K(\gamma, \alpha) = 0$  for all  $\alpha \in \Phi$ , and equivalently,  $\alpha(h_\gamma)$  for all  $\alpha \in \Phi$ .

So,

$$[h_\gamma, x_\alpha] = \alpha(h_\gamma) x_\alpha = 0$$

if  $x_\alpha \in L_\alpha$ . Hence,  $h_\gamma$  commutes with every  $L_\alpha$  for  $\alpha \in \Phi$ , and with  $H$ . So  $h_\gamma \in Z(L) = \{0\}$ , in contradiction.

2. If  $\alpha \in \Phi$  and  $-\alpha \notin \Phi$ , then, as we saw,  $L_\alpha \perp L_\beta$  if  $\alpha + \beta \neq 0$ . But if  $-\alpha \notin \Phi$  then  $\alpha + \beta \neq 0$  for all  $\beta \in \Phi$ . So,  $L_\alpha \perp L_\beta$  for all  $\beta \in \Phi$ . Also,  $L_\alpha \perp H$  since  $\alpha + 0 \neq 0$ , and therefore  $L_\alpha \perp L^6$ . This is a contradiction, since by assumption on  $\alpha$ ,  $L_\alpha \neq \{0\}$ , and because  $K$  is non-degenerate.

3. Notice firstly that  $K(h, h_\alpha) = \alpha(h)$  implies

$$h_{-\alpha} = -h_\alpha. \tag{7.1}$$

Let  $\alpha \in \Phi$ ,  $x_\alpha \in L_\alpha$  and  $y_\alpha \in L_{-\alpha}$ . Let  $[x_\alpha, y_\alpha] \in H = L_0$ . By ad-symmetry of  $K$ , one obtains

$$\begin{aligned} K([x_\alpha, y_\alpha], y_\alpha) &= \alpha(h) K(x_\alpha, y_\alpha) \\ &= K(h, [x_\alpha, y_\alpha]) \\ &= K(h, h_\alpha) K(x_\alpha, y_\alpha) \\ &= K(h, K(x_\alpha, y_\alpha) h_\alpha) \end{aligned}$$

By non-degeneracy of  $K$ ,

$$[x_\alpha, y_\alpha] = K(x_\alpha, y_\alpha) h_\alpha,$$

since they both define the same linear functional  $K(h, \cdot)$  on  $H$ .

4. We conclude from the above that for all  $x \in L_\alpha$  and  $y \in L_{-\alpha}$ ,

$$[x, y] = K(x, y) h_\alpha$$

and so  $\dim[L_\alpha, L_{-\alpha}] = 1$  and it's spanned by  $h_\alpha$ , provided we show that for  $x \neq 0$  in  $L_\alpha$  there is some  $y \in L_{-\alpha} \setminus \{0\}$  such that  $K(x, y) \neq 0$ . But, if  $L_\alpha \perp L_{-\alpha}$  then we claim that  $L_\alpha \perp L$ . Since  $L_\alpha \perp L_\beta$  for  $\beta \neq -\alpha$ , and so if  $L_\alpha \perp L_{-\alpha}$  in addition, then  $L_\alpha \perp L$  and this contradicts non-degeneracy.

---

<sup>6</sup>by the decomposition of  $L$  to a direct sum

5. Assume towards a contradiction that

$$K(\alpha, \alpha) = \alpha(h_\alpha) = 0$$

and consider the three-dimensional algebra generated by  $x \in L_\alpha \setminus \{0\}$ ,  $y \in L_{-\alpha} \setminus \{0\}$  and  $[x, y] = c \cdot h_\alpha$  with  $c \neq 0$ .<sup>7</sup> Now,

$$\begin{aligned} [h_\alpha, x] &= \alpha(h_\alpha)x = 0 \\ [h_\alpha, y] &= -\alpha(h_\alpha)y = 0 \end{aligned}$$

since we assume  $\alpha(h_\alpha) = 0$ .

Let  $M$  be the sub-algebra and we see that  $h_\alpha$  is in its center. But  $\dim M/Z(M) \leq 2$  and  $M/Z(M)$  is solvable, and hence so is  $M$ . So  $[M, M]$  is a nilpotent Lie algebra.<sup>8</sup> Consider

$$\text{ad}_L : M \rightarrow \text{End } L.$$

$\text{ad}_L[M, M]$  is a nilpotent Lie algebra of linear transformations, so  $\text{ad}_L w$  is a nilpotent linear transformation for all  $w \in [M, M]$ . But  $ch_\alpha = [x, y] \neq 0$  is in  $[M, M]$  and  $\text{ad}_L h_\alpha$  is then both diagonalisable and nilpotent. Hence  $h_\alpha$ , in contradiction.

6. Let  $x \neq 0$  in  $L_\alpha$ , by part 4 there is  $y \in L_{-\alpha}$  such that  $K(x, y) \neq 0$ . We can choose  $x_\alpha, y_\alpha$  such that  $K(x_\alpha, y_\alpha) = \frac{2}{K(\alpha, \alpha)}$  by 5 and by multiplying  $y$  by a non-zero scalar. Now compute:

$$\begin{aligned} [x_\alpha, y_\alpha] &= K(K(x_\alpha, y_\alpha))h_\alpha \\ &= \frac{2h_\alpha}{K(\alpha, \alpha)} =: \hat{h}_\alpha \end{aligned}$$

and

$$\begin{aligned} [\hat{h}_\alpha, x_\alpha] &= \alpha(\hat{h}_\alpha)x_\alpha = \frac{2\alpha(h_\alpha)}{K(\alpha, \alpha)}x_\alpha = 2x_\alpha \\ [\hat{h}_\alpha, y_\alpha] &= \frac{-2\alpha(h_\alpha)}{K(\alpha, \alpha)}y_\alpha. \end{aligned}$$

So, the multiplication table of  $\langle x_\alpha, y_\alpha, \hat{h}_\alpha \rangle = M$  is

$$\begin{aligned} [\hat{h}_\alpha, x_\alpha] &= 2x_\alpha \\ [\hat{h}_\alpha, y_\alpha] &= -2y_\alpha \\ [x_\alpha, y_\alpha] &= \hat{h}_\alpha. \end{aligned}$$

This is exactly the multiplication table of another three-dimensional algebra we know, named  $\mathfrak{sl}_2(\mathbb{F})$ , with isomorphism given by

$$\begin{aligned} x_\alpha &\mapsto \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \\ y_\alpha &\mapsto \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \\ h_\alpha &\mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned}$$

7. We already saw that  $\hat{h}_\alpha = \frac{2h_\alpha}{K(\alpha, \alpha)}$  and  $h_{-\alpha} = -h_\alpha$  implies  $\hat{h}_{-\alpha} = -\hat{h}_\alpha$ .  $h_\alpha$  is defined by  $K(h, h_\alpha) = \alpha(h)$ .  
■

<sup>7</sup>This is indeed an algebra, since  $x, y$  are eigenvectors of each  $\text{ad } h$ , and in particular  $\text{ad } h_\alpha$ .

<sup>8</sup>Deduce this from Lie's theorem.

## 7.2 Rationality properties of the root system

Assume throughout  $L$  is a semi-simple Lie algebra of an algebraically closed field  $\mathbb{F}$  of characteristic zero.

**Proposition 7.2.1.** *Let  $\alpha, \beta \in \Phi$ . Then*

$$K(\alpha, \beta) = q_{\alpha, \beta} K(\alpha, \alpha)$$

where  $q_{\alpha, \beta} \in \mathbb{Q}$ .

*Proof.* Consider the vector space<sup>9</sup>

$$L_{\alpha, \beta} = \sum_{k \in \mathbb{Z}} L_{\beta + k\alpha} \subseteq L.$$

Now take  $x_\alpha \in L_\alpha$  and  $y_\alpha \in L_{-\alpha}$ , and take  $\hat{h}_\alpha = [x_\alpha, y_\alpha]$  which we know generates a three-dimensional subalgebra,  $S_\alpha \leq L$ , which is isomorphic to  $\mathfrak{sl}_2(\mathbb{F})$ .  $L_{\alpha, \beta}$  is invariant under the action of  $S_\alpha$  by the adjoint, as shown in the following paragraph.

$$\forall w \in S_\alpha: \text{ad}_L(w) L_{\alpha, \beta} \subseteq L_{\alpha, \beta}.$$

Indeed, for all  $z \in L_\gamma$ ,

$$\text{ad}_L(z) L_{\beta + k\alpha} \subseteq L_{\beta + k\alpha + \gamma}.$$

Hence,  $\text{ad}_L x_\alpha$  and  $\text{ad}_L y_\alpha$  leave  $L_{\alpha, \beta}$  invariant, and so does  $\text{ad}_L \hat{h}_\alpha$ , since  $L_{\alpha, \beta}$  is a sum of eigenspaces.

Now,  $\text{ad}_L(S_\alpha)$  acts on  $L_{\alpha, \beta}$ , and  $\hat{h}_\alpha = [x_\alpha, y_\alpha]$  is a commutator! Henceforward,  $\text{ad}_L(h_\alpha)$  must have trace zero, when restricted to  $L_{\alpha, \beta}$ .<sup>10</sup> So, we compute using the diagonalisation of  $\text{ad}_L(h_\alpha)$  in  $L_{\alpha, \beta}$ .

$$\begin{aligned} \text{tr ad}_L(h_\alpha) &= 0 \\ &= \sum_{k \in \mathbb{Z}} (\beta + k\alpha)(h_\alpha) \dim L_{\beta + k\alpha} \\ &= \sum_{k \in \mathbb{Z}} (\beta(h_\alpha) + k\alpha(h_\alpha)) d_k \end{aligned}$$

where we denote  $d_k := \dim L_{\beta + k\alpha}$ . We get by substituting  $\beta(h_\alpha) = K(\beta, \alpha)$  and  $\alpha(h_\alpha) = K(\alpha, \alpha)$  the following.

$$\left( - \sum_{k \in \mathbb{Z}} d_k \right) K(\beta, \alpha) = \sum_{k \in \mathbb{Z}} k d_k K(\alpha, \alpha)$$

Then

$$K(\beta, \alpha) = \frac{- \sum_{k \in \mathbb{Z}} k d_k}{\sum_{k \in \mathbb{Z}} d_k} K(\alpha, \alpha) = q_{\alpha, \beta} K(\alpha, \alpha)$$

where  $\sum_{k \in \mathbb{Z}} d_k \geq 1$  since  $d_k \geq 0$  and  $d_0 > 0$ . We get  $q_{\alpha, \beta} \in \mathbb{Q}$ . ■

**Proposition 7.2.2.**  *$K(\alpha, \alpha)$  is a positive rational number, and  $K(\beta, \alpha)$  is rational for all  $\alpha, \beta \in \Phi$ .*

*Proof.* Let us compute directly from the definition of the Killing form, and the diagonalisation of  $L$  with respect to  $H$ .

$$\begin{aligned} K(\alpha, \beta) &= K(h_\alpha, h_\beta) \\ &= \text{tr}(\text{ad}_L(h_\alpha), \text{ad}_L(h_\beta)) \\ &= \sum_{\lambda \in \Phi} \lambda(h_\alpha) \lambda(h_\beta) \dim L_\lambda. \end{aligned}$$

Now,

$$\begin{aligned} K(\alpha, \alpha) &= \sum_{\lambda \in \Phi} \lambda(h_\alpha)^2 \dim L_\lambda \\ &= \sum_{\beta \in \Phi} \beta(h_\alpha)^2 \dim L_\beta \\ &= \sum_{\beta \in \Phi} K(\beta, \alpha)^2 \dim L_\beta \\ &\stackrel{\text{by previous proposition}}{=} \sum_{\beta \in \Phi} q_{\alpha, \beta}^2 K(\alpha, \alpha)^2 \dim L_\beta. \end{aligned}$$

<sup>9</sup>there are finitely many roots, hence there is much redundancy in the sum

<sup>10</sup>which is invariant also under  $\text{ad}_L x_\alpha$  and  $\text{ad}_L y_\alpha$

Isolating  $K(\alpha, \alpha)$  from this equation one gets

$$K(\alpha, \alpha) = \frac{1}{\sum_{\beta \in \Phi} q_{\alpha, \beta}^2 \dim L_\beta} \in \mathbb{Q}_{>0}$$

as required.<sup>11</sup> ■

**Corollary 7.2.3.**  $(K(\alpha, \beta))_{\alpha, \beta \in \Phi}$  is a rational matrix, with positive diagonal terms.

**Proposition 7.2.4.** 1.  $\dim L_\alpha = 1$  for all  $\alpha \in \Phi$ .

2. No multiple of  $\alpha$  by an integer is also a root, except for  $-\alpha$ .

*Proof.* Choose  $y \in L_{-\alpha}$  and consider the space

$$M_\alpha = \mathbb{F}y_\alpha + \mathbb{F}h_\alpha + \sum_{k \geq 1} L_{k\alpha}.$$

We claim that  $M_\alpha$  is a linear sub-space invariant under  $S_\alpha = \text{span}\{x_\alpha, y_\alpha, h_\alpha\} \cong \mathfrak{sl}_2(\mathbb{F})$ .

We saw  $[L_\alpha, L_{-\alpha}] = 1$  is spanned by  $h_\alpha$  and is one-dimensional. We get a few properties using this.

(i)

$$\text{ad}_L h_\alpha: M_\alpha \rightarrow M_\alpha$$

(ii)  $\text{ad}_L x_\alpha: M_\alpha \rightarrow M_\alpha$  since

$$\begin{aligned} [x_\alpha, y_\alpha] &= ch_\alpha \in \mathbb{F}h_\alpha \\ [x_\alpha, h] &= -\alpha(h) \cdot x_\alpha \in L_\alpha \\ [x_\alpha, x_{k\alpha}] &\in L_{(k+1)\alpha} \end{aligned}$$

(iii) Finally,  $\text{ad}_L y_\alpha: M_\alpha \rightarrow M_\alpha$ , and

$$\begin{aligned} [y_\alpha, y_\alpha] &= 0 \\ [y_\alpha, x_\alpha] &= c \cdot h_\alpha \end{aligned}$$

so

$$[y_\alpha, x_{k\alpha}] = \text{ad}_L(y_\alpha)(x_{k\alpha}) \in L_{(k-1)\alpha}.$$

From these,  $M_\alpha$  is  $\text{ad}_L(S_\alpha)$ -invariant. Once again  $ch_\alpha = [x_\alpha, y_\alpha]$  is a commutator, so it has trace 0 on  $M_\alpha$ . We obtain the following.

$$\begin{aligned} 0 &= \text{tr ad}_L(h_\alpha)|_{M_\alpha} \\ &= -\alpha(h_\alpha) + 0 + \sum_{k \geq 1} k\alpha(h_\alpha) \dim L_{k\alpha} \end{aligned}$$

Let  $b_k = \dim L_{k\alpha}$ , we obtain from the above equality

$$\alpha(h_\alpha) \left( -1 + \sum_{k \geq 1} kb_k \right) = 0.$$

But,  $\alpha(h_\alpha) = K(\alpha, \alpha) \neq 0$  and so  $\sum_{k \geq 1} kb_k = 1$ .

Then  $b_k \geq 0$  for  $k \geq 1$ , and  $b_1 > 0$ . Then  $b_1 = \dim L_\alpha = 1$ ,  $\dim L_{k\alpha} = 0$  for  $k > 1$ . From  $\Phi = -\Phi$  we get  $L_{k\alpha} = 0$  for all  $k \neq \pm 1, 0$ . ■

**Definition 7.2.5.** Let  $\alpha, \beta \in \Phi$ . Define the  $\alpha$ -chain through  $\beta$  to be the set of roots of the form

$$\beta + k\alpha, \quad k \in \mathbb{Z}.$$

---

<sup>11</sup>Then  $K(\beta, \alpha)$  is also rational, by the previous proposition.

**Theorem 7.2.6.** 1. The  $\alpha$ -chain through  $\beta$  has the form

$$\beta + k\alpha, \quad p \leq k \leq q.$$

It is one (uninterrupted) interval.

2.

$$p + q = \frac{-2\beta(h_\alpha)}{\alpha(h_\alpha)} = \frac{-2K(\beta, \alpha)}{K(\alpha, \alpha)}.$$

*Proof.* Let  $r$  and  $s$  define a maximal interval on the  $\alpha$ -chain through  $\beta$ . I.e.

$$\begin{aligned} \forall r \leq j \leq s: \beta + j\alpha &\in \Phi \\ \beta + (s+1)\alpha, \beta + (r-1)\alpha &\notin \Phi. \end{aligned}$$

Let

$$W_{\alpha, \beta} = \bigoplus_{j=r}^s L_{\beta+j\alpha}.$$

This is a non-trivial sub-space. Let  $x_\alpha \in L_\alpha, y_\alpha \in L_{-\alpha}$  be non-zero and  $h_\alpha = c[x_\alpha, y_\alpha]$  with  $c \neq 0$ . Now

$$S_\alpha = \mathbb{F}x_\alpha + \mathbb{F}y_\alpha + \mathbb{F}h_\alpha$$

is a Lie sub-algebra (isomorphic to  $\mathfrak{sl}_2(F)$ ).

We claim that  $W_{\alpha, \beta}$  is invariant under  $\text{ad}_L(S_\alpha)$ . Examine

$$\text{ad}_L(x_\alpha) : W_{\alpha, \beta} \rightarrow W_{\alpha, \beta}.$$

We have

$$\begin{aligned} \text{ad}_L(x_\alpha) L_\gamma &\subseteq L_{\gamma+\alpha} \\ \text{ad}_L(y_\alpha) L_\gamma &\subseteq L_{\gamma-\alpha} \end{aligned}$$

and so  $W_{\alpha, \beta}$  is invariant under  $\text{ad}_L x_\alpha$  and  $\text{ad}_L(y_\alpha)$ , then under  $\text{ad}_L(L_\alpha)$ .  $\text{ad}_L(h_\alpha)|_{W_{\alpha, \beta}}$  is a commutator, hence has trace zero.

$$\text{tr ad}_L(h_\alpha)|_{W_{\alpha, \beta}} = 0$$

Hence,

$$\sum_{j=r}^s (\beta + j\alpha)(h_\alpha) = \sum_{j=r}^s \beta(h_\alpha) + \sum_{j=r}^s j\alpha(h_\alpha) = 0.$$

Moving sides, we get the following.

$$\begin{aligned} -(s-r+1)\beta(h_\alpha) &= \left( \sum_{j=r}^s j \right) \alpha(h_\alpha) \\ &= \left( r(s-r+1) + \sum_{j=1}^{s-r} j \right) \alpha(h_\alpha) \\ &= \left( r(s-r+1) + \frac{(s-r)(s-r+1)}{2} \right) \alpha(h_\alpha) \end{aligned}$$

Moving sides we get

$$-\beta(h_\alpha) = \frac{s+r}{2} \alpha(h_\alpha).$$

Now  $\alpha(h_\alpha) \neq 0$  because we've seen  $\forall \gamma \in \Phi: \dim L_\gamma = 1$ . Hence

$$s+r = \frac{2\beta(h_\alpha)}{\alpha(h_\alpha)}.$$

This is true for all maximal intervals, and clearly this means that there is only one such an interval.<sup>12</sup> ■

<sup>12</sup>An interval to the right should have larger  $s+r$ , and one to the left should have smaller  $s+r$ .



**Theorem 7.2.7.** 1. The only roots  $c\alpha \in \Phi$ <sup>13</sup> which are multiples of  $\alpha$  are 0 and  $\pm\alpha$ .

2. For  $\alpha, \beta \in \Phi$ ,

$$L_{\alpha+\beta} = [L_\alpha, L_\beta].$$

**Remark 7.2.8.** 1. We already know that no integral multiples of  $\alpha$  are roots, except for  $\pm\alpha$  and 0.

2. We already know that  $[L_\alpha, L_\beta] \subseteq L_{\alpha+\beta}$ , and we want to show equality.

*Proof.* 1. Let us consider the  $\alpha$ -chain through  $\beta$ , which is of the form  $\beta + j\alpha$  with  $p \leq j \leq q$ , and where  $p + q = \frac{-2\beta(h_\alpha)}{\alpha(h_\alpha)}$  as we saw. Now let  $\beta = c\alpha \neq \pm\alpha, 0$ . Then,

$$p + q = \frac{-2K(\beta, \alpha)}{K(\alpha, \alpha)} = \frac{-2K(\alpha, \alpha)}{K(\alpha, \alpha)} = -2c \in \mathbb{Z}.$$

So,  $c = n + \frac{1}{2}$  for some  $n \in \mathbb{Z}$ .<sup>14</sup> So,  $(n + \frac{1}{2})\alpha \in \Phi$  then  $-(n + \frac{1}{2})\alpha \in \Phi - \Phi$ . But,

$$-\left(n + \frac{1}{2}\right)\alpha = \left(n + \frac{1}{2}\right)\alpha - (2n + 1)\alpha = \beta - (2n + 1)\alpha.$$

Therefore,  $-(n + \frac{1}{2})\alpha$  is in the  $\alpha$ -chain through  $\beta$ . Namely, it is  $\beta + k\alpha$  for some  $k \in \mathbb{Z}$ . So, the chain has  $-(2n + 1)$  in it, and also 0.

Denote

$$\begin{aligned}\gamma_1 &= \left(n + \frac{1}{2}\right)\alpha \\ \gamma_2 &= \left(n + \frac{1}{2}\right)\alpha\end{aligned}$$

both of which are in  $\Phi$ . Then

$$\gamma_1 = \gamma_2 + (2n + 1)\alpha$$

and so  $\gamma_1$  is in the  $\alpha$ -chain through  $\gamma_2$ . That  $\alpha$ -chain contains  $\gamma_2$  and  $\gamma_1 = \gamma_2 + (2n + 1)\alpha$ , and so the interval of that chain contains  $[0, 2n + 1]$ .<sup>15</sup> So 1 is also in the interval, and so

$$\gamma_2 + \alpha = \left(-n + \frac{1}{2}\right)\alpha + \alpha = \left(-(n - 1) + \frac{1}{2}\right)\alpha \in \Phi.$$

We can continue this way, and finally get that  $\frac{1}{\alpha} \in \frac{1}{2}\alpha$ . Then  $\alpha$  is a multiple by 2 of  $\frac{1}{2}\alpha$ ! this is a contradiction, since no non-trivial integer multiple of a root can be a root. We argue similarly for  $n < 0$ . We conclude that no multiple of a root is a root except 0 and  $\pm\alpha$ .

2. If  $\alpha + \beta \notin \Phi$ , then  $L_{\alpha+\beta} = 0$ .

So, assume towards a contradiction that  $[L_\alpha, L_\beta] \subsetneq L_{\alpha+\beta}$ , and  $\alpha + \beta$  is a root. Because the dimension of  $L_{\alpha+\beta}$  is 1, this means  $[L_\alpha, L_\beta] = 0$ .

Let  $p, q$  define the interval which is the  $\alpha$ -chain through  $\beta$ . We assume  $j \in [p, q]$  and  $q \geq 1$ , since  $\beta + \alpha$  is a root by assumption. We also assumed  $[L_\alpha, L_\beta] = 0$ , namely

$$\text{ad}_L(x_\alpha)(L_\beta) = 0.$$

Hence  $x_\alpha \in L_\alpha \setminus \{0\}$ . So, define

$$U_{\alpha, \beta} = \bigoplus_{j=p}^0 L_{\beta+j\alpha}$$

which is invariant under  $S_\alpha$ .<sup>16</sup>

$\text{ad}_L(x_\alpha)$  sends  $L_{\beta+j\alpha}$  to  $L_{\beta+(j+1)\alpha}$  and  $L_\beta$  to 0.  $\text{ad}_L(y_\alpha)$  sends  $L_{\beta+\alpha}$  to  $L_{\beta+(j-1)\alpha}$  and  $L_{\beta+j\alpha}$  to 0.

<sup>13</sup> $c \in \mathbb{F}$

<sup>14</sup>for we saw  $c \notin \mathbb{Z}$

<sup>15</sup>Here we assume  $n \geq 0$ , but the statement is symmetric.

<sup>16</sup>This is well-defined because  $\beta$  is a root and hence  $p \leq 0$ .

Of course,  $U_{\alpha,\beta}$  is invariant under  $[x_\alpha, y_\alpha] = ch_\alpha$ , so it is invariant under  $S_\alpha$ , and let us compute the trace.

$$\begin{aligned} 0 &= \text{tr ad}_L(h_\alpha)|_{U_{\alpha,\beta}} \\ &= \sum_{j=p}^0 (\beta + j\alpha)(h_\alpha) \\ &= \sum_{j=p}^0 \beta(h_\alpha) + \sum_{j=p}^0 j\alpha(h_\alpha) \end{aligned}$$

Then

$$-(|p|+1)\beta(h_\alpha) = -\frac{|p|(|p|+1)}{2}\alpha(h_\alpha)$$

and then

$$\frac{-2\beta(h_\alpha)}{\alpha(h_\alpha)} = -|p| = p \neq p+q$$

because  $q \geq 1$ . This is a contradiction. ■

Now we exploit the rational properties of  $\Phi \subseteq H^* = \text{Hom}(H, \mathbb{F})$ . We plan to define a positive definite bilinear form on a vector space over  $\mathbb{Q}$ , and then extend to a vector space over  $\mathbb{R}$ . Associated to this vector space will be a finite set of points in Euclidean space, the root system, with very special properties.

**Proposition 7.2.9.** *Choose  $\alpha_1, \dots, \alpha_\ell$  a basis of  $H^*$  consisting of roots. Then for every  $\beta \in \Phi$*

$$\beta = \sum_{i \in [\ell]} c_i \alpha_i$$

where  $c_i$  are rational numbers.

*Proof.* We plug the linear equation and obtain.

$$K(b, \alpha_j) = \sum_{i \in [\ell]} c_i K(\alpha_i, \alpha_j)$$

Then  $(K(\alpha_i, \alpha_j))_{i,j \in [\ell]}$  is a symmetric matrix of rational number which is *non-singular*. ■

So,  $\{c_i\}_{i \in [\ell]}$  are uniquely determined and rational. ■

*Proof.* Let  $E_{\mathbb{Q}}$  be the  $\mathbb{Q}$  span of  $\Phi$ . ■

*Proof.* Let  $\ell = \dim_{\mathbb{F}} H$ . Then  $\dim_{\mathbb{Q}} E_{\mathbb{Q}} = \ell$ . ■

*Proof.* First, since  $\alpha_1, \dots, \alpha_\ell$  are linearly independent over  $\mathbb{F}$ , they are linearly independent over  $\mathbb{Q}$ . So,  $\dim_{\mathbb{Q}} E_{\mathbb{Q}} \geq \ell$ . But each  $\beta \in \mathbb{Q}$  is a rational linear combination of  $(\alpha_i)_{i \in [\ell]}$ , as we saw, and so  $\dim_{\mathbb{Q}} E_{\mathbb{Q}} = \ell$ . ■

**Remark 7.2.10.** We already that that  $K$  restricted to  $E_{\mathbb{Q}}$  satisfies  $K(\alpha, \alpha) > 0$  for all  $\alpha \in \Phi$ .

**Proposition 7.2.11.**  *$K$  is positive definite on  $E_{\mathbb{Q}}$ .*

*Proof.* Let  $\lambda, \mu \in H^*$ . Then

$$K(\lambda, \mu) = \sum_{\alpha \in \Phi} \alpha(h_\lambda) \alpha(h_\mu) = K(h_\lambda, h_\mu) = \text{tr}(\text{ad}_L h_\lambda \text{ad}_L(h_\mu)) = \sum_{\alpha \in \Phi} K(\alpha, \lambda) K(\alpha, \mu)$$

because  $\dim L_\alpha = 1$ . For  $\lambda = \mu = E_{\mathbb{Q}}$  we have

$$K(\lambda, \mu) = \sum_{\alpha \in \Phi} K(\alpha, \lambda)^2 > 0$$

because  $K(\alpha, \lambda)$  is a rational number, and  $K$  is non-degenerate and  $\Phi$  spans  $H^*$  and  $E_{\mathbb{Q}}$ . ■

**Fact 7.2.12.** *The positive definite symmetric form  $K$  on the rational vector space  $E_{\mathbb{Q}}$  can be extended to such a form on an Euclidean space over  $\mathbb{R}$ , of the same dimension. Formally one can consider  $\mathbb{R} \otimes_{\mathbb{Q}} E_{\mathbb{Q}}$ .*

$$K(\alpha \otimes \lambda, b \otimes \mu) = abK(\lambda, \mu)$$

is the form.

Denote  $\Phi = 1 \otimes \Phi$  the set of roots.

**Proposition 7.2.13.** *We now have an  $\ell$ -dimensional Euclidean space, with a positive definite symmetric form  $(\cdot, \cdot)$ , and a finite set  $\Phi$  of roots which satisfies the following.*

- (1)  $\Phi = -\Phi$  and  $\Phi$  spans  $E$ .
- (2)  $\Phi \cap \mathbb{R}\alpha = \{\pm\alpha, 0\}$  for all  $\alpha \in \Phi$ .
- (3) If  $\alpha, \beta \in \Phi$  then  $\frac{2(\alpha, \beta)}{(\alpha, \alpha)} \in \mathbb{Z}$ .
- (4) If  $\alpha, \beta \in \Phi$  then  $\beta - \frac{2(\alpha, \beta)}{(\alpha, \alpha)}\alpha \in \Phi$ .

*Proof.* Properties one to three are clear. We shall prove (4). The  $\alpha$ -chain through  $\beta$  is

$$\beta + p\alpha, \dots, \beta, \dots, \beta + q\alpha$$

with  $p \leq 0 \leq q$ . Now we saw that  $\frac{-2(\alpha, \beta)}{(\alpha, \alpha)} = p + q$ . So all we say is that  $p \leq p + q \leq q$ , which is clear because  $p \leq 0 \leq q$ . ■

### 7.3 Geometry of Euclidean root systems

Let  $E$  be an  $\ell$ -dimensional Euclidean space, with  $(\cdot, \cdot)$  a symmetric positive-definite form.

Given any  $\alpha \in E \setminus \{0\}$ , we can define the reflection determined by  $\alpha$  on  $E$ . It is an orthogonal linear map of order 2 and determinant  $-1$ , defined by

$$\sigma_\alpha(\beta) = \beta - \frac{2(\alpha, \beta)}{(\alpha, \alpha)}\alpha.$$

Clearly,  $\sigma_\alpha(\alpha) = \alpha - 2\alpha = -\alpha$ , and on  $P_\alpha = \{\beta \mid (\beta, \alpha) = 0\}$ . The hyperplane orthogonal to  $\alpha$ , and we have

$$\sigma_\alpha(\beta) = \beta. \text{ So, } \sigma_\alpha \text{ has the matrix } \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix} \text{ in a basis of } \mathbb{R}\alpha \text{ and } P_\alpha.$$

We conclude that the reflections  $\sigma_\alpha$  with  $\alpha \in \Phi$  leave  $\Phi$  invariant.

**Definition 7.3.1.** The group generated by  $(\sigma_\alpha)_{\alpha \in \Phi}$  in the orthogonal group preserving the form, is called the **Weyl group of the root system**, and is denoted  $W(\Phi) = W$ .

$W(\Phi)$  is a finite group since each  $w \in W$  has a permutation representation on  $\Phi$ , and this representation is faithful, since  $\Phi$  spans  $E$ . So,  $W(\Phi) \hookrightarrow S(\Phi)$ .

### 7.4 Pairs of roots in a root system

Let  $\Phi$  be a root system and let  $\alpha, \beta \in \Phi$ . We know that

$$(\alpha, \beta) = \|\alpha\| \|\beta\| \cos(\theta(\alpha, \beta)).$$

We know also that

$$2 \frac{(\alpha, \beta)}{(\alpha, \alpha)} = 2 \frac{\|\beta\|}{\|\alpha\|} \cos(\theta(\alpha, \beta)) \in \mathbb{Z}.$$

So, consider  $\frac{(\alpha, \beta)}{(\alpha, \alpha)}$  and  $\frac{(\alpha, \beta)}{(\beta, \beta)}$ . We conclude that

$$4 \frac{(\alpha, \beta)}{(\alpha, \alpha)} \frac{(\alpha, \beta)}{(\beta, \beta)} = 4 \cos^2 \theta(\alpha, \beta) \in \mathbb{Z}.$$

Of course,  $(\alpha, \beta)$  and  $(\beta, \alpha)$  have the same sign, and are zero together. Hence  $4 \cos^2 \theta$  is a product of two integers of the same sign. Of course,  $0 \leq 4 \cos^2 \theta \leq 4$ . Now, 4 corresponds to  $\cos \theta = \pm 1$  so  $\theta \in \{0, \pi\}$ , then  $\alpha = \pm\beta$ . Otherwise  $4 \cos^2 \theta < 4$ . Then  $4 \cos^2 \theta \in \{0, 1, 2, 3\}$ . We classify our results in table 7.1.

**Proposition 7.4.1.** *Let  $\alpha, \beta \in \Phi$  with  $\alpha \neq \pm\beta$ . If  $(\alpha, \beta) > 0$  then  $\alpha - \beta$  is a root and if  $(\alpha, \beta) < 0$ , then  $\alpha + \beta$  is a root. So if the angle between  $\alpha, \beta$  is acute, their difference is a root, and if it's obtuse, then their sum is.*

Figure 7.1: Angles between roots.

$\frac{2(\alpha, \beta)}{(\alpha, \alpha)}$	$\frac{2(\alpha, \beta)}{(\beta, \beta)}$	$\theta$	$\frac{\ \beta\ ^2}{\ \alpha\ ^2}$
0	0	$\frac{\pi}{2}$	?
1	1	$\frac{\pi}{3}$	1
-1	-1	$\frac{2\pi}{3}$	1
1	2	$\frac{\pi}{4}$	$\frac{1}{2}$
-1	-2	$\frac{3\pi}{4}$	$\frac{1}{2}$
1	3	$\frac{\pi}{6}$	$\frac{1}{3}$
-1	-3	$\frac{5\pi}{6}$	$\frac{1}{3}$

*Proof.* First, the second part follows from the first and taking  $-\beta$  while using  $\Phi = -\Phi$ . Now, if  $(\alpha, \beta) > 0$ , the table shows that at least one of  $\frac{2(\alpha, \beta)}{(\alpha, \alpha)}$  and  $\frac{2(\alpha, \beta)}{(\beta, \beta)}$  is equal to 1.

Then, if  $\frac{2(\alpha, \beta)}{(\alpha, \alpha)} = 1$ , then  $\sigma_\alpha(\beta) = \beta - \frac{2(\alpha, \beta)}{(\alpha, \alpha)}\alpha = \beta - \alpha \in \Phi$ , because  $\Phi$  is invariant under  $\sigma_\alpha$  with  $\alpha \in \Phi$ . Then also  $-(\beta - \alpha) = \alpha \in \beta \in \Phi$  and if  $\frac{2(\alpha, \beta)}{(\beta, \beta)} = 1$  then

$$\sigma_\beta(\alpha) = \alpha - \frac{2(\alpha, \beta)}{(\beta, \beta)}\beta = \alpha - \beta \in \Phi. \quad \blacksquare$$

## 7.5 The action of the reflection $\sigma_\alpha$ on the $\alpha$ -chain through $\beta$

Recall that  $\sigma_\alpha(\beta) = \beta - \frac{2(\alpha, \beta)}{(\alpha, \alpha)}\alpha$ , so clearly  $\sigma_\alpha(\beta)$  is in the  $\alpha$ -chain through  $\beta$ <sup>17</sup>. So,  $\sigma_\alpha$  must take the  $\alpha$ -chain through  $\beta$  to itself, being an invariant finite subset.  $\sigma_\alpha$  reverses the orientation of the line  $\mathbb{R}\alpha$ , and so if the  $\alpha$ -chain through  $\beta$  is the interval  $[p, q]$ , then  $\sigma_\alpha$  flips it. So,

$$\sigma_\alpha(p + q\alpha) = \beta + p\alpha = \beta - \frac{2(\beta, \alpha)}{(\alpha, \alpha)}\alpha - q\alpha.$$

We get in particular our known fact

$$\frac{-2(\beta, \alpha)}{(\alpha, \alpha)} = (p + q).$$

**Definition 7.5.1.** A basis  $(\alpha_i)_{i \in [\ell]}$  of  $E$  is called a **basis of the root system** if every root  $\beta \in \Phi$  has a representation  $\beta = \sum_{i \in [\ell]} a_i \alpha_i$  where all  $a_i$  are non-negative or all  $a_i$  are non-positive.

**Theorem 7.5.2.** Every root system has a basis.

**Definition 7.5.3.** The roots  $(\alpha_i)_{i \in [\ell]}$  in a basis of the root system are called **simple roots**.

**Definition 7.5.4.** If  $\beta = \sum_{i \in [\ell]} a_i \alpha_i$  and  $\forall i \in [\ell] : a_i \geq 0$  then  $\beta$  is called a **positive root**. Similarly one defines **negative roots**.

**Definition 7.5.5.**  $\Phi^+$  and  $\Phi^-$  are the sets of positive and negative roots, respectively.

**Remark 7.5.6.**  $\Phi = \Phi^+ \amalg \Phi^-$ .

**Proposition 7.5.7.** If  $(\alpha_i)_{i \in [\ell]}$  form a basis of the root system, then all the mutual angles satisfy  $(\alpha_i, \alpha_j) \leq 0$ .

*Proof.* If  $(\alpha_i, \alpha_j) > 0$ , as we saw  $\alpha_i - \alpha_j$  is a root, but has neither all-positive nor all-negative coefficients, in contradiction.  $\blacksquare$

<sup>17</sup>The action of  $\sigma_\alpha$  is by adding multiples of  $\alpha$ .

To construct a basis of the root systems, consider

$$\bigcup_{\alpha \in \Phi} P_\alpha = \bigcup_{\alpha \in \Phi} \{\beta \mid (\beta, \alpha) = 0\},$$

a union of finitely many proper hyperplanes.

Choose  $\gamma$  *outside* this set, so  $\forall \alpha \in \Phi: (\gamma, \alpha) \neq 0$ .

Define

$$\Phi^+(\gamma) := \{\alpha \in \Phi \mid (\gamma, \alpha) > 0\}.$$

Clearly  $\Phi = \Phi^+(\gamma) \amalg -\Phi^+(\gamma)$ .

**Definition 7.5.8.** A root  $\alpha \in \Phi^+(\gamma)$  is called **irreducible** if it is *not* a sum of the other roots in  $\Phi^+(\gamma)$ . Equivalently,  $\alpha$  is reducible if  $\alpha = \beta + \beta'$  where  $\beta, \beta' \in \Phi^+(\gamma)$ .

**Theorem 7.5.9.** The irreducible roots in  $\Phi^+(\gamma)$  form a basis of the root system.

*Proof.* I. Every  $\alpha \in \Phi^+(\gamma)$  is a sum of irreducible roots in  $\Phi^+(\gamma)$  with non-negative coefficients.

We argue by contradiction. Otherwise there's  $\alpha \in \Phi^+(\gamma)$  which isn't a sum of irreducible roots with non-negative coefficients. Choose  $\alpha$  such that  $(\alpha, \gamma) > 0$  is minimal among those.  $\alpha$  itself cannot be irreducible, so  $\alpha$  is reducible and write  $\alpha = \beta_1 + \beta_2$  with  $\beta_1, \beta_2 \in \Phi^+(\gamma)$ . However,  $(\alpha, \gamma) = (\beta_1, \gamma) + (\beta_2, \gamma)$ . If  $\beta_1, \beta_2$  are both non-negative linear combinations of irreducible roots, the same is true for  $\alpha$  and we're done.

Assume therefore that at least one of them, WLOG  $\beta_1$  is a root, but not a non-negative linear combination of irreducible roots in  $\Phi^+(\gamma)$ . Then

$$0 < (\gamma, \beta_1) < (\gamma, \alpha)$$

in contradiction to the minimality of  $\alpha$ .

II. If  $\alpha, \beta \in \Phi^+(\gamma)$  are different and irreducible, then  $(\alpha, \beta) \leq 0$ .

Assume otherwise, i.e. that  $(\alpha, \beta) > 0$ . Then  $\alpha - \beta$  and  $\beta - \alpha$  are roots, as we saw. Hence, one of them is in  $\Phi^+(\gamma)$ . WLOG  $\alpha - \beta \in \Phi^+(\gamma)$ . Then  $\alpha = \alpha - \beta + \beta$  and so  $\alpha$  is reducible, contradicting our assumption.

III. The set of irreducible roots in  $\Phi^+(\gamma)$  is linearly independent.

Denote this set  $\Delta^+(\gamma)$  and assume towards a contradiction that

$$\sum_{\alpha \in \Delta^+(\gamma)} r_\alpha \alpha = 0.$$

Let

$$\begin{aligned} \Delta_1^+(\gamma) &= \{\alpha \mid r_\alpha < 0\} \\ \Delta_2^+(\gamma) &= \{\alpha \mid r_\alpha > 0\}. \end{aligned}$$

Changing notations we get

$$\sum_{\alpha \in \Delta_2^+(\gamma)} t_\alpha \alpha - \sum_{\beta \in \Delta_1^+(\gamma)} s_\beta \beta = 0$$

where  $t_\alpha, s_\beta \geq 0$ .

Define

$$V := \sum_{\alpha \in \Delta_2^+(\gamma)} t_\alpha \alpha - \sum_{\beta \in \Delta_1^+(\gamma)} s_\beta \beta.$$

So,

$$0 \leq (V, V) = \left( \sum_{\alpha \in \Delta_2^+(\gamma)} t_\alpha \alpha, \sum_{\beta \in \Delta_1^+(\gamma)} s_\beta \beta \right) = \sum t_\alpha s_\beta (\alpha, \beta) \leq 0$$

where the last inequality is due to  $(\alpha, \beta) \leq 0$ <sup>18</sup> and  $t_\alpha, s_\beta \geq 0$ .

So,  $V = 0$ , and so  $(\gamma, V) = 0$ , in contradiction.

We've shown that every root in  $\Phi^+(\gamma)$  is a non-negative linear combination of irreducible roots in  $\Phi^+(\gamma)$ .<sup>19</sup>

This set  $\Delta^+(\gamma)$  is also linearly independent and so  $\Delta^+(\gamma)$  spans  $\Phi = \Phi^+(\gamma) \amalg -\Phi^+(\gamma)$ , and is linearly independent, and hence is a basis of  $E$ . The above shows this is also a basis of the root system. ■

<sup>18</sup>from part II.

<sup>19</sup>in part I.

**Definition 7.5.10.** A root system is called *irreducible* if it cannot be separated to two disjoint non-empty subsets which are mutually orthogonal.

**Definition 7.5.11.** If  $\Phi$  is an irreducible root system, and  $(\alpha_i)_{i \in [\ell]}$  is a basis of the root system, then

$$C = \left( \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} \right)_{i,j \in [\ell]}$$

is called *the Cartan matrix*.

## Chapter 8

# The classification of simple Lie algebras over an algebraically closed field of characteristic 0

Let  $L$  be a simple Lie algebra and  $H$  be a Cartan sub-algebra, which is a maximal toral sub-algebra. Let  $\Phi = \Phi(L, H)$  be the corresponding root system.

Remind ourself a few facts and constructions.

The root system determines the Lie algebra up to isomorphism.<sup>1</sup>

We defined a basis of the root system  $\Phi \subseteq E^2$  as a linear basis of  $E$  which has the additional property  $\Phi = \Phi^+ \amalg \Phi^-$  and that elements of  $\Phi^+$  are represented in the basis by non-negative coefficients, and elements in  $\Phi^-$  by non-negative ones.

We saw that a root system has a basis, say  $(\alpha_i)_{i \in [\ell]}$ , and we defined the Cartan matrix associated to it by

$$C = \left( \frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} \right)_{i,j \in [\ell]}.$$

Three fundamental properties of  $C$  are the following.

a)

$$\det C \neq 0$$

since  $(\alpha_i)_{i \in [\ell]}$  is a basis.

b) The off-diagonal entries of  $C$  are non-positive, since the angles between vectors are at least  $\frac{\pi}{2}$ .

c) The entries of the Cartan matrix are integers.

**Fact 8.0.1.** *The Cartan matrix determines the root system.*

## 8.1 Coxeter graphs and Dynkin diagrams

**Definition 8.1.1.** A root system is called **irreducible** if it cannot be represented as a disjoint union of two mutually-orthogonal subsets.

**Exercise.** A decomposition of the root system into a disjoint union of mutually-orthogonal subsets implies that the Lie algebra is not simple.

We assume henceforward that  $\Phi$  is irreducible.

**Definition 8.1.2.** The **Coxeter graph associated to a Cartan matrix** is a graph with  $\ell$  vertices, and for which the pairs  $\alpha_i, \alpha_j$  with  $i \neq j$ , is connected by  $\frac{4(\alpha_i, \alpha_j)(\alpha_j, \alpha_i)}{(\alpha_i, \alpha_i)(\alpha_j, \alpha_j)}$  edges.

**Remark 8.1.3.** The number of edges between  $\alpha_i, \alpha_j$  in the Coxeter graph is 0, 1, 2 or 3, as we've seen in a table 7.1. That is zero when  $\alpha_i \perp \alpha_j$ .

---

<sup>1</sup> $\Phi$  contains a full multiplication table for the Lie brackets.

<sup>2</sup>with  $\dim_{\mathbb{R}} E = \ell$

**Remark 8.1.4.** If all roots have the same length, then the Coxeter graph determines the Cartan matrix. If  $\alpha_i$  and  $\alpha_j$  have different length, then the number of edges between them cannot be 1. Otherwise,  $\frac{4(\alpha_i, \alpha_j)(\alpha_j, \alpha_i)}{(\alpha_i, \alpha_i)(\alpha_j, \alpha_j)} = 1$  and then

$$\frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)} = \frac{2(\alpha_j, \alpha_i)}{(\alpha_j, \alpha_j)}$$

because 1 is a product of two integers only in one way. We conclude that  $(\alpha_i, \alpha_i) = (\alpha_j, \alpha_j)$ , a contradiction.


**Definition 8.1.5.** The *Dynkin diagram* is the Coxeter graph where if  $\alpha_i$  is joint with  $\alpha_j$  and they have different length, then we add an arrow from the long root to the short one.

**Remark 8.1.6.** The Coxeter graph may have more than one connected component. The roots in the two components are then mutually orthogonal, and then  $\Phi$  is reducible, a contradiction to our assumption in this chapter.

We therefore reduce to classifying connected Coxeter graphs.

Set  $n = \ell$ , we list the connected Dynkin diagrams in figure 8.1. The first four diagrams are families which appear with  $n \geq 1, 2, 3, 4$  respectively. The corresponding Lie algebras for these are  $\mathfrak{sl}(\ell + 1, \mathbb{F})$ ,  $\mathfrak{so}(2\ell + 1, \mathbb{F})$ ,  $\mathfrak{sp}(2\ell, \mathbb{F})$  and  $\mathfrak{so}(2\ell, \mathbb{F})$  respectively.

Figure 8.1: Connected Dynkin diagrams.



## 8.2 Classification of connected Coxeter graphs

1. Consider in an inner-product space a set  $\{\varepsilon_i\}_{i \in [n]}$  of linearly independent unit vectors, and form the graph where a pair  $\varepsilon_i \neq \varepsilon_j$  are connected by  $4(\varepsilon_i, \varepsilon_j)^2$  edges such that this is an integer in  $\{0, 1, 2, 3\}$ , satisfying  $(\varepsilon_i, \varepsilon_j) \leq 0$ . Call such a system of vectors *admissible*.

**Remark 8.2.1.** A basis of an irreducible root system gives us an admissible set of vectors.



**Remark 8.2.2.** Every subset of an admissible system is admissible.

Graphically, given an admissible system  $U$ , if  $U' \subseteq U$ , its graph is the induced sub-graph with vertices in  $U'$ .

2. Given an admissible system  $U = \{\varepsilon_i\}_{i \in [n]}$ , the number of pairs connected by edges is at most  $n - 1$ .

*Proof.* Let  $\varepsilon = \sum_{i \in [n]} \varepsilon_i$ , then  $\varepsilon \neq 0$  because  $U$  is linearly independent. Compute

$$\begin{aligned} (\varepsilon, \varepsilon) &= \left( \sum_{i \in [n]} \varepsilon_i, \sum_{i \in [n]} \varepsilon_i \right) \\ &= n + 2 \sum_{i < j} (\varepsilon_i, \varepsilon_j) = \star. \end{aligned}$$

We have  $4(\varepsilon_i, \varepsilon_j)^2 \in \{0, 1, 2, 3\}$  and so if  $(\varepsilon_i, \varepsilon_j) \neq 0$  then  $2(\varepsilon_i, \varepsilon_j) \leq -1$ . Substituting in  $\star$ ,

$$\begin{aligned} 0 < \delta = (\varepsilon, \varepsilon) \\ &= n + \sum_{i < j} 2(\varepsilon_i, \varepsilon_j) \\ &\leq n - \#\{i < j \mid \varepsilon_i \text{ and } \varepsilon_j \text{ are connected by an edge}\} \end{aligned}$$

Therefore

$$\#\{i < j \mid \varepsilon_i \text{ and } \varepsilon_j \text{ are connected by an edge}\} \leq n - \delta$$

and so

$$\#\{i < j \mid \varepsilon_i \text{ and } \varepsilon_j \text{ are connected by an edge}\} \leq n - 1,$$

as required. ■

3. There are no cycles involving at least 3 vertices in the graph, because a cycle has  $n$  vertices and  $n$  edges, and looking at the induced sub-graph gives a contradiction to the previous part.
4. The number of edges incident to a given vertex is at most 3.

*Proof.* Let  $\varepsilon \in U$  where  $U$  is an admissible system. Let  $\{u_i\}_{i \in [k]}$  be the distinct vertices to which  $\varepsilon$  is connected, in 1, 2, or 3 edges.

These neighbours of  $\varepsilon$  cannot be connected, because that would form a (non-degenerate) cycle, contradicting the previous part. So,  $(u_i)_{i \in [k]}$  is an orthonormal set.

Let  $u_0$  be a unit vector orthogonal to  $(u_i)_{i \in [k]}$  in the space  $V = \text{span}\{\varepsilon, u_1, \dots, u_k\}$ .<sup>3</sup>

Now,  $\varepsilon \notin \text{span}\{u_i\}_{i \in [k]}$ , due to independence, and so  $\varepsilon$  cannot be orthogonal to a unit vector orthogonal to  $\text{span}\{u_i\}_{i \in [k]}$ . I.e.  $(\varepsilon, u_0) \neq 0$ . So, we represent  $\varepsilon$  in the orthonormal basis

$$\varepsilon = \sum_{i=0}^k (\varepsilon, u_i) u_i.$$

Now

$$\begin{aligned} 1 &= (\varepsilon, \varepsilon) \\ &= \sum_{i=0}^k (\varepsilon, u_i)^2 \\ &\stackrel{(\varepsilon, u_0) \neq 0}{=} \sum_{i \in [k]} (\varepsilon, u_i)^2 \end{aligned}$$

and multiplying sides by 4 we get

$$4 > \sum_{i \in [k]} 4(\varepsilon, u_i)^2 = \#\left\{\text{edges that connect } \varepsilon \text{ to } \{u_i\}_{i \in [k]}\right\}.$$
■

---

<sup>3</sup>Notice that the spanning set is a basis, because elements of  $U$  are linearly independent.

5. By the above, the only connected graph with a triple edge is  $G_2$ .

6. Suppose that the graph contains a simple chain as a sub-graph.

Let  $\varepsilon = \sum_{i \in [k]} \varepsilon_i$  and consider the system  $U' = U \setminus (\{\varepsilon_i\}_{i \in [k]} \cup \{\varepsilon\})$ . This is an admissible system, and  $\varepsilon$  has the induced inner product with any vertex that was connected to each  $\varepsilon_i$ .

*Proof.* First,  $\varepsilon$  is a unit vector.

$$\begin{aligned} (\varepsilon, \varepsilon) &= \left( \sum_{i \in [k]} \varepsilon_i, \sum_{i \in [k]} \varepsilon_i \right) \\ &= k + \sum_{i < j} 2(\varepsilon_i, \varepsilon_j) \\ &\stackrel{\dagger}{=} k - (k-1) \\ &= 1 \end{aligned}$$


with  $\dagger$  being true because we assume  $(\varepsilon_i)_{i \in [k]}$  is a chain which means only  $(\varepsilon_j, \varepsilon_{j+1}) \neq 0$  with  $4(\varepsilon_j, \varepsilon_{j+1})^2 = 1$ <sup>4</sup> and  $2(\varepsilon_j, \varepsilon_{j+1}) = -1$ <sup>5</sup>.

Clearly,  $\varepsilon$  is linearly independent of  $U \setminus \{\varepsilon_i\}_{i \in [k]}$ .

To show that  $U'$  is an admissible system, we have to show that  $4(\varepsilon, u)^2 \in \{0, 1, 2, 3\}$  for all  $u \in U \setminus \{\varepsilon_i\}_{i \in [k]}$ . Consider  $(\varepsilon, u) = \sum_{i \in [k]} (\varepsilon_i, u)$ , we claim that at least  $k-1$  of these inner products are zero. This is true because otherwise  $u$  is connected to at least two elements of the chain, and one gets a (non-degenerate) cycle, contradicting a previous result. So, if  $u \perp \varepsilon_i$  for all  $i \in [k]$ , clearly  $u \perp \varepsilon$ . If  $\exists i \in [k]: (u, \varepsilon_i) \neq 0$ , then  $i$  is determined uniquely and  $(\varepsilon, u) = (\varepsilon_i, u)$ . ■


7. A connected graph cannot contain any of the sub-graphs in figure 8.2.

Figure 8.2: Impossible graphs containing a chain.



Otherwise, using the operation in the previous part we'd get the graphs in figure 8.3.

Figure 8.3: Impossible graphs.




<sup>4</sup>as there's one edge between  $\varepsilon_j, \varepsilon_{j+1}$

<sup>5</sup>since this expression is negative

respectively, each with a vertex connected to more than 3 other vertices.

8. All connected graph are of the following forms.

Figure 8.4: Possible graphs.



$G_2$  is the only connected with with a triple edge. See fig. 8.4.

*Proof.* I) A double edge can appear only once. Otherwise we get the forbidden graph (i).

II) A tripod cannot appear together with a double edge, since this will give forbidden graph (ii).

III) A tripod can appear only once, since otherwise we get forbidden graph (iii). ■

9. Consider the graph  $\tilde{B}(p, q)$ , in fig. 8.4.

Consider  $\varepsilon = \sum_{i \in [p]} i \varepsilon_i$  and  $\zeta = \sum_{j \in [q]} j \zeta_j$ . Then

$$(\varepsilon, \varepsilon) = \sum_{i \in [p]} i^2 - \sum_{i \in [p-1]} i(i+1)$$

since  $2(\varepsilon_i, \varepsilon_{i+1}) = -1$  and  $(\varepsilon_i, \varepsilon_j) = 0$  for  $|i - j| \geq 2$ .

This yields  $(\varepsilon, \varepsilon) = \frac{p(p+1)}{2}$ , and similarly  $(\zeta, \zeta) = \frac{q(q+1)}{2}$ .

In the two chains, only  $\varepsilon_p$  and  $\zeta_q$  are connected, so

$$\begin{aligned} (\varepsilon, \zeta) &= \left( \sum_{i \in [p]} i \varepsilon_i, \sum_{j \in [q]} j \zeta_j \right) \\ &= (p \varepsilon_p, q \zeta_q) \\ &= pq (\varepsilon_p, \zeta_q) \end{aligned}$$

and so

$$(\varepsilon, \zeta)^2 = \frac{p^2 q^2}{4} 4 (\varepsilon_p, \zeta_q)^2 = p^2 q^2$$

since  $\varepsilon_p, \zeta_q$  are connected by 2 edges.

$\varepsilon$  and  $\zeta$  are linearly independent, and by Cauchy-Schwartz

$$(\varepsilon, \zeta)^2 < (\varepsilon, \varepsilon) (\zeta, \zeta).$$

So,

$$\frac{p^2 q^2}{2} < \frac{p(p+1)}{2} \frac{q(q+1)}{2}.$$

Hence

$$pq < \frac{(p+1)(q+1)}{2} = \frac{1}{2}(pq + p + q + 1).$$

Then

$$\frac{1}{2}pq - \frac{1}{2}p - \frac{1}{2}q < \frac{1}{2}$$

and then

$$(p-1)(q-1) < 2.$$

Either  $p = q = 2$ , in which case we get  $F_4$ , or  $q = 1$  and  $p$  is arbitrary, in which case we get  $B_\ell$ .

10. We show that in case (iv) we have *only*  $D_\ell$ ,  $E_6$ ,  $E_7$ , or  $E_8$ .<sup>6</sup>

*Proof.* Define the following vectors.

$$\begin{aligned}\varepsilon &= \sum_{i \in [p-1]} i \varepsilon_i \\ \eta &= \sum_{j \in [q-1]} j \eta_j \\ \zeta &= \sum_{k \in [r-1]} k \zeta_k\end{aligned}$$

$\{\varepsilon, \eta, \zeta, \psi\}$  are linearly independent.

Let  $\tau$  be a unit vector orthogonal to  $\{\varepsilon, \eta, \zeta\}$  in the space  $\text{span}\{\varepsilon, \eta, \zeta, \psi\}$ . So,  $(\psi, \tau) \neq 0$ .

$\{\varepsilon, \eta, \zeta, \tau\}$  is an orthogonal basis of this space, because clearly  $\{\varepsilon, \eta, \zeta\}$  is orthogonal.

Write

$$\psi = \frac{(\psi, \varepsilon)}{(\varepsilon, \varepsilon)^{\frac{1}{2}}} \varepsilon + \frac{(\psi, \eta)}{(\eta, \eta)^{\frac{1}{2}}} \eta + \frac{(\psi, \zeta)}{(\zeta, \zeta)^{\frac{1}{2}}} \zeta + (\psi, \tau) \tau.$$

Now  $(\psi, \psi) = 1$  because  $\psi$  is in the system.

So,

$$\frac{(\psi, \varepsilon)}{(\varepsilon, \varepsilon)^{\frac{1}{2}}} \varepsilon + \frac{(\psi, \eta)}{(\eta, \eta)^{\frac{1}{2}}} \eta + \frac{(\psi, \zeta)}{(\zeta, \zeta)^{\frac{1}{2}}} \zeta < 1.$$

As before,

$$\begin{aligned}(\varepsilon, \varepsilon) &= \frac{p(p-1)}{2} \\ (\eta, \eta) &= \frac{q(q-1)}{2} \\ (\zeta, \zeta) &= \frac{r(r-1)}{2}.\end{aligned}$$

We now compute.

$$\begin{aligned}4(\psi, \varepsilon) &= 4 \left( \psi, \sum_{i \in [p-1]} i \varepsilon_i \right)^2 \\ &= 4(\psi, (p-1) \varepsilon_{p-1})^2 \\ &= (p-1)^2 4(\psi, \varepsilon_{p-1})^2 \\ &= (p-1)^2\end{aligned}$$

so,

$$\frac{(\psi, \varepsilon)^2}{(\varepsilon, \varepsilon)} = \frac{\frac{(p-1)^2}{4}}{\frac{p(p-1)}{2}} = \frac{1}{2} \frac{p-1}{p} = \frac{1}{2} \left( 1 - \frac{1}{p} \right).$$

---

<sup>6</sup>We assume  $r, q, p \geq 2$ , because otherwise there is no tripod.

Similarly for the other two, so we conclude that

$$\frac{1}{2} \left(1 - \frac{1}{p}\right) + \frac{1}{2} \left(1 - \frac{1}{q}\right) + \frac{1}{2} \left(1 - \frac{1}{r}\right) < 1.$$

Multiplying by 2,

$$3 - \left(\frac{1}{p} + \frac{1}{q} + \frac{1}{r}\right) < 2$$

and so

$$\frac{1}{p} + \frac{1}{q} + \frac{1}{r} > 1.$$

**Exercise.** The solutions to the above equation are

$$D_n = (p, 2, 2)$$

$$E_6 = (3, 3, 2)$$

$$E_7 = (4, 3, 2)$$

$$E_8 = (5, 3, 2).$$