All the rules we know

13 October 2023

Chapter 1.A

Definition 1.1 (Complex Numbers)**.**
A **complex number** is an ordered pair (a, b), where $a, b \in \mathbb{R}$, but we will write this as $a + bi$.

The set of all complex numbers is denoted by $\mathbb{C}$:

$$\mathbb{C} = \{a + bi \mid a, b \in \mathbb{R}\}.$$

Addition and multiplication on $\mathbb{C}$ are defined by

$$\begin{aligned}(a + bi) + (c + di) &= (a + c) + (b + d)i,\\ (a + bi)(c + di) &= (ac - bd) + (ad + bc)i;\end{aligned}$$

here $a, b, c, d \in \mathbb{R}$.

Property 1.3 (Properties of real arithmetic)**.**
Intentionally restricted to $\mathbb{R}$ for the sake of the 1.A exercises. Equivalent properties can be derived for $\mathbb{C}$.

commutativity

$\alpha + \beta = \beta + \alpha$ and $\alpha\beta = \beta\alpha$ for all $\alpha, \beta \in \mathbb{R}$;

associativity

$(\alpha + \beta) + \lambda = \alpha + (\beta + \lambda)$ and $(\alpha\beta)\lambda = \alpha(\beta\lambda)$ for all $\alpha, \beta, \lambda \in \mathbb{R}$;

identities

$\lambda + 0 = \lambda$ and $\lambda 1 = \lambda$ for all $\lambda \in \mathbb{R}$;

additive inverse

for every $\alpha \in \mathbb{R}$, there exists a unique $\beta \in \mathbb{R}$ such that $\alpha + \beta = 0$;

multiplicative inverse

for every $\alpha \in \mathbb{R}$ with $\alpha \neq 0$, there exists a unique $\beta \in \mathbb{R}$ such that $\alpha\beta = 1$;

distributive property

$\lambda(\alpha + \beta) = \lambda\alpha + \lambda\beta$ for all $\lambda, \alpha, \beta \in \mathbb{R}$.

Definition 1.5 ($-\alpha$, subtraction, $1/\alpha$, division)**.**
Let $\alpha, \beta \in \mathbb{C}$.

Let $-\alpha$ denote the additive inverse of α. Thus $-\alpha$ is the unique complex number such that

$$\alpha + (-\alpha) = 0.$$

Subtraction on $\mathbb{C}$ is defined by

$$\beta - \alpha = \beta + (-\alpha).$$

For $\alpha \neq 0$, let $1/\alpha$ denote the multiplicative inverse of α. Thus $1/\alpha$ is the unique complex number such that

$$\alpha(1/\alpha) = 1.$$

Division on $\mathbb{C}$ is defined by

$$\beta/\alpha = \beta(1/\alpha).$$

Notation 1.6 ($\mathbb{F}$)**.**
$\mathbb{F}$ stands for either $\mathbb{R}$ or $\mathbb{C}$.

Definition 1.8 (list, length)**.**
Suppose n is a nonnegative integer. A **list** of **length** n is an ordered collection of n elements separated by commas and surrounded by parentheses. A list of length n looks like this:

$$(x_1, \ldots, x_n).$$

Two lists are equal if and only if they have the same length and the same elements in the same order.

Notation 1.10 (notation: n)**.**
n represents a positive integer.

Definition 1.11 ($\mathbb{F}^n$, coordinate)**.**
$\mathbb{F}^n$ is the set of all lists of length n of elements of F:

$$\mathbb{F}^n = \{(x_1, \ldots, x_n) \mid x_j \in \mathbb{F} \text{ for } j = 1, \ldots, n\}.$$

For $(x_1, \ldots, x_n) \in \mathbb{F}^n$ and $j \in \{1, \ldots, n\}$, we say that x_j is the jth **coordinate** of $(x_1, \ldots, x_n)$.

Definition 1.13 (addition in $\mathbb{F}^n$)**.**
Addition in $\mathbb{F}^n$ is defined by adding corresponding coordinates:

$$(x_1, \ldots, x_n) + (y_1, \ldots, y_n) = (x_1 + y_1, \ldots, x_n + y_n).$$

Definition 1.15 (0)**.**
Let 0 denote the list of length n whose coordinates are all 0:

$$(0, \ldots, 0).$$

Definition 1.17 (additive inverse in $\mathbb{F}^n$)**.**
For $x \in \mathbb{F}^n$, the **additive inverse** of x, denoted $-x$, is the vector $-x \in \mathbb{F}^n$ such that

$$x + (-x) = 0.$$

In other words, if $x = (x_1, \ldots, x_n)$, then $-x = (-x_1, \ldots, -x_n)$.

Definition 1.18 (scalar multiplication in $\mathbb{F}^n$)**.**
The **product** of a number λ and a vector in $\mathbb{F}^n$ is computed by multiplying each coordinate of the vector by λ:

$$\lambda(x_1, \ldots, x_n) = (\lambda x_1, \ldots, \lambda x_n);$$

Here $\lambda \in \mathbb{F}$ and $(x_1, \ldots, x_n) \in \mathbb{F}^n$.

Chapter 1.B

Definition 1.19 (addition, scalar multiplication).
An **addition** on a set V is a function that assigns an element $u + v \in V$ to each pair of elements $u, v \in V$.

A **scalar multiplication** on a set V is a function that assigns an element $\lambda v \in V$ to each $\lambda \in \mathbb{F}$ and each $v \in V$.

Definition 1.20 (vector space).
A **vector space over** $\mathbb{F}$ is a set V along with an addition on V and a scalar multiplication on V such that the following properties hold:

commutativity

$u + v = v + u$ for all $u, v \in V$;

associativity

$(u + v) + w = u + (v + w)$ and $(ab)v = a(bv)$ for all $u, v, w \in V$ and all $a, b \in \mathbb{F}$;

additive identity

there exists an element $0 \in V$ such that $v + 0 = v$ for all $v \in V$;

additive inverse

for every $v \in V$, there exists $w \in V$ such that $v + w = 0$;

multiplicative identity

$1v = v$ for all $v \in V$;

distributive property

$a(u+v) = au + av$ and $(a+b)v = av + bv$ for all $a, b \in \mathbb{F}$ and all $u, v \in V$.

Definition 1.21 (vector, point).
Elements of a vector space are called **vectors** or **points**.

Definition 1.22 (real vector space, complex vector space).
A vector space over $\mathbb{R}$ is called a **real vector space**.

A vector space over $\mathbb{C}$ is called a **complex vector space**.

Notation 1.24 ($\mathbb{F}^S$).
If S is a set, $\mathbb{F}^S$ denotes the set of functions from S to $\mathbb{F}$.

For $f, g \in \mathbb{F}^S$ the **sum** $f + g \in \mathbb{F}^S$ is the function defined by

$$(f + g)(x) = f(x) + g(x)$$

for all $x \in S$.

For $\lambda \in \mathbb{F}$ and $f \in \mathbb{F}^S$, the **product** $\lambda f \in \mathbb{F}^S$ is the function defined by

$$(\lambda f)(x) = \lambda f(x)$$

for all $x \in S$.

Notation 1.28 ($-v, w - v$).
Let $v, w \in V$. Then

1. $-v$ denotes the additive inverse of v;
2. $w - v$ is defined to be $w + (-v)$.

Notation 1.29 (V).
V denotes a vector space over $\mathbb{F}$.

The following rules can all derived from the definition of a vector space.

Result 1.26 (unique additive identity).
A vector space has a unique additive identity.

Result 1.27 (unique additive inverse).
Every element in a vector space has a unique additive inverse.

Result 1.30 (the number 0 times a vector).
$0v = 0$ for every $v \in V$.

Result 1.31 (a number times the vector 0).
$a0 = 0$ for every $a \in \mathbb{F}$.

Result 1.32 (the number -1 times a vector).
$(-1)v = -v$ for every $v \in V$.

Chapter 1.C

Definition 1.33 (subspace)**.**
A subset U of V is called a **subspace** of V if U is also a vector space with the same additive identity, addition, and scalar multiplication as on V.

Definition 1.36 (sum of subspaces)**.**
Suppose $V_1, \ldots, V_m$ are subspaces of V. The **sum** of $V_1, \ldots, V_m$, denoted by $V_1 + \cdots + V_m$, is the set of all possible sums of elements of $V_1, \ldots, V_m$. More precisely:

$$V_1 + \cdots + V_m = \{v_1 + \cdots + v_m \mid v_1 \in V_1, \ldots, v_m \in V_m\}.$$

Definition 1.41 (direct sum, $\oplus$)**.**
Suppose $V_1, \ldots, V_m$ are subspaces of V.

1. The sum $V_1 + \cdots + V_m$ is called a **direct sum** if each element of $V_1 + \cdots + V_m$ can be written in only one way as a sum $v_1 + \cdots + v_m$, where each $v_k \in V_k$.
2. If $V_1 + \cdots + V_m$ is a direct sum, then $V_1 \oplus \cdots \oplus V_m$ denotes $V_1 + \cdots + V_m$, with the $\oplus$ notation serving as an indication that this is a direct sum.

The following rules can all be derived from the definitions.

Result 1.34 (conditions for a subspace)**.**
A subset U of V is a subspace of V if and only if U satisfies the following three conditions.

additive identity

$0 \in U$.

closed under addition

$u, w \in U$ implies $u + w \in U$.

closed under scalar multiplication

$a \in \mathbb{F}$ and $u \in U$ implies $au \in U$.

Result 1.40 (sum of subspaces is the smallest containing subspace)**.**
Suppose $V_1, \ldots, V_m$ are subspaces of V. Then $V_1 + \cdots + V_m$ is the smallest subspace of V containing $V_1, \ldots, V_m$.

Result 1.45 (condition for a direct sum)**.**
Suppose $V_1, \ldots, V_m$ are subspaces of V. Then $V_1 + \cdots + V_m$ is a direct sum if and only if the only way to write 0 as a sum $v_1 + \cdots + v_m$, where each $v_k \in V_k$, is by taking each v_k equal to 0.

Result 1.46 (direct sum of two subspaces)**.**
Suppose U and W are subspaces of V. Then

$$U + W \text{ is a direct sum} \iff U \cap W = \{0\}.$$

Chapter 2.A

Notation 2.1 (list of vectors).
We write lists of vectors without surrounding parentheses.

Definition 2.2 (linear combination).
A **linear combination** of a list $v_1, \ldots, v_m$ of vectors in V is a vector of the form

$$a_1 v_1 + \cdots + a_m v_m,$$

where $a_1, \ldots, a_m \in \mathbb{F}$.

Definition 2.4 (span).
The set of all linear combinations of a list of vectors $v_1, \ldots, v_m$ in V is called the **span** of $v_1, \ldots, v_m$, denoted by $\operatorname{span}(v_1, \ldots, v_m)$. In other words

$$\operatorname{span}(v_1, \ldots, v_m) = \{a_1 v_1 + \cdots + a_m v_m \mid a_1, \ldots, a_m \in \mathbb{F}\}.$$

The span of the empty list ( ) is defined to be $\{0\}$.

Definition 2.7 (spans).
If $\operatorname{span}(v_1, \ldots, v_m)$ equals V, we say that the list $v_1, \ldots, v_m$ **spans** V.

Definition 2.9 (finite-dimensional vector space).
A vector space is called **finite-dimensional** if some list of vectors in it spans the space.

Definition 2.10 (polynomial, $\mathcal{P}(\mathbb{F})$).

1. A function $p : \mathbb{F} \to \mathbb{F}$ is called a **polynomial** with coefficients in $\mathbb{F}$ if there exist $a_0, \ldots, a_m \in \mathbb{F}$ such that

$$p(z) = a_0 + a_1 z + a_2 z^2 + \cdots + a_m z^m$$

   for all $z \in \mathbb{F}$.

2. $\mathcal{P}(\mathbb{F})$ is the set of all polynomials with coefficients in $\mathbb{F}$.

Definition 2.11 (degree of a polynomial, deg p).

1. A polynomial $p \in \mathcal{P}(\mathbb{F})$ is said to have **degree** m if there exist scalars $a_0, a_1, \ldots, a_m \in \mathbb{F}$ with $a_m \neq 0$ such that for every $z \in \mathbb{F}$, we have

$$p(z) = a_0 + a_1 z + \cdots + a_m z^m.$$

2. The polynomial that is identically 0 is said to have degree $-\infty$.
3. The degree of polynomial p is denoted by deg p.

Notation 2.12 ($\mathcal{P}_m(\mathbb{F})$).
For m a nonnegative integer, $\mathcal{P}_m(\mathbb{F})$ denotes the set of all polynomials with coefficients in $\mathbb{F}$ and degree at most m.

Definition 2.13 (infinite-dimensional vector space).
A vector space is called **infinite-dimensional** if it is not finite-dimensional.

Definition 2.15 (linearly independent).

1. A list $v_1, \ldots, v_m$ of vectors in V is called **linearly independent** if the only choice of $a_1, \ldots, a_m \in \mathbb{F}$ that makes

$$a_v v_1 + \cdots + a_m v_m = 0$$

   is $a_1 = \cdots = a_m = 0$.

2. The empty list ( ) is also declared to be linearly independent.

Definition 2.17 (linearly dependent).

1. A list of vectors V is called **linearly dependent** if it is not linearly independent.
2. In other words, a list $v_1, \ldots, v_m$ of vectors in V is linearly dependent if there exist $a_1, \ldots, a_m \in \mathbb{F}$, not all 0, such that $a_1 v_1 + \cdots + a_m v_m = 0$.

The following can be derived from the definitions.

Result 2.6 (span is the smallest containing subspace).
The span of a list of vectors in V is the smallest subspace of V containing all vectors in the list.

Lemma 2.19 (linear dependence lemma).
Suppose $v_1, \ldots, v_m$ is a linearly dependent list in V. Then there exists $k \in \{1, 2, \ldots, m\}$ such that

$$v_k \in \operatorname{span}(v_1, \ldots, v_{k-1}).$$

Furthermore, if k satisfies the condition above and the kth term is removed from $v_1, \ldots, v_m$, then the span of the remaining list equals $\operatorname{span}(v_1, \ldots, v_m)$.

Result 2.22 (length of linearly independent list $\leq$ length of spanning list).
In a finite-dimensional vector space, the length of every linearly independent list of vectors is less than or equal to the length of every spanning list of vectors.

Result 2.25 (finite-dimensional subspaces).
Every subspace of a finite-dimensional vector space is finite-dimensional.

Chapter 2.B

Definition 2.26 (basis)**.**
A **basis** of V is a list of vectors in V that is linearly dependent and spans V.

Result 2.28 (criterion for basis)**.**
A list $v_1, \ldots, v_n$ of vectors of V is a basis of V if and only if every $v \in V$ can be written uniquely in the form

$$v = a_1 v_1 + \cdots + a_n v_n,$$

where $a_1, \ldots, a_n \in \mathbb{F}$.

Result 2.30 (every spanning list contains a basis)**.**
Every spanning list in a vector space can be reduced to a basis of the vector space.

Corollary 2.31 (basis of finite-dimensional vector space)**.**
Every finite-dimensional vector space has a basis.

Result 2.32 (every linearly independent list extends to a basis)**.**
Every linearly independent list of vectors in a finite-dimensional vector space can be extended to a basis of the vector space.

Corollary 2.33 (every subspace of V is part of a direct sum equal to V)**.**
Suppose V is finite-dimensional and U is a subspace of V. Then there is a subspace W of V such that $V = U \oplus W$.

Chapter 2.C

Definition 2.35 (dimension, $\dim V$)**.**
The **dimension** of a finite-dimensional vector space is the length of any basis of the vector space.

The dimension of a finite-dimensional vector space V is denoted by $\dim V$.

Result 2.34 (basis length does not depend on basis)**.**
Any two bases of a finite-dimensional vector space have the same length.

Result 2.37 (dimension of a subspace)**.**
If V is finite-dimensional and U is a subspace of V, then $\dim U \leq \dim V$.

Result 2.38 (linearly independent list of the right length is a basis)**.**
Suppose V is finite-dimensional. Then every linearly independent list of vectors in V of length $\dim V$ is a basis of V.

Corollary 2.39 (subspace of full dimension equals the whole space)**.**
Suppose that V is finite-dimensional and U is a subspace of V such that $\dim U = \dim V$. Then $U = V$.

Result 2.42 (spanning list of the right length is a basis)**.**
Suppose V is finite-dimensional. Then every spanning list of vectors in V of length $\dim V$ is a basis of V.

Result 2.43 (dimension of sum)**.**
If V_1 and V_2 are subspaces of a finite-dimensional vector space, then

$$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2).$$

Result Page 48 (Sets and vector spaces)**.**

sets	**vector spaces**
S is a finite set	V is a finite-dimensional vector space
$\#S$	$\dim V$
for subsets S_1, S_2 of S, the union $S_1 \cup S_2$ is the smallest subset of S containing S_1 and S_2	for subspaces V_1, V_2 of V, the sum $V_1 + V_2$ is the smallest subspace of V containing V_1 and V_2
$\#(S_1 \cup S_2) = \#S_1 + \#S_2 - \#(S_1 \cap S_2)$	$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 - \dim(V_1 \cap V_2)$
$\#(S_1 \cup S_2) = \#S_1 + \#S_2 \iff \#(S_1 \cap S_2) =$	$\dim(V_1 + V_2) = \dim V_1 + \dim V_2 \iff \dim(V_1 \cap V_2) =$
$S_1 \cup \cdots \cup S_m$ is a disjoint union $\iff \#(S_1 \cup \cdots \cup S_M) = \#S_1 + \cdots + \#S_M$	$V_1 \cdots + V_m$ is a direct sum $\iff \dim(V_1 + \cdots + V_m) = \dim V_1 + \cdots + \dim V_m$

Chapter 3.A

Definition 3.1 (linear map)**.**
A **linear map** from V to W is a function $T : V \to W$ with the following properties.

additivity

$T(u + v) = Tu + Tv$ for all $u, v \in V$.

homogeneity

$T(\lambda v) = \lambda(Tv)$ for all $\lambda \in \mathbb{F}$ and all $v \in V$.

Notation 3.2 ($\mathcal{L}(V, W), \mathcal{L}(V)$)**.**

1. The set of linear maps from V to W is denoted by $\mathcal{L}(V, W)$.
2. The set of linear maps from V to V is denoted by $\mathcal{L}(V)$. In other words, $\mathcal{L}(V) = \mathcal{L}(V, V)$.

Definition 3.5 (addition and scalar multiplication on $\mathcal{L}(V, W)$)**.**
Suppose $S, T \in \mathcal{L}(V, w)$ and $\lambda \in \mathbb{F}$. The **sum** $S+T$ and the **product** λT are the linear maps from V to W defined by

$$(S + T)(v) = Sv + Tv \quad \text{and} \quad (\lambda T)(v) = \lambda(Tv)$$

for all $v \in V$.

Definition 3.7 (product of linear maps)**.**
If $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$, then the **product** $ST \in \mathcal{L}(U, V)$ is defined by

$$(ST)(u) = S(Tu)$$

for all $u \in U$.

Lemma 3.4 (linear map lemma)**.**
Suppose $v_1, \ldots, v_n$ is a basis of V and $w_1, \ldots, w_m \in W$. Then there exists a unique linear map $T : V \to W$ such that

$$Tv_k = w_k$$

for each $k = 1, \ldots, n$.

Result 3.6 ($\mathcal{L}(V, W)$ is a vector space)**.**
With the operations of addition and scalar multiplication as in Definition 3.5, $\mathcal{L}(V, W)$ is a vector space.

Result 3.8 (algebraic properties of products of linear maps)**.**

associativity

$(T_1T_2)T_3 = T_1(T_2T_3)$ whenever T_1, T_2 and T_3 are linear maps such that the products make sense (meaning T_3 maps into the domain of T_2 and T_2 maps into the domain of T_1).

identity

$TI = IT = T$ whenever $T \in \mathcal{L}(V, W)$; here the first I is the identity operator on V and the second I is the identity operator on W.

distributive properties

$(S_1 + S_2)T = S_1T + S_2T$ and $S(T_1 + T_2) = ST_1 + ST_2$ whenever $T, T_1, T_2 \in \mathcal{L}(U, V)$ and $S, S_1, S_2 \in \mathcal{L}(V, W)$.

Result 3.10 (linear maps take 0 to 0)**.**
Suppose T is a linear map from V to W. Then $T(0) = 0$.

Result Ex. 3A, 13 (Linear maps on a subspace can be extended to a map on the whole vector space)**.**
Suppose V is finite-dimensional. Prove that every linear map on a subspace of V can be extended to a linear map on V. In other words, show that if U is a subspace of V and $S \in \mathcal{L}(U, V)$, then there exists $T \in \mathcal{L}(V, W)$ such that $Tu = Su$ for all $u \in E$.

Chapter 3.B

Definition 3.11 (null space, $\operatorname{null} T$)**.**
For $T \in \mathcal{L}(V, W)$, the **null space** of T, denoted by null T, is the subset of V consisting of those vectors that T maps to 0:

$$\operatorname{null} T = \{v \in V \mid Tv = 0\}.$$

Definition 3.14 (injective)**.**
A function $T : V \to W$ is called **injective** if $Tu = Tv$ implies $u = v$.

Definition 3.16 (range)**.**
For $T \in \mathcal{L}(V, W)$, the **range** of T is the subset W consisting of those vectors that are equal to Tv for some $v \in V$:

$$\operatorname{range} T = \{Tv \mid v \in V\}.$$

Definition 3.19 (surjective)**.**
A function $T : V \to W$ is called **surjective** if its range equals W.

Result 3.13 (the null space is a subspace)**.**
Suppose $T \in \mathcal{L}(V, W)$. Then $\operatorname{null} T$ is a subspace of V.

Result 3.15 (injectivity $\Leftrightarrow$ null space equals $\{0\}$)**.**
Let $T \in \mathcal{L}(V, W)$. Then T is injective if and only if $\operatorname{null} T = \{0\}$.

Result 3.18 (the range is a subspace)**.**
If $T \in \mathcal{L}(V, W)$, then the range T is a subspace of W.

Theorem 3.21 (fundamental theorem of linear maps)**.**
Suppose V is finite-dimensional and $T \in \mathcal{L}(V, W)$. Then $\operatorname{range} T$ is finite-dimensional and

$$\dim V = \dim \operatorname{null} T + \dim \operatorname{range} T.$$

Result 3.22 (linear map to a lower-dimensional space is not injective)**.**
Suppose V and W are finite-dimensional vector spaces such that $\dim V > \dim W$. Then no linear map from V to W is injective.

Result 3.24 (linear map to a higher-dimensional space is not surjective)**.**
Suppose V and W are finite-dimensional vector spaces such that $\dim V < \dim W$. Then no linear map from V to W is surjective.

Result 3.26 (homogeneous system of linear equations)**.**
A homogeneous system of linear equations with more variables than equations has nonzero solutions.

Result 3.28 (inhomogeneous systems of linear equations)**.**
An inhomogeneous system of linear equations with more equations than variables has no solution for some choice of the constant terms.

Chapter 3.C

Definition 3.29 (matrix, $A_{j,k}$)**.**
Suppose m and n are nonnegative integers. An m-by-n **matrix** A is a rectangular array of elements of $\mathbb{F}$ with m rows and n columns:

$$A = \begin{pmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{pmatrix}.$$

The notation $A_{j,k}$ denotes the entry in row j, column k of A.

Definition 3.31 (matrix of a linear map, $\mathcal{M}(T)$)**.**
Suppose $T \in \mathcal{L}(V, W)$ and $v_1, \ldots, v_n$ is a basis of V and $w_1, \ldots, w_m$ is a basis of W. The **matrix of** T with respect to these bases is the m-by-n matrix $\mathcal{M}(T)$ whose entries $A_{j,k}$ are defined by

$$Tv_k = A_{1,k}w_1 + \cdots + A_{m,k}w_m.$$

If the basis $v_1, \ldots, v_n$ and $w_1, \ldots, w_m$ are not clear from the context, then the notation $\mathcal{M}(T, (v_1, \ldots, v_n), (w_1, \ldots, w_m))$ is used.

Definition 3.34 (matrix addition)**.**
The **sum of two matrices of the same size** is the matrix obtained by adding corresponding entries in the matrices:

$$\begin{pmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{pmatrix} + \begin{pmatrix} C_{1,1} & \cdots & C_{1,n} \\ \vdots & & \vdots \\ C_{m,1} & \cdots & C_{m,n} \end{pmatrix}$$
$$= \begin{pmatrix} A_{1,1} + C_{1,1} & \cdots & A_{1,n} + C_{1,n} \\ \vdots & & \vdots \\ A_{m,1} + C_{m,1} & \cdots & A_{m,n} + C_{m,n} \end{pmatrix}.$$

Definition 3.36 (scalar multiplication of a matrix)**.**
The product of a scalar and a matrix is the matrix obtained by multiplying each entry in the matrix by the scalar:

$$\lambda \begin{pmatrix} A_{1,1} & \cdots & A_{1,n} \\ \vdots & & \vdots \\ A_{m,1} & \cdots & A_{m,n} \end{pmatrix} = \begin{pmatrix} \lambda A_{1,1} & \cdots & \lambda A_{1,n} \\ \vdots & & \vdots \\ \lambda A_{m,1} & \cdots & \lambda A_{m,n} \end{pmatrix}.$$

Notation 3.39 ($\mathbb{F}^{m,n}$)**.**
For m and n positive integers, the set of all m-by-n matrices with entries in $\mathbb{F}$ is denoted by $\mathbb{F}^{m,n}$.

Definition 3.41 (matrix multiplication)**.**
Suppose A is an m-by-n matrix and B is an n-by-p matrix. Then AB is defined to be the m-by-p matrix whose entry in row j, column k, is given by the equation

$$(AB)_{j,k} = \sum_{r=1}^{n} A_{j,r}B_{r,k}.$$

Thus the entry in row j, column k, of AB is computed by taking row i of A and column k of B, multiplying together corresponding entries, and then summing.

Notation 3.44 ($A_{j,\cdot}, A_{\cdot,k}$)**.**
Suppose A is an m-by-n matrix.

1. If $1 \leq j \leq m$, then $A_{j,\cdot}$ denotes the 1-by-n matrix consisting of row j of A.
2. If $1 \leq k \leq n$, then $A_{\cdot,k}$ denotes the m-by-1 matrix consisting of column k of A.

Definition 3.52 (column rank, row rank)**.**
Suppose A is an m-by-n matrix with entries in $\mathbb{F}$.

1. The **column rank** of A is the dimension of the span of the columns of A in $\mathbb{F}^{m,1}$.
2. The **row rank** of A is the dimension of the span of the rows of A in $\mathbb{F}^{1,n}$.

Definition 3.54 (transpose, A^{t})**.**
The **transpose** of a matrix A, denoted by A^{t}, is the matrix obtained from A by interchanging rows and columns. Specifically, if A is an m-by-n matrix, then A^{t} is an n-by-m matrix whose entries are given by the equation

$$(A^{\mathrm{t}})_{k,j} = A_{j,k}.$$

Definition 3.58 (rank)**.**
The **rank** of a matrix $A \in \mathbb{F}^{m,n}$ is the column rank of A.

Result 3.35 (matrix of the sum of linear maps).
Suppose $S, T \in \mathcal{L}(V, W)$. Then $\mathcal{M}(S+T) = \mathcal{M}(S) + \mathcal{M}(T)$.

Result 3.38 (the matrix of a scalar times a linear map).
Suppose $\lambda \in \mathbb{F}$ and $T \in \mathcal{L}(V, W)$. Then $\mathcal{M}(\lambda T) = \lambda \mathcal{M}(T)$.

Result 3.40 ($\dim \mathbb{F}^{m,n} = mn$).
Suppose m and n are positive integers. With addition and scalar multiplication defined as above, $\mathbb{F}^{m,n}$ is a vector space of dimension mn.

Result 3.43 (matrix of product of linear maps).
If $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$, then $\mathcal{M}(ST) = \mathcal{M}(S)\mathcal{M}(T)$.

Result 3.46 (entry of matrix product equals row times column).
Suppose A is an m-by-n matrix and B is an n-by-p matrix. Then

$$(AB)_{j,k} = A_{j,\cdot} B_{\cdot,k}$$

if $1 \le j \le m$ and $1 \le k \le p$. In other words, the entry in row j, column k, of AB equals (row j of A) times (column k of B).

Result 3.48 (column of matrix product equals matrix times column).
Suppose A is an m-by-n matrix and B is an n-by-p matrix. Then

$$(AB)_{\cdot,k} = AB_{\cdot,k}$$

if $1 \le k \le p$. In other words, column k of AB equals A times column k of B.

Result Ex. 3C, 8 (row of matrix product equals matrix times row).
Suppose A is an m-by-n matrix and B is an n-by-p matrix. Then

$$(AB)_{j,\cdot} = A_{j,\cdot} B$$

if $1 \le j \le m$. In other words, row j of AB equals row j of A times B.

This is the row version of Result 3.48.

Result 3.50 (linear combination of columns).
Suppose A is an m-by-n matrix and $b = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}$ is an n-by-1 matrix. Then

$$Ab = b_1 A_{\cdot,1} + \cdots + b_n A_{\cdot,n}.$$

In other words, Ab is a linear combination of the columns of A, with the scalars that multiply the columns coming from b.

Result Ex. 3C, 9 (linear combination of rows).
Suppose $a = \begin{pmatrix} a_1 & \cdots & a_n \end{pmatrix}$ is a 1-by-n matrix and B is an n-by-p matrix. Then

$$aB = a_1 B_{1,\cdot} + \cdots + a_n B_{n,\cdot}.$$

In other words, aB is a linear combination of the rows of B, with the scalars that multiply the rows coming from a.

This is the row version of Result 3.50.

Result 3.51 (matrix multiplication as linear combinations of columns).
Suppose C is an m-by-c matrix and R is a c-by-n matrix.

(a) If $k \in \{1, \ldots, n\}$, then column k of CR is a linear combination of the columns of C, with the coefficients of this linear combination coming from columns k of R.

(b) If $j \in \{1, \ldots, m\}$, then row j of CR is a linear combination of the rows of R, with the coefficients of this linear combination coming from row j of C.

Result Ex. 3C, 14 (transpose is a linear map).
Suppose m and n are positive integers. Then the function $A \mapsto A^{\mathrm{t}}$ is a linear map from $\mathbb{F}^{m,n}$ to $\mathbb{F}^{n,m}$.

In other words $(A+B)^{\mathrm{t}} = A^{\mathrm{t}} + B^{\mathrm{t}}$, $(\lambda A)^{\mathrm{t}} = \lambda A^{\mathrm{t}}$ for all m-by-n matrices A, B and all $\lambda \in \mathbb{F}$.

Result Ex. 3C, 15 (The transpose of the product is the product of the transposes in the opposite order).
If A is an m-by-n matrix and C is an n-by-p matrix, then

$$(AC)^{\mathrm{t}} = C^{\mathrm{t}} A^{\mathrm{t}}.$$

Result 3.56 (column-row factorisation).
Suppose A is an m-by-n matrix with entries in $\mathbb{F}$ and column rank $c \ge 1$. Then there exist an m-by-c matrix C and a c-by-n matrix R, both with entries in $\mathbb{F}$, such that $A = CR$.

Result 3.57 (column rank equals row rank).
Suppose $A \in \mathbb{F}^{m,n}$. Then the column rank of A equals the row rank of A.

Chapter 3.D

Definition 3.59 (invertible, inverse)**.**

1. A linear map $T \in \mathcal{L}(V, W)$ is called **invertible** if there exists a linear map $S \in \mathcal{L}(W, V)$ such that ST equals the identity operator on V and TS equals the identity operator on W.
2. A linear map $S \in \mathcal{L}(W, V)$ satisfying $ST = I$ and $TS = I$ is called an **inverse** of T (note that the first I is the identity operator on V and the second I is the identity operator on W.

Notation 3.61 (T^{-1})**.**
If T is invertible, then its inverse is denoted by T^{-1}. In other words, if $T \in \mathcal{L}(V, W)$ is invertible, then T^{-1} is the unique element of $\mathcal{L}(W, V)$ such that $T^{-1}T = I$ and $TT^{-1} = I$.

Definition 3.69 (isomorphism, isomorphic)**.**

1. An **isomorphism** is an invertible linear map.
2. Two vector spaces are called **isomorphic** if there is an isomorphism from one vector space onto the other one.

Definition 3.73 (matrix of a vector, $\mathcal{M}(v)$)**.**
Suppose $v \in V$ and $v_1, \ldots, v_n$ is a basis of V. The **matrix of** V with respect to this basis is the n-by-1 matrix

$$\mathcal{M}(v) = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix},$$

where $b_1, \ldots, b_n$ are the scalars such that

$$v = b_1 v_1 + \cdots + b_n v_n.$$

Definition 3.79 (identity matrix, I)**.**
Suppose n is a positive integer. The n-by-n matrix

$$\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}$$

with 1's on the diagonal (the entries where the row number equals the column number) and 0's elsewhere is called the **identity matrix** and is denoted by I.

Definition 3.80 (invertible, inverse, A^{-1})**.**
A square matrix A is called **invertible** if there is a square matrix B of the same size such that $AB = BA = I$; we call B the **inverse** of A and denote it by A^{-1}.

Result 3.60 (inverse is unique)**.**
An invertible linear map has a unique inverse.

Result 3.63 (invertibility $\iff$ injectivity and surjectivity)**.**
A linear map is invertible if and only if it is injective and surjective.

Result 3.65 (injectivity is equivalent to surjectivity (if $\dim V = \dim W < \infty$))**.**
Suppose that V and W are finite-dimensional vector spaces, $\dim V = \dim W$, and $T \in \mathcal{L}(V, W)$. Then

$$T \text{ is invertible } \iff T \text{ is injective } \iff T \text{ is surjective.}$$

Result 3.68 ($ST = I \iff TS = I$ (on vector spaces of the same dimension))**.**
Suppose V and W are finite-dimensional vector spaces of the same dimension, $S \in \mathcal{L}(V, W)$, and $T \in \mathcal{L}(W, V)$. Then $ST = I$ if and only if $TS = I$.

Result 3.70 (dimension shows whether vector spaces are isomorphic)**.**
Two finite-dimensional vector spaces over $\mathbb{F}$ are isomorphic if and only if they have the same dimension.

Result 3.71 ($\mathcal{L}(V, W)$ and $\mathbb{F}^{m,n}$ are isomorphic)**.**
Suppose $v_1, \ldots, v_n$ is a basis of V and $w_1, \ldots, w_m$ is a basis of W. Then M is an isomorphism between $\mathcal{L}(V, W)$ and $\mathbb{F}^{m,n}$.

Result 3.72 ($\dim \mathcal{L}(V, W) = (\dim V)(\dim W)$)**.**
Suppose V and W are finite-dimensional. Then $\mathcal{L}(V, W)$ is finite-dimensional and

$$\dim \mathcal{L}(V, W) = (\dim V)(\dim W).$$

Result 3.75 ($\mathcal{M}(T)_{\cdot,k} = \mathcal{M}(Tv_k)$)**.**
Suppose $T \in \mathcal{L}(V, W)$ and $v_1, \ldots, v_n$ is a basis of V and $w_1, \ldots, w_m$ is a basis of W. Let $1 \leq k \leq n$. Then the kth column of $\mathcal{M}(T)$, which is denoted by $\mathcal{M}(T)_{\cdot,k}$, equals $\mathcal{M}(Tv_k)$.

Result 3.76 (linear maps act like matrix multiplication)**.**
Suppose $T \in \mathcal{L}(V, W)$ and $v \in V$. Suppose $v_1, \ldots, v_n$ is a basis of V and $w_1 \ldots, w_m$ is a basis of W. Then

$$\mathcal{M}(Tv) = \mathcal{M}(T)\mathcal{M}(v).$$

Result 3.78 (dimension of range T equals column rank of $\mathcal{M}(T)$)**.**
Suppose V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$. Then $\dim \operatorname{range} T$ equals the column rank of $\mathcal{M}(T)$.

Result 3.81 (matrix of product of linear maps)**.**
Suppose $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$. If $u_1, \ldots, u_m$ is a basis of U, $v_1, \ldots, v_n$ is a basis of V and $w_1, \ldots, w_p$ is a basis for W, then

$$\begin{aligned} \mathcal{M}(ST, (u_1, \ldots, u_m), (w_1, \ldots, w_p)) = \\ \mathcal{M}(S, (v_1, \ldots, v_n), (w_1, \ldots, w_p)) \\ \mathcal{M}(U, (u_1, \ldots, u_m), (v_1, \ldots, v_n)). \end{aligned}$$

Result 3.82 (matrix of identity operator with respect to two bases)**.**
Suppose that $u_1, \ldots, u_n$ and $v_1, \ldots, v_n$ are bases of V. Then the matrices $\mathcal{M}(I, (u_1, \ldots, u_n), (v_1, \ldots, v_n))$ and $\mathcal{M}(I, (v_1, \ldots, v_n), (u_1, \ldots, u_n))$ are invertible, and each is the inverse of the other.

Result 3.84 (change-of-basis formula)**.**
Suppose $T \in \mathcal{L}(V)$. Suppose $u_1, \ldots, u_n$ and $v_1, \ldots, v_n$ are bases of V. Let $A = \mathcal{M}(T, (u_1, \ldots, u_n))$ and $B = \mathcal{M}(T, (v_1, \ldots, v_n))$ and $C = \mathcal{M}(I, (u_1, \ldots, u_n), (v_1 \ldots, v_n))$. Then

$$A = C^{-1}BC.$$

Result 3.86 (matrix of inverse equals inverse of matrix)**.**
Suppose that $v_1, \ldots, v_n$ is a basis of V and $T \in \mathcal{L}(V)$ is invertible. Then $\mathcal{M}(T^{-1}) = (\mathcal{M}(T))^{-1}$, where both matrices are with respect to the basis $v_1, \ldots, v_n$.

Chapter 3.E

Definition 3.87 (product of vector spaces).
Suppose $V_1, \ldots, V_m$ are vector spaces over $\mathbb{F}$.

1. The **product** $V_1 \times \cdots \times V_m$ is defined by
$$V_1 \times \cdots \times V_m = \{(v_1, \ldots, v_m) \mid v_1 \in V_1, \ldots, v_m \in V_m\}.$$
2. Addition on $V_1 \times \cdots \times V_m$ is defined by
$$(u_1, \ldots, u_m) + (v_1, \ldots, v_m) = (u_1 + v_1, \ldots, u_m + v_m).$$
3. Scalar multiplication on $V_1 \times \cdots \times V_m$ is defined by
$$\lambda(v_1, \ldots, v_m) = (\lambda v_1, \ldots, \lambda v_m).$$

Notation 3.95 ($v + U$).
Suppose $v \in V$ and $U \subseteq V$. Then $v + U$ is the subset of V defined by
$$v + U = \{v + u \mid u \in U\}$$

Definition 3.97 (translate).
For $v \in V$ and U a subset of V, the set $v + U$ is said to be a **translate** of U.

Definition 3.99 (quotient space, V/U).
Suppose U is a subspace of V. The **quotient space** V/U is the set of all translates of U. Thus
$$V/U = \{v + U \mid v \in V\}.$$

Definition 3.102 (addition and scalar multiplication on V/U).
Suppose U is a subspace of V. The **addition** and **scalar multiplication** are defined on V/U by
$$(v + U) + (w + U) = (v + w) + U$$
$$\lambda(v + U) = (\lambda v) + U$$
for all $v, w \in V$ and all $\lambda \in \mathbb{F}$.

Definition 3.104 (quotient map, π).
Suppose U is a subspace of V. The **quotient map** $\pi : V \to V/U$ is the linear map defined by
$$\pi(v) = v + U$$
for each $v \in V$.

Notation 3.106 ($\widetilde{T}$).
Suppose $T \in \mathcal{L}(V, W)$. Define $\widetilde{T} : V/(\operatorname{null} T) \to W$ by
$$\widetilde{T}(v + \operatorname{null} T) = Tv.$$

Notation (composition of linear maps, $\circ$).
If $T \in \mathcal{L}(U, V)$ and $S \in \mathcal{L}(V, W)$, then the **composition** notation $S \circ T$ is an alternative way of writing the product ST of S and T, as given in Definition 3.7.

Result 3.89 (product of vector spaces is a vector space).
Suppose $V_1, \ldots, V_n$ are vector spaces over $\mathbb{F}$. Then $V_1 \times \cdots \times V_m$ is a vector space over $\mathbb{F}$.

Result 3.92 (dimension of a product is the sum of dimensions).
Suppose $V_1, \ldots, V_m$ are finite-dimensional vector spaces. Then $V_1 \times \cdots \times V_m$ is finite-dimensional and
$$\dim(V_1 \times \cdots \times V_m) = \dim V_1 + \cdots + \dim V_m.$$

Result 3.93 (products and direct sums).
Suppose that $V_1, \ldots, V_m$ are subspaces of V. Define a linear map $\Gamma : V_1 \times \cdots \times V_m \to V_1 + \cdots + V_m$ by
$$\Gamma(v_1, \ldots, v_m) = v_1 + \ldots + v_m.$$
Then $V_1 + \cdots + V_m$ is a direct sum if and only if Γ is injective.

Result 3.94 (a sum is a direct sum if and only if dimensions add up).
Suppose V is finite-dimensional and $V_1, \ldots, V_m$ are subspaces of V. Then $V_1 + \cdots + V_m$ is a direct sum if and only if
$$\dim(V_1 + \cdots + V_m) = \dim V_1 + \cdots + \dim V_m.$$

Result 3.101 (two translates of a subspace are equal or disjoint).
Suppose U is a subspace of V and $v, w \in V$. Then
$$v - w \in U \iff v + U = w + U \iff (v + U) \cap (w + U) \neq \emptyset.$$

Result 3.103 (quotient space is a vector space).
Suppose U is a subspace of V. Then V/U, with the operations of addition and scalar multiplication as defined in Definition 3.102, is a vector space.

Result 3.105 (dimension of quotient space).
Suppose V is finite-dimensional and U is a subspace of V. Then
$$\dim V/U = \dim U - \dim V.$$

Result 3.107 (null space and range of $\widetilde{T}$).
Suppose $T \in \mathcal{L}(V, W)$. Then

(a) $\widetilde{T} \circ \pi = T$, where π is the quotient map of V onto $V/(\operatorname{null} T)$;

(b) $\widetilde{T}$ is injective;

(c) $\operatorname{range} \widetilde{T} = \operatorname{range} T$;

(d) $V/(\operatorname{null} T)$ and $\operatorname{range} T$ are isomorphic vector spaces.

Result Ex. 3E, 18 (Direct sum of a quotient).
Suppose U is a subspace of V such that V/U is finite-dimensional. Then there exists a finite-dimensional subspace W of V such that $\dim W = \dim V/U$ and $V = U \oplus W$.

Chapter 3.F

Definition 3.108 (linear functional).
A **linear functional** on V is a linear map from V to $\mathbb{F}$. In other words, a linear functional is an element of $\mathcal{L}(V, \mathbb{F})$.

Definition 3.110 (dual space V').
The **dual space** of V, denoted by V', is the vector space of all linear functionals on V. In other words, $V' = \mathcal{L}(V, \mathbb{F})$.

Definition 3.112 (dual basis).
If $v_1, \ldots, v_n$ is a basis of V, then the **dual basis** of $v_1, \ldots, v_n$ is the list $\varphi_1, \ldots, \varphi_n$ of elements of V', where each φ_j is the linear functional on V such that

$$\varphi_j(v_k) = \begin{cases} 1 & \text{if } k = j, \\ 0 & \text{if } k \neq j. \end{cases}$$

Definition 3.118 (dual map, T').
Suppose $T \in \mathcal{L}(V, W)$. The **dual map** of T is the linear map $T' \in \mathcal{L}(W', V')$ defined for each $\varphi \in W'$ by

$$T'(\varphi) = \varphi \circ T.$$

Definition 3.121 (annihilator, U^0).
For $U \subseteq V$, the **annihilator** of U, denoted by U^0, is defined by

$$U^0 = \{\varphi \in V' \mid \varphi(u) = 0 \text{ for all } u \in U\}.$$

Result 3.111 ($\dim V' = \dim V$).
Suppose V is finite-dimensional. Then V' is also finite-dimensional and

$$\dim V' = \dim V.$$

Result 3.114 (dual basis gives coefficients for linear combination).
Suppose $v_1, \ldots, v_n$ is a basis of V and $\varphi_i, \ldots, \varphi_n$ is the dual basis. Then for each $v \in V$

$$v = \varphi_1(v)v_1 + \cdots + \varphi_n(v)v_n$$

Result 3.116 (dual basis is a basis of the dual space).
Suppose V is finite-dimensional. Then the dual basis of a basis of V is a basis of V'.

Result 3.120 (algebraic properties of dual maps).
Suppose $T \in \mathcal{L}(V, W)$. Then

(a) $(S + T)' = S' + T'$ for all $S \in \mathcal{L}(V, W)$;

(b) $(\lambda T)' = \lambda T'$ for all $\lambda \in \mathbb{F}$;

(c) $(ST)' = T'S'$ for all $S \in \mathcal{L}(W, U)$.

Result 3.124 (the annihilator is a subspace).
Suppose $U \subseteq V$. Then U^0 is a subspace of V'.

Result 3.125 (dimension of the annihilator).
If V is finite-dimensional and U is a subspace of V then

$$\dim U^0 = \dim V - \dim U.$$

Result 3.127 (condition for the annihilator to equal $\{0\}$ or the whole space).
If V is finite-dimensional and U is a subspace of V then

(a) $U^0 = \{0\} \iff U = V$;

(b) $U^0 = V' \iff U = \{0\}$.

Result 3.128 (the null space of T').
Suppose $T \in \mathcal{L}(V, W)$. Then

(a) $\operatorname{null} T' = (\operatorname{range} T)^0$.

Suppose further that V and W are finite-dimensional. Then

(b) $\dim \operatorname{null} T' = \dim \operatorname{null} T + \dim W - \dim W$.

Result 3.129 (T surjective is equivalent to T' injective).
If V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$ then

$$T \text{ is surjective } \iff T' \text{ is injective.}$$

Result 3.130 (the range of T').
If V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$ then

(a) $\dim \operatorname{range} T' = \dim \operatorname{range} T$;

(b) $\operatorname{range} T' = (\operatorname{null} T)^0$.

Result 3.131 (T injective is equivalent to T' surjective).
If V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$ then

$$T \text{ is injective } \iff T' \text{ is surjective.}$$

Result 3.132 (matrix of T' is transpose of matrix of T).
If V and W are finite-dimensional and $T \in \mathcal{L}(V, W)$ then

$$\mathcal{M}(T') = (\mathcal{M}(T))^{\mathrm{t}}.$$