

STT 3850 : Chi-Square Tests

Fall 2023

Appalachian State University

Section 1

Chi-Square Goodness-of-Fit Tests

Background

- Many statistical procedures require knowledge of the population from which the sample is taken. For example, using Student's t -distribution for testing a hypothesis or constructing a confidence interval for μ assumes that the parent population is normal.
- **Goodness-of-fit** (GOF) procedures are presented that will help to identify the distribution of the population from which the sample is drawn.
- The null hypothesis in a goodness-of-fit test is a statement about the form of the cumulative distribution. When all the parameters in the null hypothesis are specified, the hypothesis is called **simple**.
- Recall that in the event the null hypothesis does not completely specify all of the parameters of the distribution, the hypothesis is said to be **composite**.

Background

- Goodness-of-fit tests are typically used when the form of the population is in question. In contrast to most of the statistical procedures discussed so far, where the goal has been to **reject** the null hypothesis, in a GOF test one hopes to **retain** the null hypothesis.
- Given a single random sample of size n from an unknown population F_X , one may wish to test the hypothesis that F_X has some known distribution $F_0(x)$ for all x .

Background

- For example, using the data frame `SOCGER` from the `PASWR2` package, is it reasonable to assume the number of goals scored during regulation time for the 232 soccer matches has a Poisson distribution with $\lambda = 2.5$?
- Before applying the chi-square goodness-of-fit test, the data must be grouped according to some scheme to form k mutually exclusive categories. When the null hypothesis completely specifies the population, the probability that a random observation will fall into each of the chosen or fixed categories can be computed.

Background

- Once the probabilities for a data point to fall into each of the chosen or fixed categories is computed, multiplying the probabilities by n produces the expected counts for each category under the null distribution.
- If the null hypothesis is true, the differences between the counts observed in the k categories and the counts expected in the k categories should be small.

Background

- The test criterion for testing $H_0 : F_X(x) = F_0(x)$ for all x against the alternative $H_1 : F_X(x) \neq F_0(x)$ for some x when the null hypothesis is completely specified is

$$\chi_{\text{obs}}^2 = \sum_{i=1}^k \frac{(O_k - E_k)^2}{E_k}, \quad (1)$$

where χ_{obs}^2 is the sum of the squared deviations between what is observed (O_k) and what is expected (E_k) in each of the k categories divided by what is expected in each of the k categories. Large values of χ_{obs}^2 occur when the observed data are inconsistent with the null hypothesis and thus lead to rejection of the null hypothesis. The exact distribution of χ_{obs}^2 is very complicated; however, for large n , provided all expected categories are at least 5, χ_{obs}^2 is distributed approximately χ^2 with $k - 1$ degrees of freedom.

Background

- NOTE: When the null hypothesis is composite, that is, not all of the parameters are specified, the degrees of freedom for the random variable χ^2_{obs} are reduced by one for each parameter that must be estimated.

Soccer Example

Test the hypothesis that the number of goals scored during regulation time for the 232 soccer matches stored in the data frame `SOCCER` has a Poisson cdf with $\lambda = 2.5$ with the chi-square goodness-of-fit test and an α level of 0.05. Produce a histogram showing the number of observed goals scored during regulation time and superimpose on the histogram the number of goals that are expected to be made when the distribution of goals follows a Poisson distribution with $\lambda = 2.5$.

Soccer Solution

- Since the number of categories for a Poisson distribution is theoretically infinite, a table is first constructed of the observed number of goals to get an idea of reasonable categories.

```
library(PASWR2)
xtabs(~goals, data = SOCCER)
```

```
goals
 0   1   2   3   4   5   6   7   8
19 49 60 47 32 18   3   3   1
```

Soccer Solution

Based on the table, a decision is made to create categories for 0, 1, 2, 3, 4, 5, and 6 or more goals. Under the null hypothesis that $F_0(x)$ is a Poisson distribution with $\lambda = 2.5$, the probabilities of scoring 0, 1, 2, 3, 4, 5, and 6 or more goals are computed with R as follows:

```
PX <- c(dpois(0:5, 2.5), ppois(5, 2.5, lower = FALSE))
PX[1:4] # Probabilities for categories 0, 1, 2, 3
```

```
[1] 0.0820850 0.2052125 0.2565156 0.2137630
```

```
PX[4:6] # Probabilities for categories 4, 5, and 6 or more
```

```
[1] 0.21376302 0.13360189 0.06680094
```

Soccer Solution

Since there were a total of $n = 232$ soccer games, the expected number of goals for the six categories is simply $232 \times PX$.

```
EX <- 232*PX
OB <- c(as.vector(xtabs(~goals, data = SOCCER)[1:6]),
        sum(xtabs(~goals, data = SOCCER)[7:9]))
OB
```

```
[1] 19 49 60 47 32 18 7
```

```
ans <- cbind(PX, EX, OB)
row.names(ans) <- c(" X=0", " X=1", " X=2",
                    " X=3", " X=4", " X=5", "X>=6")
```

Soccer Solution

ans

	PX	EX	OB
X=0	0.08208500	19.043720	19
X=1	0.20521250	47.609299	49
X=2	0.25651562	59.511624	60
X=3	0.21376302	49.593020	47
X=4	0.13360189	30.995638	32
X=5	0.06680094	15.497819	18
X>=6	0.04202104	9.748881	7

Soccer Solution

The null and alternative hypotheses for using the chi-square goodness-of-fit test to test the hypothesis that the number of goals scored during regulation time for the 232 soccer matches stored in the data frame `SOCGER` has a Poisson cdf with $\lambda = 2.5$ are

$$H_0 : F_X(x) = F_0(x) \sim \text{Pois}(\lambda = 2.5) \text{ for all } x \text{ versus}$$

$$H_1 : F_X(x) \neq F_0(x) \text{ for some } x.$$

Soccer Solution

- The test statistic chosen is χ_{obs}^2 .
- Reject if $\chi_{\text{obs}}^2 > \chi_{1-\alpha; k-1}^2$.

```
chi.obs <- sum((OB-EX)^2/EX)
chi.obs
```

```
[1] 1.39194
```

Soccer Solution

```
chisq.test(x = OB, p = PX)
```

Chi-squared test for given probabilities

data: OB

X-squared = 1.3919, df = 6, p-value = 0.9663

Soccer Solution

$$1.3919402 = \chi_{\text{obs}}^2 \stackrel{?}{>} \chi_{0.95;6}^2 = 12.5915872.$$

The p -value is 0.9663469.

```
p.val <- pchisq(chi.obs, 7-1, lower = FALSE)
p.val
```

```
[1] 0.9663469
```

Soccer Solution

- Since $\chi_{\text{obs}}^2 = 1.3919402$ is not greater than $\chi_{0.95;6}^2 = 12.5915872$, fail to reject H_0 .
- Since the p -value = 0.9663469 is greater than 0.05, fail to reject H_0 .

Soccer Solution


English Conclusion: There is no evidence to suggest that the true **cdf** does not equal the Poisson distribution with $\lambda = 2.5$ for at least one x .

Soccer Solution

The following code can be used to create a histogram with superimposed expected goals.

```
hist(SOCCER$goals, breaks = c((-0.5 + 0):(8 + 0.5)),
     col = "lightblue",
     xlab = "Goals scored", ylab = "",
     freq = TRUE, main = "")
x <- 0:8
fx <- (dpois(0:8, lambda = 2.5))*232
lines(x, fx, type = "h")
lines(x, fx, type = "p", pch = 16)
```

Soccer Solution



All Parameters Known

- Bansley et al. (1992) investigated the relationship between month of birth and achievement in sport. Birth dates were collected for players in teams competing in the 1990 World Cup soccer games.

```
Observed <- c(150, 138, 140, 100)
names(Observed) <- c("Aug-Oct", "Nov-Jan",
                    "Feb-April", "May-July")
```

Observed

Aug-Oct	Nov-Jan	Feb-April	May-July
150	138	140	100

All Parameters Known

We wish to test whether these data are consistent with the hypothesis that birthdays of soccer players are uniformly distributed across the four quarters of the year. Let P_i denote the probability of a birth occurring in the i^{th} quarter; the hypotheses are as follows:

$H_0 : p_1 = \frac{1}{4}, p_2 = \frac{1}{4}, p_3 = \frac{1}{4}, p_4 = \frac{1}{4}$ versus $H_A : p_i \neq \frac{1}{4}$ for at least one i .

There were a total of $n = 528$ players considered for this study, so the expected count for each quarter is $528/4 = 132$.

All Parameters Known

$$\chi_{obs}^2 = \sum_{i=1}^k \frac{(O_i - E_i)^2}{E_i} = \frac{(150-132)^2}{132} + \frac{(138-132)^2}{132} + \frac{(140-132)^2}{132} + \frac{(100-132)^2}{132} = 10.97$$

```
(chi_obs <- sum((Observed - 132)^2/132))
```

```
[1] 10.9697
```

Or

```
chisq.test(Observed, p = c(1/4, 1/4, 1/4, 1/4))$stat
```

X-squared

10.9697

All Parameters Known

```
chisq.test(Observed, p = c(1/4, 1/4, 1/4, 1/4)) -> CST
CST
```

Chi-squared test for given probabilities

data: Observed

X-squared = 10.97, df = 3, p-value = 0.01189

CST\$observed

Aug-Oct	Nov-Jan	Feb-April	May-July
150	138	140	100

CST\$expected

Aug-Oct	Nov-Jan	Feb-April	May-July
132	132	132	132

All Parameters Known

```
(pvalue <- pchisq(CST$stat, 3, lower = FALSE))
```

```
  X-squared  
0.01189087
```

```
# Or  
CST$p.value
```

```
[1] 0.01189087
```

All Parameters Known - Conclusion

Given the p - *value* of 0.012 evidence suggests birthdays for World Cup soccer players are not uniformly distributed.

All Parameters Known - Example 2

Suppose you draw 100 numbers at random from an unknown distribution. Thirty values fall in the interval $(0, 0.25]$, 30 fall in $(0.25, 0.75]$, 22 fall in $(0.75, 1.25]$, and the rest fall in $(1.25, \infty]$. Your friend claims that the distribution is exponential with parameter $\lambda = 1$. Do you believe her?

- A random variable X has the exponential distribution with parameter $\lambda > 0$ if its **pdf** is

$$f(x) = \lambda e^{-\lambda x}, \quad x \geq 0.$$

All Parameters Known - Example 2

We wish to test the following:

H_0 : The data are from an exponential distribution with $\lambda = 1$.

H_A : The data are not from an exponential distribution with $\lambda = 1$.

All Parameters Known - Example 2

Given $X \sim \text{Exp}(\lambda = 1)$. The probabilities for each interval are as follows:

$$p_1 = P(0 \leq X \leq 0.25) = \int_0^{0.25} e^{-x} dx = 0.2211992$$

$$p_2 = P(0.25 \leq X \leq 0.75) = \int_{0.25}^{0.75} e^{-x} dx = 0.3064342$$

$$p_3 = P(0.75 \leq X \leq 1.25) = \int_{0.75}^{1.25} e^{-x} dx = 0.1858618$$

$$p_4 = P(1.25 \leq X \leq \infty) = \int_{1.25}^{\infty} e^{-x} dx = 0.2865048$$

All Parameters Known - Example 2

```
p1 <- pexp(0.25, 1)
p2 <- pexp(0.75, 1) - pexp(0.25, 1)
p3 <- pexp(1.25, 1) - pexp(0.75, 1)
p4 <- pexp(1.25, 1, lower = FALSE)
ps <- c(p1, p2, p3, p4)
ps
```

[1] 0.2211992 0.3064342 0.1858618 0.2865048

All Parameters Known - Example 2

```
EXP <- ps*100  
EXP
```

```
[1] 22.11992 30.64342 18.58618 28.65048
```

```
OBS <- c(30, 30, 22, 18)  
test_stat <- sum((OBS - EXP)^2/EXP)  
test_stat
```

```
[1] 7.406963
```


All Parameters Known - Example 2

```
# Another approach  
chisq.test(OBS, p = ps)
```

Chi-squared test for given probabilities

```
data:  OBS  
X-squared = 7.407, df = 3, p-value = 0.06  
  
pvalue <- chisq.test(OBS, p = ps)$p.value  
pvalue  
  
[1] 0.05999777
```

All Parameters Known - Example 2 - Conclusion

If you test using $\alpha = 0.05$, you will fail to reject the null hypothesis since the $p - value = 0.0599 > \alpha = 0.05$. There is not convincing evidence that the data do not come from an $\text{Exp}(\lambda = 1)$.

Section 2

Chi-Square Tests of Independence

Example

```
library(PASWR2)
(xtabs(~sex + survived, data = TITANIC3) -> T1)
```

	survived	
sex	0	1
female	127	339
male	682	161

```
chisq.test(T1, correct = FALSE) -> CST
CST
```

Pearson's Chi-squared test

```
data:  T1
X-squared = 365.89, df = 1, p-value < 2.2e-16
```

Example

```
(EXP <- CST$expected)
```

	survived	
sex	0	1
female	288.0015	177.9985
male	520.9985	322.0015

```
(OBS <- CST$observed)
```

	survived	
sex	0	1
female	127	339
male	682	161

```
(chi_obs <- sum((OBS - EXP)^2/EXP))
```

```
[1] 365.8869
```

Section 3

Chi-Square Tests of Homogeneity

Example

- Data will often come summarized in contingency tables.

```
DP <- c(67, 76, 57, 48, 73, 79)
MDP <- matrix(data = DP, nrow = 2, byrow = TRUE)
dimnames(MDP) <- list(Pop = c("Drug", "Placebo"),
  Status = c("Improve", "No Change", "Worse"))
TDP <- as.table(MDP)
TDP
```

Pop	Status		
	Improve	No Change	Worse
Drug	67	76	57
Placebo	48	73	79

Putting the data back in a tidy format

```
library(tidyverse)
NT <- TDP %>%
  tibble::as_tibble() %>%
  uncount(n)
head(NT, 3)
```

```
# A tibble: 3 x 2
  Pop    Status
<chr> <chr>
1 Drug  Improve
2 Drug  Improve
3 Drug  Improve
```