

ME 760 Homework 4

Alan Burl

October 31, 2020

1. Verify the contour integral $\int_C [2xy^2dx + 2x^2ydy + dz]$ is independent of the path. Evaluate this integral between the points $(0, 0, 0)$ and (a, b, c) .

A line integral with the vector function of the form $\mathbf{F} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ is path independent if $\text{curl}\mathbf{F} = 0$. The curl of the vector is found and evaluated below using equation 1.

$$\text{curl}\mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = 0 \quad (1)$$

Expansion of the above yields the following:

$$\text{curl}\mathbf{F} = \frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}\mathbf{i} - \frac{\partial R}{\partial x} - \frac{\partial P}{\partial z}\mathbf{j} + \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\mathbf{k} \quad (2)$$

The two non-trivial partial derivatives are shown below:

$$\frac{\partial Q}{\partial x} = 4xy$$

$$\frac{\partial P}{\partial y} = 4xy$$

Substituting the values into equation 2, yields $\text{curl}\mathbf{F} = 0$, thus the contour integral is independent of the path.

Evaluation of this leads to $f = x^2 + y^2 + z$. Thus evaluating using the form $f(B) - f(A)$ yields:

$$\boxed{a^2b^2 + c}$$

2. Given the parametric form of a cone $r(u, v) = [u \cos v, u \sin v, cu]$

- (a) find an explicit representation of the form $z = f(x, y)$
 - (b) find and identify the paramer curves definid as $u = \text{const}$ and $v = \text{const}$
 - (c) find the normal vector N to the conical surface
- (a) Let $x, y, z = u \cos v, u \sin v, cu$ respectively. Squaring x and y and summing them yields: $x^2 + y^2 = u^2 (\cos^2 v + \sin^2 v)$. Solving for u from this: $u = \sqrt{x^2 + y^2}$. Using this to substitute into z results in the explicity form of the equation:

$$\boxed{z = c\sqrt{x^2 + y^2}} \quad (3)$$

- (b) From (a), it is seen that $\boxed{u = \sqrt{x^2 + y^2}}$. v is found from the division of x and y . Using this and trigonemetric definitions yields: $\boxed{v = \tan^{-1} \frac{x}{y}}$.

- (c) The normal vector is found using equation 4

$$\nabla f = 0 \quad (4)$$

Differentiation of the explicit form with respect to x, y, z results in the normal vector:

$$\boxed{\mathbf{N} = \frac{2xc}{\sqrt{x^2 + y^2}} \mathbf{i} + \frac{2yc}{\sqrt{x^2 + y^2}} \mathbf{j} - \mathbf{k}}$$

3. In class we discussed surface integrals without regard to orientation. By reparameterizing the surface integral could be written as

$$I = \int \int_s G(s) dS = \int \int_R G(r(u, v)) |N(u, v)| du dv$$

- (a) Consider the case $G = z$ and the surface S is the hemisphere $x^2 + y^2 + z^2 = 9$ with $z \geq 0$. Use polar coordinates and evaluate the right hand side of the above result.
- (b) The surface S is also given explicitly by $z = f(x, y) = \sqrt{9 - x^2 - y^2}$. For such cases the surface integral can be rewritten as

$$\int \int_S G(r) dA = \int \int_{R^*} G(x, y, f(x, y)) \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dx dy \quad (5)$$

Evaluate the right-hand side for this result.