

Quantum States of Light in Cavity QED: A Computational Study of Wigner Function Dynamics and Atom–Field Entanglement in the Jaynes–Cummings Model

Nguyen Khoi Nguyen (Alan)*

Department of Electrical and Computer Engineering, Boston University, Boston, MA 02215

(Dated: September 17, 2026)

We present a computational study of quantum light–matter interaction in the Jaynes–Cummings (JC) model, combining analytical derivations with original numerical simulations. Starting from the electric-dipole coupling Hamiltonian, we derive the JC model under the rotating-wave approximation, solve for the dressed eigenstates and their limiting behavior, and obtain analytical expressions for the collapse time $t_c = O(1/g)$ and the revival time $t_r = 2\pi\sqrt{n}/g$. We introduce Fock, coherent, squeezed, and thermal states and their Wigner function representations, then solve the full JC dynamics using the Lindblad master equation. The principal computational results are: (i) time-resolved Wigner function snapshots revealing the formation of a Schrödinger cat state in the cavity field at half the revival time, with Wigner negativity volume $\delta = 0.85$, field purity 0.96, and fidelity 0.78 with respect to a two-component cat state; at this instant the atom and field approximately *disentangle* (reduced atomic entropy $S \approx 0.14$ bit), so the cat is a nearly pure state of the field alone; (ii) a systematic comparison of the reduced atomic entropy and of the logarithmic negativity across coherent, thermal, squeezed, and Fock initial field states, demonstrating that the half-revival disentanglement occurs only for the coherent state, and that for a thermal field a saturated entropy of one bit coexists with a logarithmic negativity of only ≈ 0.2 ; and (iii) a quantitative study of decoherence, showing that a cavity decay rate as low as $\kappa/g = 0.02$ reduces the cat-state Wigner negativity by over 80%. All simulations are performed in QuTiP with reproducible code available at https://github.com/alanknguyen/QSOL_CQED.

I. INTRODUCTION

Coherent manipulation of quantum two-level systems is a cornerstone of quantum technology. Rabi oscillations, the periodic cycling of population between two quantum states under a resonant driving field, provide the physical mechanism for controlling qubit states [1, 2]. When a two-level system is driven on resonance, its state oscillates between the ground and excited state at a rate known as the Rabi frequency, which is set by the coupling strength to the field [3, 4]. By choosing the duration and phase of the driving pulse, one can prepare arbitrary superpositions or implement reliable state flips, operations essential for quantum gates.

The Jaynes–Cummings (JC) model [5], which describes the interaction of a single two-level atom with a single quantized cavity mode, is the simplest fully quantum model of light–matter interaction. Despite its simplicity, the JC model predicts a rich set of phenomena with no classical analog: vacuum Rabi splitting, collapse and revival of Rabi oscillations [6], and the dynamical generation of Schrödinger cat states in the cavity field [7]. These predictions have been confirmed in pioneering cavity quantum electrodynamics (QED) experiments with Rydberg atoms in microwave cavities [8, 9] and, more recently, in circuit QED with superconducting qubits coupled to microwave resonators [10].

While the JC model is a textbook topic, most treatments present the collapse-revival phenomenon qualita-

tively and do not explore the associated phase-space dynamics or entanglement structure in detail. In this paper, we combine a self-contained theoretical introduction with original computational results that go beyond standard treatments in two ways. First, we compute the time-resolved Wigner function of the cavity field during JC dynamics, directly visualizing the formation and destruction of cat states in phase space and quantifying their non-classicality via the Wigner negativity volume. Second, we systematically compare the atom–field entanglement across four classes of initial field states (coherent, thermal, squeezed vacuum, and Fock), using the reduced entropy where the global state is pure and the logarithmic negativity where it is mixed, revealing how the field’s photon statistics govern the entanglement dynamics and, in particular, that the atom and field approximately disentangle at half the revival time only for a coherent field. We also study the effect of cavity dissipation via the Lindblad master equation, quantifying how rapidly decoherence destroys the non-classical features.

The paper is organized as follows. Section II reviews the quantization of the electromagnetic field, introduces the principal quantum states of light and the Wigner function formalism, derives the Jaynes–Cummings Hamiltonian from first principles including the rotating-wave approximation, solves for the dressed states and their limiting behavior, and presents analytical derivations of the collapse and revival timescales. Section III describes the computational methods. Section IV presents the simulation results. Section V discusses criteria for distinguishing classical and quantum light. Section VI reviews the experimental observation

* Also at Physics Department, Boston University.

of these phenomena across four platforms. Section VII concludes.

II. THEORETICAL FRAMEWORK

A. Quantum harmonic oscillator and field quantization

The quantization of the harmonic oscillator underpins all of quantum optics, since each mode of the electromagnetic field can be treated as a harmonic oscillator. Classically, a harmonic oscillator of mass m and angular frequency ω has the Hamiltonian $H = p^2/(2m) + \frac{1}{2}m\omega^2 x^2$. In quantum mechanics, we promote x and p to operators satisfying $[\hat{x}, \hat{p}] = i\hbar$ and introduce ladder operators

$$a = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} + \frac{i}{m\omega} \hat{p} \right), \quad a^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} - \frac{i}{m\omega} \hat{p} \right), \quad (1)$$

which satisfy $[a, a^\dagger] = 1$. The Hamiltonian becomes $H = \hbar\omega(a^\dagger a + \frac{1}{2})$, and the number operator $\hat{n} = a^\dagger a$ has eigenstates $|n\rangle$ (Fock states) with $n = 0, 1, 2, \dots$. The ladder operators act as $a^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle$ and $a|n\rangle = \sqrt{n}|n-1\rangle$, giving the quantized energy spectrum $E_n = \hbar\omega(n + \frac{1}{2})$ with equally spaced levels separated by $\hbar\omega$. The ground state $|0\rangle$ carries zero-point energy $\frac{1}{2}\hbar\omega$, reflecting irreducible vacuum fluctuations.

To describe light quantum mechanically, we expand the electromagnetic field in a complete set of modes (plane waves or cavity standing waves). Each mode of frequency ω_k becomes a quantum harmonic oscillator with operators a_k and $a_k^\dagger$. The electric field operator for a single mode takes the form

$$\hat{E}(t) = i\mathcal{E}_0 \left(a_k e^{-i\omega_k t} - a_k^\dagger e^{i\omega_k t} \right), \quad (2)$$

where $\mathcal{E}_0 = \sqrt{\hbar\omega_k/(2\varepsilon_0 V)}$ is the single-photon field amplitude and V is the quantization volume. The operator a_k annihilates one photon in mode k , while $a_k^\dagger$ creates one. The total field Hamiltonian is

$$H_{\text{field}} = \sum_k \hbar\omega_k \left(a_k^\dagger a_k + \frac{1}{2} \right). \quad (3)$$

The vacuum state has zero-point energy $\frac{1}{2} \sum_k \hbar\omega_k$, and excitations are described by multi-mode Fock states $|n_1, n_2, \dots\rangle$ specifying the photon count in each mode.

B. Quantum states of light

Fock states $|n\rangle$ are eigenstates of the photon number operator with a definite number of photons. They have no classical analog: a classical field cannot have an exact integer photon count. Fock states exhibit the quantum granularity of the field; for example, a single-photon state

$|1\rangle$ produces at most one detection event, and the vacuum $|0\rangle$ produces none.

Coherent states $|\alpha\rangle$, for complex amplitude α , are the most important states in quantum optics. They are defined as eigenstates of the annihilation operator, $a|\alpha\rangle = \alpha|\alpha\rangle$, or equivalently as displaced vacuum states $|\alpha\rangle = D(\alpha)|0\rangle$ with displacement operator $D(\alpha) = \exp(\alpha a^\dagger - \alpha^* a)$. Expanding in the Fock basis yields a Poissonian photon-number distribution:

$$P_n = |\langle n|\alpha\rangle|^2 = e^{-|\alpha|^2} \frac{|\alpha|^{2n}}{n!}, \quad (4)$$

with mean $\bar{n} = |\alpha|^2$ and variance $\text{Var}(n) = \bar{n}$. The equality of variance and mean is a hallmark of classical (shot-noise-limited) light. Coherent states minimize the Heisenberg uncertainty relation for field quadratures, with equal noise in both conjugate components at the vacuum level. They are often called the “most classical” quantum states because their mean electric field follows a classical sinusoid and their Glauber–Sudarshan P -distribution is a delta function, $P(\beta) = \delta^2(\beta - \alpha)$ [11]. Laser light above threshold is well described by a coherent state with large $\bar{n}$.

Squeezed states are non-classical states in which the quantum noise in one field quadrature is reduced below the vacuum level, at the expense of increased noise in the conjugate quadrature. Defining the quadrature $X_\phi = \frac{1}{2}(ae^{-i\phi} + a^\dagger e^{i\phi})$, a vacuum or coherent state has $\langle(\Delta X_\phi)^2\rangle = \frac{1}{4}$ independent of ϕ (in units with $\hbar = 1$). A squeezed state has $\langle(\Delta X_{\phi_0})^2\rangle < \frac{1}{4}$ for some optimal phase ϕ_0 . The squeeze operator $S(\zeta) = \exp[\frac{1}{2}(\zeta^* a^2 - \zeta a^{\dagger 2})]$ with $\zeta = re^{i\varphi}$ applied to the vacuum yields uncertainties

$$\langle(\Delta X_{\varphi/2})^2\rangle = \frac{1}{4}e^{-2r}, \quad \langle(\Delta X_{\varphi/2+\pi/2})^2\rangle = \frac{1}{4}e^{2r}. \quad (5)$$

Squeezed states can be produced by parametric down-conversion or four-wave mixing. Because they exhibit noise below the vacuum level in one observable, they have no classical description (their P -representation involves derivatives of delta functions). Since 2019, squeezed light has been injected into the LIGO and Virgo gravitational-wave observatories to reduce quantum noise, significantly improving sensitivity [13, 14], a direct application of non-classical light to precision measurement.

C. Phase-space representations: the Wigner function

The Wigner function provides a quasi-probability distribution in phase space that is particularly useful for visualizing quantum states and identifying non-classical features. For a single-mode field with density operator ρ , it is defined as [15]

$$W(x, p) = \frac{1}{\pi\hbar} \int_{-\infty}^{\infty} \langle x - y | \rho | x + y \rangle e^{2ipy/\hbar} dy, \quad (6)$$

where x and p are conjugate phase-space quadratures. Unlike a true probability distribution, $W(x, p)$ can take negative values; such negative regions are a sufficient (though not necessary) indicator of non-classicality [16].

For a coherent state $|\alpha\rangle$, the Wigner function is a Gaussian centered at $(\sqrt{2} \operatorname{Re} \alpha, \sqrt{2} \operatorname{Im} \alpha)$:

$$W_\alpha(x, p) = \frac{1}{\pi} \exp \left[- (x - \sqrt{2} \operatorname{Re} \alpha)^2 - (p - \sqrt{2} \operatorname{Im} \alpha)^2 \right], \quad (7)$$

which is everywhere non-negative. For a Fock state $|n\rangle$,

$$W_n(x, p) = \frac{(-1)^n}{\pi} L_n(2(x^2 + p^2)) e^{-(x^2 + p^2)}, \quad (8)$$

where L_n is the n -th Laguerre polynomial; for $n \geq 1$ this exhibits pronounced negative regions. A squeezed vacuum has an elliptical Gaussian Wigner function with widths e^{-r} and e^r along the squeezed and anti-squeezed quadratures; it remains non-negative, though its P -representation is singular.

A quantitative measure of non-classicality is the Wigner negativity volume [17]:

$$\delta = \int |W(x, p)| dx dp - 1. \quad (9)$$

Since $\int W dx dp = 1$, δ equals twice the integrated volume of the negative part of W . Every state with a non-negative Wigner function has $\delta = 0$; this includes all classical states but also, for example, the squeezed vacuum, so $\delta > 0$ is a sufficient but not a necessary witness of non-classicality. We use δ as a quantitative probe throughout the simulations in Sec. IV.

D. Derivation of the Jaynes–Cummings Hamiltonian

We derive the Jaynes–Cummings model [5] from the electric-dipole form of the light–matter interaction, the dipole-gauge counterpart of the minimal-coupling Hamiltonian $(\mathbf{p} - q\mathbf{A})^2/2m$ to which it is related by the Power–Zienau–Woolley transformation [2]. In the dipole approximation, the interaction between a two-level atom and a single quantized cavity mode of frequency ω_c is

$$H_{\text{int}} = -\mathbf{d} \cdot \mathbf{E}(t), \quad (10)$$

where $\mathbf{d}$ is the atomic dipole operator and $\mathbf{E}(t) \propto \boldsymbol{\varepsilon}(a + a^\dagger)$ is the quantized electric field, with $\boldsymbol{\varepsilon}$ the polarization vector. Restricting to two atomic levels $|g\rangle$ and $|e\rangle$ with transition frequency ω_a , the dipole operator is $\mathbf{d} = \mathbf{d}_0(\sigma^+ + \sigma^-)$, where $\sigma^+ = |e\rangle\langle g|$, $\sigma^- = |g\rangle\langle e|$, and $\mathbf{d}_0$ is the transition dipole matrix element. The total Hamiltonian is then

$$H = \hbar\omega_c a^\dagger a + \frac{\hbar\omega_a}{2} \sigma_z + \hbar g(\sigma^+ + \sigma^-)(a + a^\dagger), \quad (11)$$

where $\sigma_z = |e\rangle\langle e| - |g\rangle\langle g|$ and the coupling constant g is proportional to $\mathbf{d}_0 \cdot \boldsymbol{\varepsilon}$ and the single-photon field amplitude. Equation (11) is the quantum Rabi model. It contains both energy-conserving terms ($a\sigma^+$ and $a^\dagger\sigma^-$) and counter-rotating terms ($a^\dagger\sigma^+$ and $a\sigma^-$) that do not conserve excitation number.

In the regime $g \ll \omega_a, \omega_c$ and near resonance ($|\omega_a - \omega_c| \ll \omega_a + \omega_c$), the counter-rotating terms oscillate at frequency $\omega_a + \omega_c$ in the interaction picture and average to zero. Dropping them gives the rotating-wave approximation (RWA) and the Jaynes–Cummings Hamiltonian:

$$H_{\text{JC}} = \hbar\omega_c a^\dagger a + \frac{\hbar\omega_a}{2} \sigma_z + \hbar g (a^\dagger \sigma^- + a \sigma^+). \quad (12)$$

The term $a^\dagger\sigma^-$ describes atomic de-excitation with photon emission, and $a\sigma^+$ describes photon absorption with atomic excitation. A key symmetry of H_{JC} is conservation of the total excitation number $N = a^\dagger a + \sigma^+ \sigma^-$, which commutes with H_{JC} . The Hilbert space therefore decomposes into invariant two-dimensional subspaces $\{|e, n\rangle, |g, n+1\rangle\}$ for each $n \in \mathbb{N}$, enabling exact analytic solutions.

E. Dressed states and energy eigenvalues

Within the subspace of $n+1$ total excitations, the Hamiltonian in the basis $\{|e, n\rangle, |g, n+1\rangle\}$ reduces (after subtracting the common energy $(n + \frac{1}{2})\hbar\omega_c$) to the 2×2 matrix

$$H^{(n)} = \frac{\hbar}{2} \begin{pmatrix} \Delta & \Omega_0 \sqrt{n+1} \\ \Omega_0 \sqrt{n+1} & -\Delta \end{pmatrix}, \quad (13)$$

where $\Delta = \omega_a - \omega_c$ is the detuning and $\Omega_0 = 2g$ is the single-photon Rabi frequency. The coupling matrix element $\langle e, n | H_{\text{JC}} | g, n+1 \rangle = \hbar g \sqrt{n+1}$ follows from the bosonic commutation relations. The eigenvalues are

$$E_\pm(n) = \hbar\omega_c (n + \frac{1}{2}) \pm \frac{\hbar}{2} \Omega_n(\Delta), \quad (14)$$

where the generalized n -photon Rabi frequency is

$$\Omega_n(\Delta) = \sqrt{\Delta^2 + \Omega_0^2(n+1)}. \quad (15)$$

The corresponding eigenstates (dressed states) are

$$|n, +\rangle = \cos \frac{\theta_n}{2} |e, n\rangle + \sin \frac{\theta_n}{2} |g, n+1\rangle, \quad (16)$$

$$|n, -\rangle = \sin \frac{\theta_n}{2} |e, n\rangle - \cos \frac{\theta_n}{2} |g, n+1\rangle, \quad (17)$$

with mixing angle defined by $\tan \theta_n = \Omega_0 \sqrt{n+1} / \Delta$. These are entangled atom–photon states whose character interpolates between atom-like and photon-like as Δ is varied. The energy splitting between the two dressed states is $\hbar\Omega_n(\Delta)$.

F. Limiting cases

Resonance ($\Delta = 0$). The mixing angle is $\theta_n = \pi/2$ and the dressed states become equal superpositions:

$$|n, \pm\rangle = \frac{1}{\sqrt{2}} (|e, n\rangle \pm |g, n+1\rangle). \quad (18)$$

The splitting reduces to $\hbar\Omega_0\sqrt{n+1}$. For $n = 0$ (one total excitation), the splitting is $\hbar\Omega_0 = 2\hbar g$, the vacuum Rabi splitting. This is directly analogous to the normal-mode splitting of two coupled oscillators, and its spectroscopic observation [19] was among the first confirmations of the JC model.

Large detuning ($|\Delta| \gg \Omega_0\sqrt{n+1}$). The mixing angle is small and the dressed states are only weakly entangled. Expanding $\Omega_n \approx |\Delta| + \Omega_0^2(n+1)/(2|\Delta|)$ gives energy shifts

$$E_{\pm}(n) \approx \hbar\omega_c (n + \frac{1}{2}) \pm \frac{\hbar}{2} \left(|\Delta| + \frac{\Omega_0^2(n+1)}{2|\Delta|} \right). \quad (19)$$

The second-order corrections $\propto g^2/\Delta$ are the ac Stark (light) shifts [9]. In this regime the eigenstates approach the bare states $|e, n\rangle$ and $|g, n+1\rangle$, and the atom-field coupling manifests primarily as dispersive frequency shifts rather than coherent population transfer. The transition from resonant Rabi splitting to dispersive shifts as Δ is varied produces an avoided crossing of width $\hbar\Omega_0\sqrt{n+1}$ at $\Delta = 0$.

G. Rabi oscillations from a Fock-state field

Consider the atom initially excited and the field in a Fock state with n photons: $|\Psi(0)\rangle = |e, n\rangle$. This state is not an eigenstate of the coupled system; it is a superposition of the two dressed states in the $(n+1)$ -excitation manifold. On resonance,

$$|e, n\rangle = \frac{1}{\sqrt{2}} (|n, +\rangle + |n, -\rangle). \quad (20)$$

Evolving each eigenstate with its eigenphase gives the time-dependent state (dropping the common phase):

$$|\Psi(t)\rangle = \cos(\frac{1}{2}\Omega_n t) |e, n\rangle - i \sin(\frac{1}{2}\Omega_n t) |g, n+1\rangle, \quad (21)$$

where $\Omega_n = \Omega_0\sqrt{n+1}$ and the sign of the second term follows from $e^{-iH_{JC}t/\hbar}$ with the coupling $+\hbar g(a^\dagger\sigma^- + a\sigma^+)$. The excited-state probability oscillates sinusoidally:

$$P_e(t) = \cos^2\left(\frac{\Omega_0\sqrt{n+1}}{2} t\right), \quad (22)$$

at the n -photon Rabi frequency $\Omega_n = \Omega_0\sqrt{n+1}$. For $n = 0$, this is the vacuum Rabi oscillation: an excited atom in an empty cavity emits and reabsorbs a single

photon periodically at frequency Ω_0 . For $n \gg 1$, the Rabi frequency $\Omega_0\sqrt{n+1} \approx \Omega_0\sqrt{n}$ is proportional to the classical field amplitude $\propto \sqrt{n}$, recovering the semiclassical limit. These coherent oscillations require the strong-coupling regime ($g \gg \kappa, \gamma$) so that the photon is not lost from the cavity during a Rabi cycle [9].

H. Collapse and revival of Rabi oscillations

Now suppose the atom is initially excited and the field is in a coherent state $|\alpha\rangle$ with mean photon number $\bar{n} = |\alpha|^2$. Expanding the coherent state in the Fock basis,

$$|\Psi(0)\rangle = |e\rangle \otimes |\alpha\rangle = e^{-\bar{n}/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |e, n\rangle, \quad (23)$$

and evolving each Fock component independently via Eq. (21), the excited-state probability at time t is

$$P_e(t) = \sum_{n=0}^{\infty} p(n) \cos^2\left(\frac{\Omega_0\sqrt{n+1}}{2} t\right), \quad (24)$$

where $p(n) = e^{-\bar{n}}\bar{n}^n/n!$ is the Poisson distribution. Using the identity $\cos^2\theta = \frac{1}{2}(1 + \cos 2\theta)$,

$$P_e(t) = \frac{1}{2} + \frac{1}{2} \sum_{n=0}^{\infty} p(n) \cos(\Omega_0\sqrt{n+1} t). \quad (25)$$

Equation (25) is a weighted superposition of cosines at incommensurate frequencies $\Omega_0\sqrt{n+1}$. The Poisson weights are peaked around $n \approx \bar{n}$ with width $\sqrt{\bar{n}}$, so at short times all components oscillate nearly in phase at the dominant frequency $\Omega_0\sqrt{\bar{n}+1}$.

Collapse. As time progresses, the components dephase because neighboring Fock states oscillate at slightly different frequencies. The spread of Rabi frequencies across the Poisson distribution (width $\sigma_n = \sqrt{\bar{n}}$) is

$$\delta\Omega \sim \left. \frac{\partial\Omega_n}{\partial n} \right|_{\bar{n}} \cdot \sigma_n = \frac{\Omega_0}{2\sqrt{\bar{n}}} \cdot \sqrt{\bar{n}} = \frac{\Omega_0}{2}. \quad (26)$$

The oscillations dephase when $\delta\Omega \cdot t_c \sim 1$, giving the collapse time

$$t_c \sim \frac{2}{\Omega_0} = \frac{1}{g}, \quad (27)$$

independent of $\bar{n}$ [6]. The numerical prefactor depends on the criterion adopted: linearizing $\sqrt{n+1}$ about $\bar{n}$ and summing Eq. (25) over the approximately Gaussian Poisson weights gives the envelope $P_e(t) - \frac{1}{2} \approx \frac{1}{2} \cos(\Omega_0\sqrt{\bar{n}} t) e^{-g^2 t^2/2}$, whose $1/e$ time is $\sqrt{2}/g$. Throughout this paper we use the operational definition $t_c \equiv 1/g$; in the simulations the oscillations are fully damped by $t \approx 2t_c$. After the collapse, $P_e(t)$ fluctuates around $1/2$ with negligible amplitude.

Revival. Despite the apparent loss of coherence, the oscillations return at a later time. Because the Rabi frequencies are $\Omega_0\sqrt{n+1}$ (a discrete set, not a continuum), the phases can realign modulo 2π . To estimate the revival time, we require adjacent Fock components near $n \approx \bar{n}$ to accumulate a phase difference of exactly 2π :

$$(\Omega_0\sqrt{n+2} - \Omega_0\sqrt{n+1})t_r = 2\pi. \quad (28)$$

For large n , $\sqrt{n+2} - \sqrt{n+1} \approx 1/(2\sqrt{n+1})$, giving

$$t_r \approx \frac{4\pi\sqrt{\bar{n}}}{\Omega_0} = \frac{2\pi\sqrt{\bar{n}}}{g}. \quad (29)$$

The ratio $t_r/t_c \approx 2\pi\sqrt{\bar{n}}$ shows that for large $\bar{n}$, the revival occurs many collapse times later, with a broad plateau of near-zero oscillation amplitude in between.

The revival is a purely quantum effect. A classical field with a continuous amplitude distribution would produce irreversible dephasing with no recurrence. The revival requires the *discrete* spectrum of Fock-state Rabi frequencies, and its observation constitutes direct evidence that the intracavity field is quantized [6, 9].

After the first revival, the components dephase again on a timescale $\sim t_c$, producing a second collapse. Further revivals occur quasi-periodically at integer multiples of t_r , though higher-order revivals become progressively less sharp as the accumulated phase errors grow [4].

For a thermal field (photon statistics given by the Bose-Einstein distribution, $p(n) = \bar{n}^n/(1+\bar{n})^{n+1}$), the distribution is much broader than Poisson ($\sigma_n \sim \bar{n}$ rather than $\sqrt{\bar{n}}$). The dephasing is correspondingly faster, and no pronounced revival is resolved within the simulated window; the evolution nevertheless remains unitary, so this is a loss of visibility rather than a strictly irreversible process. This contrast between coherent and thermal fields is a central result of our simulations (Sec. IV B).

During the collapse the atom and field become strongly entangled. Remarkably, at half the revival time, $t_r/2 = \pi\sqrt{\bar{n}}/g$, the two branches of the atomic state reconverge onto a common state that is independent of the initial atomic state, so that the atom and field approximately *disentangle*, and the field is left in a nearly pure Schrödinger cat state, a superposition of two coherent states with opposite phases [27–29]. This connection between Rabi collapse, transient entanglement, half-revival disentanglement, and cat-state formation is a central theme of our computational results.

I. Dressed-state spectroscopy

The dressed-state picture also explains spectroscopic signatures of strong atom–field coupling. In resonance fluorescence of a strongly driven two-level atom, the emission spectrum exhibits a central peak at the drive frequency and sidebands at $\omega \pm \Omega$, the Mollow triplet [18], which can be interpreted as transitions between dressed

states separated by $\hbar\Omega$. In cavity QED, a strongly coupled atom–cavity system shows normal-mode splitting: the cavity transmission spectrum splits into two peaks at $\omega_c \pm g$ for one excitation, a direct observation of the $n = 1$ dressed-state splitting of $2g$ [19]. As n increases, the splitting grows as $\sqrt{n+1}$, though thermal fluctuations may blur individual n -dependent peaks in practice.

J. Open quantum systems: the Lindblad master equation

Real cavity QED systems are coupled to their environment. The dominant dissipation channels are cavity photon loss at rate κ and atomic spontaneous emission into non-cavity modes at rate γ . Within the Markov approximation, the density matrix evolves according to the Lindblad master equation [20, 21]:

$$\frac{d\rho}{dt} = -\frac{i}{\hbar}[H_{\text{JC}}, \rho] + \kappa \mathcal{D}[a]\rho + \gamma \mathcal{D}[\sigma^-]\rho, \quad (30)$$

where the Lindblad dissipator is

$$\mathcal{D}[L]\rho = L\rho L^\dagger - \frac{1}{2}(L^\dagger L\rho + \rho L^\dagger L). \quad (31)$$

The strong-coupling regime, where quantum coherent effects dominate, requires $g \gg \kappa, \gamma$ so that the atom and field exchange energy many times before dissipation destroys coherence. In this regime, vacuum Rabi splitting is spectroscopically resolvable and collapse-revival dynamics can unfold before being damped.

Dissipation has a particularly dramatic effect on the cat states generated during JC dynamics. The decoherence rate of the interference fringes of a superposition of two coherent states separated by $\Delta\alpha$ in phase space is $\kappa|\Delta\alpha|^2/2$ [9, 22]; for the JC cat, whose components sit at $\pm i\sqrt{\bar{n}}$ (Sec. IV A), this is $2\kappa\bar{n}$, far exceeding the bare cavity decay rate κ . This rapid fragility of mesoscopic superpositions is a central result of our simulations.

K. Atom–field entanglement: von Neumann entropy

For a bipartite pure state $|\Psi\rangle_{AF}$ of the atom–field system, the degree of entanglement is uniquely quantified by the von Neumann entropy of either reduced subsystem [23]:

$$S(\rho_{\text{atom}}) = -\text{Tr}[\rho_{\text{atom}} \log_2 \rho_{\text{atom}}], \quad (32)$$

where $\rho_{\text{atom}} = \text{Tr}_{\text{field}}(|\Psi\rangle\langle\Psi|)$. For a two-level atom, S ranges from 0 (product state) to 1 bit (maximally entangled). The entropy is related to the purity of the reduced field state,

$$\mathcal{P} = \text{Tr}[\rho_{\text{field}}^2], \quad (33)$$

which equals unity for product states and decreases with increasing entanglement.

Equation (32) is an entanglement measure only when the global atom–field state is pure. If the global state is mixed, as for a thermal initial field or whenever dissipation acts, $S(\rho_{\text{atom}})$ counts classical as well as quantum correlations and can saturate at 1 bit for a weakly entangled state. In those cases we use the logarithmic negativity [30, 31]

$$E_{\mathcal{N}} = \log_2 \|\rho^{T_{\text{atom}}}\|_1, \quad (34)$$

where T_{atom} denotes partial transposition with respect to the atom and $\|\cdot\|_1$ the trace norm. $E_{\mathcal{N}}$ is an entanglement monotone for mixed states; it vanishes for separable states and equals 1 for a maximally entangled atom–field pair.

In the JC model, the entanglement dynamics are directly correlated with the collapse and revival of Rabi oscillations [24]. As the oscillations collapse, the atom becomes strongly entangled with the field ($S \rightarrow 1$ bit). At half the revival time, however, the system approximately refactorizes: S passes through a minimum, the atom approaches a pure state, and the field is left in a nearly pure cat state [27, 29]. Around the full revival the branches rephase and the entanglement grows again. This entropy–inversion correspondence was first analyzed by Phoenix and Knight [24, 29], and we verify it computationally in Sec. IV.

III. COMPUTATIONAL METHODS

All simulations are performed using QuTiP (Quantum Toolbox in Python, version 5.2) [25, 26], an open-source framework for simulating open and closed quantum systems. The cavity field is represented in a truncated Fock basis $\{|0\rangle, |1\rangle, \dots, |N_{\text{cav}} - 1\rangle\}$, and the joint atom–field Hilbert space has dimension $2N_{\text{cav}}$.

Fock space truncation. The truncation is chosen per initial state. For the coherent and Fock states with $\bar{n} = 10$ we use $N_{\text{cav}} = 35$ –50: the Poisson weight beyond $n = 35$ is below 10^{-9} , and every quantity reported at $t_r/2$ (entropy, purity, δ) agrees to four digits between $N_{\text{cav}} = 35$, 50, and 70. Thermal and squeezed-vacuum fields have much heavier photon-number tails (geometric, and $\propto \tanh^{2m} r/\sqrt{m}$ for $n = 2m$, respectively): at $N_{\text{cav}} = 50$ the truncated states would have $\langle n \rangle = 9.57$ and 9.15 instead of 10, so for these we use $N_{\text{cav}} = 150$, which reproduces $\langle n \rangle = 10$ to four digits and leaves a truncated weight below 10^{-6} . For the $\bar{n}$ scan we use $N_{\text{cav}} = 3\bar{n} + 20$. Trace normalization alone is not used as a convergence criterion.

Time evolution. All propagation uses QuTiP’s `mesolve` (version 5.2.3) with its default adaptive-step Adams–Moulton multistep integrator (`method='adams'`, absolute and relative tolerances 10^{-8} and 10^{-6}). Pure initial states without dissipation are propagated as state

vectors; the thermal initial field and all dissipative runs [Eq. (30)] propagate the full density matrix.

Wigner functions. At each snapshot time, we obtain the reduced field state $\rho_{\text{field}}(t) = \text{Tr}_{\text{atom}}[\rho(t)]$ via partial trace, then compute $W(x, p)$ on a 200×200 grid over $[-7, 7]^2$ (spacing 0.07, i.e. ten points per interference fringe of spacing $\pi/\sqrt{2\bar{n}} \approx 0.70$) using QuTiP’s `wigner` function [25] in the convention $a = (x + ip)/\sqrt{2}$, for which $\int W dx dp = 1$ (verified to 10^{-4} on the grid). The negativity δ of Eq. (9) is evaluated on the same grid.

Entanglement and cat-state measures. The von Neumann entropy $S(\rho_{\text{atom}})$ is computed at each time step from the eigenvalues of the 2×2 reduced atomic density matrix, the field purity $\mathcal{P}(t) = \text{Tr}[\rho_{\text{field}}^2]$ directly, and the logarithmic negativity of Eq. (34) from the trace norm of the partially transposed joint state. The cat-state fidelity is $F_{\text{cat}} = \max \langle \text{cat} | \rho_{\text{field}} | \text{cat} \rangle$ over normalized states $|\text{cat}\rangle \propto |\beta\rangle + e^{i\theta} |-\beta\rangle$, with β scanned in a neighborhood of $i\sqrt{\bar{n}}$ and $\theta \in [0, 2\pi)$.

Parameters. We work in dimensionless units with $g = 1$. The characteristic timescales are the collapse time $t_c \equiv 1/g$ and the revival time $t_r = 2\pi\sqrt{\bar{n}}/g$. Table I summarizes the simulation parameters.

TABLE I. Simulation parameters. All quantities in units where $g = 1$ and $\hbar = 1$.

Simulation	$\bar{n}$	N_{cav}	κ/g	γ/g
Wigner evolution	10	35	0	0
Entanglement (coherent, Fock)	10	50	0	0
Entanglement (thermal, squeezed)	10	150	0	0
$\bar{n}$ scaling	1–36	$3\bar{n} + 20$	0	0
Dissipative dynamics	10	35–50	0–0.2	0

All simulation code is available as Python scripts at https://github.com/alanknguyen/QSOL_CQED, and the release used for this article is archived on Zenodo [38].

IV. RESULTS AND DISCUSSION

A. Wigner function evolution and cat-state formation

We first consider unitary JC dynamics with the atom initially excited and the field in a coherent state $|\alpha\rangle$ with $\bar{n} = |\alpha|^2 = 10$. Figure 1 shows the atomic inversion $\langle \sigma_z(t) \rangle$, exhibiting the collapse–revival structure: rapid Rabi oscillations that dephase by $t \approx 2/g$, a quiescent interval, and a revival centered at $t_r \approx 2\pi\sqrt{10}/g \approx 19.9/g$.

The central computational result of this section is shown in Fig. 2, which displays the Wigner function of the reduced cavity field at six characteristic times. At $t = 0$, the Wigner function is a Gaussian displaced to $x \approx 4.5$, consistent with the initial coherent state. As the system evolves through the collapse region ($t \approx 2/g$), the distribution spreads along the circle $x^2 + p^2 = 2\bar{n}$ in phase space. At the midpoint of the collapse–revival

cycle ($t = t_r/2$), the Wigner function develops two well-separated lobes connected by interference fringes with alternating positive and negative values, the signature of a Schrödinger cat state, $|\Psi_{\text{cat}}\rangle \approx \mathcal{N}(|\beta\rangle + e^{i\theta}|\beta\rangle)$ with $\beta \approx i\sqrt{\bar{n}}$: the two components sit on the p axis, rotated by $\pm\pi/2$ from the initial coherent amplitude.

Figure 3 provides a detailed view of this cat state. The cross-section $W(x, p=0)$ shows deep negative excursions reaching $W \approx -0.25$, with Wigner negativity volume $\delta = 0.85$ (compared to $\delta = 0$ at $t = 0$). The best-fitting two-component cat has fidelity $F_{\text{cat}} = 0.78$ with the reduced field state, whose purity is $\mathcal{P} = 0.96$: the field is a nearly pure cat state rather than one branch of an entangled atom–field pair (Sec. IV B). The relative phase of the fit is $\theta \approx 1.2\pi$ and the photon-number parity is $\langle(-1)^{\hat{n}}\rangle = -0.64$, so the JC cat is closer to an odd than to an even cat, and its photon-number distribution displays the corresponding odd- n comb; its fidelity with the even cat $|\alpha\rangle + |-\alpha\rangle$ on the real axis is essentially zero. The interference fringes have characteristic spacing $\Delta x \approx \pi/\sqrt{2\bar{n}} \approx 0.7$, set by the separation of the two coherent components. These fringes are the phase-space manifestation of quantum interference between macroscopically distinguishable field states and serve as a definitive marker of non-classicality.

As the system approaches the revival time, the two lobes merge and the Wigner function partially re-localizes, though with residual non-classical features ($\delta = 0.31$ at $t = t_r$). The revival is not exact: higher-order terms in the photon-number distribution prevent perfect recurrence.

B. Atom–field entanglement dynamics

We now quantify atom–field entanglement, comparing four initial field states with the same mean photon number $\bar{n} = 10$: coherent, thermal, squeezed vacuum, and Fock. For the three pure initial fields the global state remains pure and $S(\rho_{\text{atom}})$ is the entanglement entropy; for the thermal field we report both S and the logarithmic negativity $E_{\mathcal{N}}$ of Eq. (34).

Figure 5 shows the inversion, entropy, and negativity for all four cases. The principal observations are as follows.

Coherent state. The entropy rises from $S = 0$ to ≈ 1 bit within $t \approx 2/g$, as the Rabi oscillations collapse. It then decreases steadily and reaches a minimum $S \approx 0.14$ bit ($E_{\mathcal{N}} = 0.36$) at $t = t_r/2 \approx 9.9/g$, in the middle of the quiescent interval. This is the half-revival disentanglement predicted by Gea-Banacloche [27]: the atom returns close to a pure state while the field is in the cat state of Fig. 3. Toward the revival the entanglement grows again; at t_r the entropy is $S \approx 0.79$ bit and oscillates between about 0.5 and 1 at the Rabi frequency. The cat state and the minimum of entanglement therefore coincide: the field is a nearly pure superposition, not one branch of an entangled pair.

Thermal state. The reduced entropy rises to $S = 1$ bit within $t \sim 1/g$ and stays within a few percent of the maximum thereafter, with no dip at $t_r/2$. This does not indicate maximal entanglement: the global state is mixed, and the logarithmic negativity remains small, $E_{\mathcal{N}} \approx 0.23$ on average (maximum 0.33). Most of the reduced-atom entropy is classical correlation inherited from the broad Bose–Einstein photon distribution, whose spread of Rabi frequencies ($\sigma_n \sim \bar{n}$) dephases the oscillations within a Rabi period and leaves no resolved revival in the simulated window.

Squeezed vacuum. The dynamics are intermediate: a rapid rise to $S \approx 1$ bit followed by quasi-periodic partial dips (to $S \approx 0.7$ –0.8), but no clean half-revival disentanglement. The super-Poissonian photon statistics of the squeezed vacuum (only even n populated, variance $2\bar{n}(\bar{n}+1)$) spread the Rabi frequencies over a much wider range than the Poisson distribution.

Fock state. With a definite photon number there is a single Rabi frequency, $\Omega_{10} = 2g\sqrt{11}$ [Eq. (21)]. The inversion oscillates with period $\pi/(g\sqrt{11}) \approx 0.95/g$, and the entropy, which depends on the populations only through the binary entropy $h(P_e)$ with $h(p) = h(1-p)$, oscillates between 0 and 1 bit with half that period, $\pi/(2g\sqrt{11}) \approx 0.47/g$ [Fig. 5(b)]. No collapse or revival occurs because there is no dephasing.

Figure 6 shows the coherent-state case in greater detail, plotting inversion, entropy, and field purity $\mathcal{P}(t)$ together. During the collapse the field purity drops to $\mathcal{P} \approx 0.5$: the reduced field is then an incoherent *mixture* of two nearly orthogonal branches, each correlated with a different atomic state (a coherent superposition of the two branches would have $\mathcal{P} = 1$). As the atomic branches reconverge, the purity climbs to $\mathcal{P} = 0.96$ at $t_r/2$, where the field is the nearly pure cat, and falls again toward the revival ($\mathcal{P} = 0.64$ at t_r). The three observables thus tell a single story: entanglement during the collapse turns the reduced field into a two-branch mixture, and disentanglement at $t_r/2$ restores the field’s coherence in the form of a cat state.

Figure 7 shows how these dynamics depend on $\bar{n}$. For $\bar{n} = 1$ the entropy oscillates quasi-periodically without a clean collapse. As $\bar{n}$ increases, the collapse still occurs at $t \sim 1/g$, but the half-revival minimum at $\pi\sqrt{\bar{n}}/g$ moves out and deepens ($S_{\text{min}} \approx 0.16, 0.11, 0.07$, and 0.05 bit for $\bar{n} = 9, 16, 25$, and 36), and the entangled plateau around the revival broadens. For $\bar{n} \gtrsim 9$ the disentanglement and the subsequent re-entanglement are cleanly resolved, consistent with the large- $\bar{n}$ analysis of Gea-Banacloche [28].

C. Decoherence of non-classical features

We now include cavity photon loss ($\kappa > 0$) via Eq. (30) and study its effect on the Wigner function and entanglement.

Figure 4 shows the Wigner function at the cat-state time $t = t_r/2$ for $\kappa/g = 0, 0.02, 0.05, 0.10$. The interfer-

ence fringes are progressively destroyed: at $\kappa/g = 0.02$, the Wigner negativity drops from $\delta = 0.85$ to $\delta = 0.14$; at $\kappa/g = 0.10$, the negativity is effectively zero ($\delta = 0.002$). Table II summarizes these results; the values are converged to three digits between $N_{\text{cav}} = 35$ and 50.

TABLE II. Wigner negativity δ and field purity $\mathcal{P}$ at $t = t_r/2$ for varying cavity decay rate, with $\bar{n} = 10$.

κ/g	Wigner negativity δ	Field purity $\mathcal{P}$	Cat state visible?
0.00	0.851	0.960	Yes
0.02	0.143	0.460	Marginal
0.05	0.022	0.418	No
0.10	0.002	0.386	No

This extreme sensitivity is expected: the fringe decoherence rate is $2\kappa\bar{n}$ (Sec. II J). For $\bar{n} = 10$, even $\kappa/g = 0.02$ gives a fringe-decoherence time $1/(2\kappa\bar{n}) = 2.5/g$, a quarter of the cat-formation time $t_r/2 \approx 10/g$, which suffices to wash out most of the interference while the two Gaussian lobes, which decay only at the rate κ , persist.

Figure 8 shows the inversion, the reduced entropy, and the logarithmic negativity under dissipation. Three effects are visible. (i) The Rabi revival is damped and the inversion relaxes toward $\langle\sigma_z\rangle \rightarrow -1$ as photons leak out; for $\kappa/g = 0.2$ the atom is essentially in its ground state by $gt \approx 30$. (ii) The reduced entropy no longer measures entanglement: for $\kappa/g \geq 0.05$ it saturates near 1 bit at late times while the negativity decays toward zero ($E_{\mathcal{N}} = 0.14, 0.02$, and 0.00 at $gt = 30$ for $\kappa/g = 0.05, 0.10$, and 0.20). (iii) The atom-field entanglement at $t_r/2$ is comparatively robust: $E_{\mathcal{N}}(t_r/2) \approx 0.35$ for all $\kappa/g \leq 0.1$, close to the unitary value 0.36 , even though the entropy minimum is progressively filled in [$S(t_r/2) = 0.14, 0.16, 0.27$, and 0.49 for $\kappa/g = 0, 0.01, 0.05$, and 0.10] and the cat fringes of Table II are already gone at $\kappa/g = 0.05$. Cavity loss thus destroys the internal coherence of the field cat long before it changes the (small) residual entanglement between atom and field.

These results demonstrate the challenge of maintaining quantum coherence. By the operational criterion $\delta > 0.05$ at $t_r/2$, cat-state formation with $\bar{n} = 10$ requires $\kappa/g \lesssim 0.03$, i.e. $g/\kappa \gtrsim 30$; the threshold depends on $\bar{n}$, on the observation time, and on the chosen figure of merit (a full map of δ and F_{cat} over the $(\bar{n}, \kappa/g)$ plane is provided with the code). This condition is routinely achieved in microwave cavity QED [9] and circuit QED [10], but remains an engineering challenge in optical cavities.

V. CLASSICAL VS. QUANTUM LIGHT: CRITERIA FOR QUANTUM ENGINEERING

The computational results above illustrate the distinction between classical and quantum light in a dynamical setting. Here we summarize the formal criteria and their relevance to quantum technologies.

The Glauber–Sudarshan P -distribution [11, 12] provides the gold-standard test. If a state’s density matrix can be written as $\rho = \int P(\alpha)|\alpha\rangle\langle\alpha| d^2\alpha$ with a non-negative, well-behaved $P(\alpha)$, the state is classical (a mixture of coherent states). States for which $P(\alpha)$ is more singular than a delta function or takes negative values, including Fock states and squeezed states, are non-classical. The Wigner function being negative is a sufficient but not necessary indicator; the Husimi Q -function is always positive and thus less sensitive.

Photon statistics provide experimentally accessible tests. Sub-Poissonian statistics (Mandel $Q < 0$) or photon antibunching ($g^{(2)}(0) < 1$) cannot occur with classical fields. The observation of $g^{(2)}(0) \approx 0$ in single-atom fluorescence [32] was the first definitive proof of quantum light. Conversely, photon bunching ($g^{(2)}(0) = 2$) is expected for thermal light and does not imply non-classicality.

Squeezing below the vacuum level is another definitive quantum signature. No classical state can have less noise in any quadrature than the vacuum. The injection of squeezed vacuum into the Advanced LIGO detectors has reduced the quantum shot noise by up to ≈ 3 dB at frequencies above ~ 50 Hz, which was estimated to raise the detection rate for binary-neutron-star mergers by 40–50% [14].

These criteria matter because non-classical light enables quantum technologies that surpass classical limits: squeezed states improve interferometric sensitivity beyond the standard quantum limit; entangled photon pairs enable quantum key distribution [33]; and NOON states can achieve Heisenberg-limited phase sensitivity scaling as $1/N$ rather than the shot-noise $1/\sqrt{N}$.

VI. CONNECTION TO EXPERIMENT

The collapse-revival phenomenon remained a theoretical prediction for over two decades after the original JC model [5] and the analysis by Eberly *et al.* [6]. Its observation required isolating a single two-level atom in a single-mode cavity in the strong-coupling regime ($g \gg \kappa, \gamma$), a formidable experimental challenge.

Microwave cavity QED (1987). The first observation of quantum revivals was achieved by Rempe, Walther, and Klein [8] using a one-atom maser: Rydberg atoms (with transition frequencies in the microwave range) passed one at a time through a high- Q superconducting niobium cavity. The cavity field was prepared in a coherent or thermal state, the atom entered in $|e\rangle$, and after a controlled interaction time its state was measured by field ionization. The measured $P_e(t)$ exhibited the predicted collapse of Rabi oscillations followed by a revival, in quantitative agreement with JC theory. In subsequent experiments by Haroche, Raimond, and collaborators [9], the same microwave cavity QED platform was used to observe quantum jumps of the photon number, generate and detect Schrödinger cat states [7], and perform

quantum non-demolition measurements of the intracavity photon number. Typical parameters in these experiments are $g/2\pi \sim 50$ kHz with $\kappa/2\pi \sim 1$ Hz, giving $g/\kappa \sim 10^4$, well within the strong-coupling regime.

Trapped-ion analog (1996). The JC Hamiltonian also describes a trapped ion whose internal (electronic) levels are coupled to quantized motional (vibrational) levels by a laser tuned to a motional sideband. In 1996, Meekhof *et al.* [36] (Wineland’s group at NIST) observed collapse and revival of Rabi oscillations in a single trapped $^9\text{Be}^+$ ion. The ion’s vibrational mode played the role of the cavity field, and by preparing it in a coherent state of motion and driving the blue sideband, they reproduced the full collapse-revival sequence. This demonstration showed that the physics is universal to any two-level system coupled to a quantized bosonic mode, not specific to cavity photons.

Optical cavities (1992 onwards). In the optical domain, Thompson, Rempe, and Kimble [19] observed vacuum Rabi splitting in the transmission spectrum of a Fabry–Pérot cavity containing single cesium atoms, confirming the dressed-state structure at $n = 1$. Full time-domain collapse-revival experiments in optical cavities are more challenging due to shorter cavity lifetimes, but the spectroscopic signatures of the quantized JC interaction have been well established.

Circuit QED (2000s onwards). In circuit QED, a superconducting qubit (typically a transmon) is coupled to a microwave resonator on a chip [10]. Typical parameters are $g/2\pi \sim 100\text{--}300$ MHz and $\kappa/2\pi \sim 1$ MHz, giving $g/\kappa \sim 100\text{--}300$. This satisfies the $g/\kappa \gtrsim 30$ threshold identified in Sec. IV C for observing cat-state formation with $\bar{n} = 10$. Hofheinz *et al.* [34] generated arbitrary Fock states and their superpositions in a coplanar waveguide resonator coupled to a phase qubit, verifying the $\sqrt{n+1}$ scaling of the Rabi frequency. Ofek *et al.* [35] used the JC interaction to implement quantum error correction with cat states encoded in a superconducting cavity, reaching the break-even point where error correction extends the logical qubit lifetime beyond the physical components.

Our dimensionless simulation results (with $g = 1$) can be directly mapped to any of these platforms by substituting the physical value of g . For the microwave cavity QED parameters above ($g = 2\pi \times 50$ kHz), the revival time for $\bar{n} = 10$ is $t_r = 2\pi\sqrt{10}/g \approx 60$ μs , much shorter than the cavity lifetime $1/\kappa \approx 0.16$ s. For circuit QED with $g/2\pi = 100\text{--}300$ MHz, $t_r \approx 10\text{--}30$ ns, compared with a cavity lifetime $1/\kappa \approx 160$ ns for $\kappa/2\pi = 1$ MHz. In both cases the first revival occurs well within the coherence time of the system; the cat-state fringes, which decohere at the enhanced rate $2\kappa\bar{n}$, survive only where $\kappa/g \lesssim 0.01\text{--}0.03$, as in the microwave and circuit-QED regimes.

VII. CONCLUSIONS

We have presented a computational study of quantum light–matter interaction in the Jaynes–Cummings model that goes beyond standard textbook treatments in two respects. First, time-resolved Wigner function snapshots directly visualize the formation of a Schrödinger cat state at half the revival time, with Wigner negativity volume $\delta = 0.85$, purity 0.96, and cat fidelity 0.78. The cat coincides with a *minimum* of atom–field entanglement ($S \approx 0.14$ bit), so it is a nearly pure state of the field alone, whereas during the preceding collapse the atom and field are maximally entangled and the reduced field is a two-branch mixture. The interference fringes in the Wigner function provide a quantitative, phase-space signature of non-classicality that is destroyed by cavity decay rates as low as $\kappa/g = 0.02$, while the residual atom–field negativity at $t_r/2$ is much less affected. Second, a systematic comparison across coherent, thermal, squeezed, and Fock initial fields demonstrates that the half-revival disentanglement first analyzed by Gea-Banacloche [27] and by Phoenix and Knight [24, 29] is unique to the coherent state, and that for a thermal field a saturated reduced entropy of one bit conceals a logarithmic negativity of only ≈ 0.2 : once the global state is mixed, the reduced entropy must not be read as an entanglement measure.

These results illustrate how the quantum statistics of the field, encoded in the photon-number distribution, determine the entanglement dynamics and the fragility of non-classical features under decoherence. The computational framework is straightforward to extend to multi-mode cavities, ultrastrong coupling regimes, and quantum trajectory simulations of individual measurement records.




FIG. 1. Atomic inversion $\langle \sigma_z(t) \rangle$ for a two-level atom interacting with a coherent-state field ($\bar{n} = 10$) in the resonant Jaynes–Cummings model. Colored markers indicate the six snapshot times used for the Wigner function evolution in Fig. 2. The collapse of oscillations near $t \approx 2/g$ and the first revival at $t_r \approx 19.9/g$ are clearly visible.

ACKNOWLEDGMENTS

The author thanks Prof. Luca Dal Negro for guidance and feedback during the preparation of this work (EC 585 and EC 777, Boston University).

-
- [1] I. I. Rabi, Phys. Rev. **51**, 652 (1937).
 - [2] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, 1997).
 - [3] L. Allen and J. H. Eberly, *Optical Resonance and Two-Level Atoms* (Dover, 1975).
 - [4] C. C. Gerry and P. L. Knight, *Introductory Quantum Optics* (Cambridge University Press, 2004).
 - [5] E. T. Jaynes and F. W. Cummings, Proc. IEEE **51**, 89 (1963).
 - [6] J. H. Eberly, N. B. Narozhny, and J. J. Sanchez-Mondragon, Phys. Rev. Lett. **44**, 1323 (1980).
 - [7] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **77**, 4887 (1996).
 - [8] G. Rempe, H. Walther, and N. Klein, Phys. Rev. Lett. **58**, 353 (1987).
 - [9] S. Haroche and J.-M. Raimond, *Exploring the Quantum: Atoms, Cavities, and Photons* (Oxford University Press, 2006).
 - [10] A. Blais, A. L. Grimsmo, S. M. Girvin, and A. Wallraff, Rev. Mod. Phys. **93**, 025005 (2021).
 - [11] R. J. Glauber, Phys. Rev. **131**, 2766 (1963).
 - [12] E. C. G. Sudarshan, Phys. Rev. Lett. **10**, 277 (1963).
 - [13] J. Aasi *et al.* (LIGO Scientific Collaboration), Nat. Photonics **7**, 613 (2013).
 - [14] M. Tse *et al.* (LIGO Scientific Collaboration), Phys. Rev. Lett. **123**, 231107 (2019).
 - [15] E. Wigner, Phys. Rev. **40**, 749 (1932).
 - [16] R. L. Hudson, Rep. Math. Phys. **6**, 249 (1974).
 - [17] A. Kenfack and K. Życzkowski, J. Opt. B: Quantum Semiclass. Opt. **6**, 396 (2004).
 - [18] B. R. Mollow, Phys. Rev. **188**, 1969 (1969).
 - [19] R. J. Thompson, G. Rempe, and H. J. Kimble, Phys. Rev. Lett. **68**, 1132 (1992).
 - [20] G. Lindblad, Commun. Math. Phys. **48**, 119 (1976).
 - [21] H. M. Wiseman and G. J. Milburn, *Quantum Measurement and Control* (Cambridge University Press, 2009).
 - [22] W. H. Zurek, Rev. Mod. Phys. **75**, 715 (2003).
 - [23] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000).
 - [24] S. J. D. Phoenix and P. L. Knight, Ann. Phys. (N.Y.) **186**, 381 (1988).
 - [25] J. R. Johansson, P. D. Nation, and F. Nori, Comp. Phys. Comm. **183**, 1760 (2012).
 - [26] J. R. Johansson, P. D. Nation, and F. Nori, Comp. Phys. Comm. **184**, 1234 (2013).
 - [27] J. Gea-Banacloche, Phys. Rev. Lett. **65**, 3385 (1990).
 - [28] J. Gea-Banacloche, Phys. Rev. A **44**, 5913 (1991).
 - [29] S. J. D. Phoenix and P. L. Knight, Phys. Rev. A **44**, 6023 (1991).
 - [30] G. Vidal and R. F. Werner, Phys. Rev. A **65**, 032314 (2002).
 - [31] M. B. Plenio, Phys. Rev. Lett. **95**, 090503 (2005).
 - [32] H. J. Kimble, M. Dagenais, and L. Mandel, Phys. Rev. Lett. **39**, 691 (1977).
 - [33] A. K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
 - [34] M. Hofheinz, H. Wang, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O’Connell, D. Sank, J. Wenner, J. M. Martinis, and A. N. Cleland, Nature **459**, 546 (2009).
 - [35] P. Ofek *et al.*, Nature **536**, 441 (2016).
 - [36] D. M. Meekhof, C. Monroe, B. E. King, W. M. Itano, and D. J. Wineland, Phys. Rev. Lett. **76**, 1796 (1996).
 - [37] D. F. Walls and G. J. Milburn, *Quantum Optics*, 2nd ed. (Springer, 2008).
 - [38] N. K. Nguyen, *QSOL_CQED*, version 1.1.0, Zenodo (2026), <https://doi.org/10.5281/zenodo.22806076>.

Wigner Function Evolution of Cavity Field
($\bar{n} = 10, g = 1.0$, resonant)




FIG. 2. Wigner function $W(x, p)$ of the reduced cavity field state at six characteristic times during Jaynes–Cummings evolution ($\bar{n} = 10, \Delta = 0$). At $t = 0$, the field is a coherent-state Gaussian. By $t = 2 t_c$ (collapse onset), the distribution begins to spread. At $t = t_r/2$, a Schrödinger cat state has formed: two distinct lobes at $p \approx \pm \sqrt{2\bar{n}}$ connected by interference fringes with strong Wigner negativity ($\delta = 0.85$); at this time the atom and field are approximately disentangled (Sec. IV B). By $t = t_r$, the field partially re-localizes at revival ($\delta = 0.31$).





FIG. 4. Decoherence of the Schrödinger cat state. Wigner function at $t = t_r/2$ for increasing cavity decay rates $\kappa/g = 0, 0.02, 0.05, 0.10$ (left to right). The interference fringes are progressively destroyed; quantitative values of the Wigner negativity δ are given in Table II.




FIG. 5. Comparison of Jaynes–Cummings dynamics for four initial field states with $\bar{n} = 10$. (a) Atomic inversion $\langle \sigma_z \rangle$ for the coherent (blue), thermal (red), and squeezed-vacuum (green) fields. (b) Fock state $|10\rangle$ over a short window: the inversion oscillates with period $\pi/(g\sqrt{11})$ and the entropy (plotted as $2S - 1$) with half that period. (c) Reduced-atom entropy $S(\rho_{\text{atom}})$ (solid) and logarithmic negativity $E_{\mathcal{N}}$ (dashed). The coherent state collapses to $S \approx 1$ bit and disentangles at $t_r/2$ (dotted line); the thermal state saturates at $S = 1$ bit but has $E_{\mathcal{N}} \approx 0.2$ because its global state is mixed; the squeezed vacuum shows only partial dips. The dashed purple line marks the revival time t_r .




FIG. 6. Detailed coherent-state dynamics ($\bar{n} = 10$). Top: atomic inversion. Middle: entanglement entropy $S(\rho_{\text{atom}})$, peaking near 1 bit during the collapse and reaching its minimum (≈ 0.14 bit) at $t_r/2$, where the atom and field disentangle and the field is the cat state of Fig. 3. Bottom: field purity $\mathcal{P} = \text{Tr}[\rho_{\text{field}}^2]$, dropping to ≈ 0.5 during the collapse (a two-branch mixture, the field being entangled with the atom) and rising to 0.96 at $t_r/2$ (nearly pure cat). Dotted and dashed lines mark $t_r/2$ and t_r .




FIG. 7. Entanglement entropy $S(\rho_{\text{atom}})$ for coherent-state fields with $\bar{n} = 1, 4, 9, 16, 25, 36$. As $\bar{n}$ increases, the collapse time remains $\sim 1/g$ while the half-revival time $\pi\sqrt{\bar{n}}/g$ (dotted) and the revival time $t_r = 2\pi\sqrt{\bar{n}}/g$ (dashed) grow as $\sqrt{\bar{n}}$. The entropy minimum at $t_r/2$ (atom-field disentanglement) deepens with $\bar{n}$ and is cleanly resolved for $\bar{n} \gtrsim 9$.




FIG. 8. Effect of cavity decay on JC dynamics ($\bar{n} = 10$). Top: atomic inversion for $\kappa/g = 0-0.20$. Middle: reduced-atom entropy, which for $\kappa > 0$ measures total (classical plus quantum) correlations. Bottom: logarithmic negativity $E_{\mathcal{N}}$, a proper entanglement monotone for the mixed states produced by dissipation. Increasing κ damps the revival, drives the atom toward its ground state, and suppresses the late-time negativity even where the reduced entropy stays near 1 bit. Dotted and dashed lines mark $t_r/2$ and t_r .




FIG. 9. Atomic inversion $\sum_n p(n) \cos(2g\sqrt{n+1}t)$ for coherent (left) and thermal (right) field states at $\bar{n} = 4, 9, 14, 19, 24$; time in units of $1/g$. Coherent states show collapse-revival dynamics whose revivals, at multiples of $t_r = 2\pi\sqrt{\bar{n}}/g$, become more widely separated and better resolved as $\bar{n}$ increases; thermal states dephase within a Rabi period and show no resolved revival.

Wigner functions of Fock, squeezed-vacuum and coherent states




FIG. 10. Static Wigner functions $W(x, p)$ for Fock states ($|0\rangle$, $|1\rangle$, $|6\rangle$, $|7\rangle$), squeezed vacuum ($r = 1.0$, $\phi = 0$ and $\pi/2$), and coherent states ($\alpha = 2$ and $2 + 2i$). Fock states exhibit ring-like structures and negativity; squeezed states show elliptical distributions (squeezing axis at angle $\phi/2$); coherent states are displaced Gaussians with no negativity. Phase-space convention $a = (x + ip)/\sqrt{2}$.




FIG. 11. Contour plots of the Wigner functions from Fig. 10. The color scale highlights negative regions (blue) as signatures of non-classicality.




FIG. 12. Mollow triplet spectra for a driven two-level system at varying Rabi frequencies $\Omega/\gamma = 0.5$ –15. As the driving strength increases, the single peak splits into three: a central Rayleigh line and two sidebands at $\omega - \omega_L = \pm\Omega$.