


New Approach of Numerical Relativity

Implementation, tests and applications



Alan Tsz-Lok Lam

Supervisor: Prof. Tjonnie G.F. LI

Department of Physics
The Chinese University of Hong Kong

This thesis is submitted for the degree of
Master of Philosophy in Physics

July 2021

I would like to dedicate this thesis to my loving parents . . .

Declaration

I hereby declare that except where specific reference is made to the work of others, the contents of this dissertation are original and have not been submitted in whole or in part for consideration for any other degree or qualification in this, or any other university. This dissertation is my own work and contains nothing which is the outcome of work done in collaboration with others, except as specified in the text and Acknowledgements. This dissertation contains fewer than 65,000 words including appendices, bibliography, footnotes, tables and equations and has fewer than 150 figures.

Alan Tsz-Lok Lam

July 2021

Acknowledgements

And I would like to acknowledge ...

Abstract

This is where you write your abstract ...

Table of contents

List of figures	xiii
List of tables	xv
1 Introduction	1
1.1 Introduction to General Relativity	1
1.1.1 Einstein Field Equation	1
1.1.2 Relativistic Star	1
1.1.3 Gravitational Wave	1
1.2 Gmunu: A general-relativistic electro-magneto-hydrodynamics code for generic astrophysical simulations	1
1.3 Outline of this thesis	1
1.4 Units and Convention	1
2 Formulations of Einstein Field Equations	3
2.1 Introduction	3
2.2 The 3+1 decomposition of spacetime	3
2.2.1 Foliation of spacetime	3
2.2.2 Derivative operator	5
2.2.3 Extrinsic curvature	6
2.2.4 The Gauss, Codazzi and Ricci equations	7
2.2.5 Constraint and evolution equations	7
2.2.6 The Arnowitt, Deser and Misner equations	9
2.3 Conformal Decomposition	9
2.3.1 Conformal transformation of the spatial metric	9
2.3.2 Conformal transformation of the extrinsic curvature	10
2.3.3 Conformal transverse-traceless decomposition	13
2.4 Gauge Condition	14

2.4.1	Maximal Slicing	14
2.4.2	Generalized Dirac Gauge	14
2.5	Constrained scheme for the Einstein equations	14
2.5.1	The Conformal Flatness Condition	14
2.5.2	The Fully Constrained Formulation	14
3	Formulations of the Relativistic Hydrodynamics	15
3.1	The 3+1 "Valencia" formulation	15
3.2	The reference-metric formulism	15
3.3	Conserved to Primitive variables conversion	15
4	NUMERICAL Methods and Tests	17
4.1	Multigrid Method for elliptic equations	17
4.1.1	Overview	17
4.1.2	Smoother	17
4.1.3	Restriction and Prolongation	17
4.1.4	Multigrid cycle	17
4.1.5	Grid Structure	17
4.1.6	Ghost cells and refinement boundary	17
4.1.7	Numerical Tests	17
4.2	Numerical method for hydrodynamics	17
4.2.1	Finite volume methods	17
4.2.2	Time Discretization	17
4.2.3	Atmosphere Treatment	17
4.2.4	Numerical Tests	17
	References	19
	Appendix A Useful relations for implementation of constrained scheme	21
A.1	The elliptic equations in constrained scheme	21
A.2	Generalized Dirac gauge conditions	21
	Appendix B Reference flat metric in 3D	23

List of figures

List of tables

Chapter 1

Introduction

1.1 Introduction to General Relativity

1.1.1 Einstein Field Equation

$$G_{\mu\nu} = 8\pi T_{\mu\nu} \tag{1.1}$$

1.1.2 Relativistic Star

1.1.3 Gravitational Wave

1.2 Gmunu: A general-relativistic electro-magneto-hydrodynam code for generic astrophysical simulations

1.3 Outline of this thesis

1.4 Units and Convention

Chapter 2

Formulations of Einstein Field Equations

2.1 Introduction

Due to the complexity and nonlinearity of Einstein field equations, it is extremely difficult to obtain analytical solution even for the simplest dynamical evolution systems. Therefore, the accurate discription of the such systems can only be derived through numerical simulation. For this, we need to reformulate the Einstein equations as an initial-value problem or Cauchy problem. In this chapter, we will introduction the Arnowitt-Deser-Misner (ADM) formulation, which is the foundation of the 3+1 numerical relativity. In particular, we will focus on the constrained scheme for the Einstein equations.

2.2 The 3+1 decomposition of spacetime

2.2.1 Foliation of spacetime

In the 3+1 decomposition, the spacetime manifold \mathcal{M} is foliated into a set of non-intersecting spacelike hypersurfaces Σ_t parameterized by the coordinate time t [5]. We denote a future-directed timelike unit four-vector n^μ normal to the hypersurface Σ_t (i.e. $n_\mu \propto \nabla_\mu t$). The induced spacetime metric $\gamma_{\mu\nu}$ on each hypersurface can then be defined as

$$\gamma_{\mu\nu} := g_{\mu\nu} + n_\mu n_\nu. \quad (2.1)$$

Thus, we can construct spatial projection tensor $\gamma^\mu{}_\nu$ and time projection tensor $N^\mu{}_\nu$ as

$$\gamma^\mu{}_\nu := \delta^\mu{}_\nu + n^\mu n_\nu, \quad N^\mu{}_\nu := -n^\mu n_\nu, \quad (2.2)$$

which decompose any generic four-vector U^μ into spatial part $\gamma^\mu{}_\nu U^\nu$ and timelike part $N^\mu{}_\nu U^\nu$. Therefore, we can decompose the timelike vector field

$$t^\mu = \alpha n^\mu + \beta^\mu \quad (2.3)$$

into two components as

$$\alpha := -t^\mu n_\mu, \quad \beta^\mu := t^\nu \gamma^\mu{}_\nu, \quad (2.4)$$

where the lapse function α measures the physical proper time ($\alpha \Delta t$) between two neighboring spatial hypersurface Σ_t and $\Sigma_{t+\Delta t}$, and the shift vector β^i measures the changes of spatial coordinates on $\Sigma_{t+\Delta t}$.

Here, we summarise several useful relations. The timelike normal vector n^μ and its corresponding one-form n_μ can be expressed as

$$n^\mu = \frac{1}{\alpha} (1, \beta^i), \quad n_\mu = (\alpha, \vec{0}). \quad (2.5)$$

The generic line element in 3+1 decomposition is given by

$$ds^2 = -(\alpha^2 - \beta^i \beta_i) dt^2 + \beta_i dx^i dt + \gamma_{ij} dx^i dx^j \quad (2.6)$$

The covariant and contravariant components of the metric can be written as

$$g_{\mu\nu} = \begin{pmatrix} -\alpha^2 + \beta^i \beta_i & \beta_j \\ \beta_i & \gamma_{ij} \end{pmatrix}, \quad g^{\mu\nu} = \begin{pmatrix} -\frac{1}{\alpha^2} & \beta^j \\ \beta^i & \gamma^{ij} \end{pmatrix}. \quad (2.7)$$

From equation(2.7), we can conclude that

$$\sqrt{-g} = \alpha \sqrt{\gamma}, \quad (2.8)$$

where $g := \det(g_{\mu\nu})$ and $\gamma := \det(\gamma_{ij})$.

2.2.2 Derivative operator

With the 3+1 decomposition, we can now construct the 3-dimensional covariant derivative D_α associated with $\gamma_{\mu\nu}$ by projecting the 4-dimensional covariant derivative ∇_α onto Σ_t , which is given by

$$D_\alpha T^{\mu_1\mu_2\cdots}_{\nu_1\nu_2\cdots} = \gamma_\alpha^\beta \gamma_{\rho_1}^{\mu_1} \gamma_{\rho_2}^{\mu_2} \cdots \gamma_{\nu_1}^{\sigma_1} \gamma_{\nu_2}^{\sigma_2} \cdots \nabla_\beta T^{\rho_1\rho_2\cdots}_{\sigma_1\sigma_2\cdots}, \quad (2.9)$$

for arbitrary tensor $T^{\mu_1\mu_2\cdots}_{\nu_1\nu_2\cdots}$ on spatial hypersurface Σ_t . Using equation(2.9), it can be shown that the covariant derivative of $\gamma_{\mu\nu}$ vanishes

$$\begin{aligned} D_\alpha \gamma_{\mu\nu} &= \gamma_\alpha^\beta \gamma_\rho^\mu \gamma_\nu^\sigma \nabla_\beta (g_{\rho\sigma} + n_\rho n_\sigma) \\ &= \gamma_\alpha^\beta \gamma_\rho^\mu \gamma_\nu^\sigma (n_\rho \nabla_\beta n_\sigma + n_\sigma \nabla_\beta n_\rho) = 0 \end{aligned} \quad (2.10)$$

The components of 3-dimensional connection coefficients $\Gamma^\alpha_{\mu\nu}$ in coordinate basis can be expressed as

$$\Gamma^\alpha_{\mu\nu} = \frac{1}{2} \gamma^{\alpha\beta} (\partial_\nu \gamma_{\beta\mu} + \partial_\mu \gamma_{\beta\nu} - \partial_\beta \gamma_{\mu\nu}). \quad (2.11)$$

Here, the upper left index ⁽⁴⁾ marks the 4-dimensional tensors while the unmarked one represents purely spatial 3-dimensional tensors. Similarly, the 3-dimensional Riemann tensor $R^\alpha_{\beta\mu\nu}$ associated with $\gamma_{\mu\nu}$ is defined by requiring that

$$2D_{[\nu} D_{\mu]} W_\beta = W_\alpha R^\alpha_{\beta\mu\nu}, \quad R^\alpha_{\beta\mu\nu} n_\alpha = 0, \quad (2.12)$$

which can be explicitly expressed in coordinate basis as

$$R^\alpha_{\beta\mu\nu} = \partial_\mu \Gamma^\alpha_{\beta\nu} - \partial_\nu \Gamma^\alpha_{\beta\mu} + \Gamma^\alpha_{\mu\rho} \Gamma^\rho_{\beta\nu} - \Gamma^\alpha_{\nu\rho} \Gamma^\rho_{\beta\mu}. \quad (2.13)$$

The 3-dimensional Ricci tensor $R_{\mu\nu}$ and Ricci scalar R are defined in a similar manner as their 4-dimensional counterparts

$$R_{\mu\nu} := R^\alpha_{\mu\alpha\nu}, \quad R := R^\mu_{\mu}. \quad (2.14)$$

Since $R^\alpha_{\beta\mu\nu}$ is purely spatial and can be computed by the spatial derivatives of the spatial metric alone, it only contains about information about the curvature intrinsic to the hypersurface Σ_t , but cannot contain all the information of ⁽⁴⁾ $R^\alpha_{\beta\mu\nu}$ which includes time derivative of the 4-dimensional metric. The missing information can be found in a purely spatial symmetric tensor called the extrinsic curvature $K_{\mu\nu}$.

2.2.3 Extrinsic curvature

The extrinsic curvature $K_{\mu\nu}$ is related to the time derivative of the spatial metric $\gamma_{\mu\nu}$. Therefore, the spatial metric and extrinsic curvature $(\gamma_{\mu\nu}, K_{\mu\nu})$ are equivalent to the positions and velocities in classical mechanics, which describe the instantaneous state of the gravitational field. It can be obtained by projecting of the gradient of the normal vector $\gamma_\mu^\lambda \gamma_\nu^\rho \nabla_\lambda n_\rho$ into the hypersurface Σ_t , and then taking the negative expression of the symmetric part

$$\begin{aligned} K_{\mu\nu} &:= -\gamma_\mu^\lambda \gamma_\nu^\rho \nabla_\lambda n_\rho \\ &= -\gamma_\mu^\lambda (\delta_\nu^\rho + n_\nu n^\rho) \nabla_\lambda n_\rho \\ &= -\gamma_\mu^\lambda \nabla_\lambda n_\nu, \end{aligned} \tag{2.15}$$

where the identity $n^\rho \nabla_\lambda n_\rho = 0$ is used.

We can also define an spatial acceleration a_ν

$$a_\nu := n^\mu \nabla_\mu n_\nu, \tag{2.16}$$

satisfying the identities

$$a_\nu = D_\nu \ln \alpha, \tag{2.17}$$

to rewrite equation(2.15) as

$$K_{\mu\nu} = -\nabla_\mu n_\nu - n_\mu a_\nu \tag{2.18}$$

Finally, we can write the extrinsic curvature $K_{\mu\nu}$ as the Lie derivative of the spatial metric along the local normal n^μ

$$K_{\mu\nu} = -\frac{1}{2} \mathcal{L}_n \gamma_{\mu\nu}. \tag{2.19}$$

Using equation(2.3), we can express the Lie derivative \mathcal{L}_n as

$$\mathcal{L}_n = \frac{1}{\alpha} (\mathcal{L}_t - \mathcal{L}_\beta), \tag{2.20}$$

and thus obtain the evolution equation for the spatial metric

$$\mathcal{L}_t \gamma_{\mu\nu} = -2\alpha K_{\mu\nu} + \mathcal{L}_\beta \gamma_{\mu\nu}. \tag{2.21}$$

2.2.4 The Gauss, Codazzi and Ricci equations

To express the Einstein field equations in term of the spatial variables $(\gamma_{\mu\nu}, K_{\mu\nu})$ we defined previous, we first have to relate 3-dimensional Riemann tensor $R^\alpha_{\beta\mu\nu}$ on Σ_t to the 4-dimensional Riemann tensor ${}^{(4)}R^\alpha_{\beta\mu\nu}$ on \mathcal{M} , The relation between $R^\alpha_{\beta\mu\nu}$ and the full spatial projection of ${}^{(4)}R^\alpha_{\beta\mu\nu}$ is given by the Gauss' equation

$$R_{\alpha\beta\mu\nu} + K_{\alpha\mu}K_{\beta\nu} - K_{\alpha\nu}K_{\beta\mu} = \gamma_\alpha{}^\rho\gamma_\beta{}^\sigma\gamma_\mu{}^\lambda\gamma_\nu{}^\delta{}^{(4)}R_{\rho\sigma\lambda\delta}, \quad (2.22)$$

while the projection of ${}^{(4)}R^\alpha_{\beta\mu\nu}$ with one index projected in the normal direction is given by the Codazzi equation

$$D_\nu K_{\mu\alpha} - D_\mu K_{\nu\alpha} = \gamma_\mu{}^\rho\gamma_\nu{}^\sigma\gamma_\alpha{}^\lambda n^\delta{}^{(4)}R_{\rho\sigma\lambda\delta}. \quad (2.23)$$

Finally, by projecting two indices of ${}^{(4)}R_{\rho\sigma\lambda\delta}$ in the normal direction, we can relate it to the time derivative of $K_{\mu\nu}$

$$\mathcal{L}_n K_{\mu\nu} = n^\alpha n^\beta \gamma_\mu{}^\lambda \gamma_\nu{}^\delta{}^{(4)}R_{\alpha\beta\lambda\delta} - \frac{1}{\alpha} D_\mu D_\nu \alpha - K_\nu{}^\lambda K_{\mu\lambda}, \quad (2.24)$$

which is called the Ricci equation.

2.2.5 Constraint and evolution equations

Using the Gauss, Codazzi and Ricci equations, the Einstein fields equations can be decomposed into a set of evolution equations and a set of constraint equations of $(\gamma_{\mu\nu}, K_{\mu\nu})$. To begin with, we define the following matter quantities

$$S_{\mu\nu} := \gamma^\alpha{}_\mu \gamma^\beta{}_\nu T_{\alpha\beta}, \quad (2.25)$$

$$S_\mu := -\gamma^\alpha{}_\mu n^\beta T_{\alpha\beta}, \quad (2.26)$$

$$S := S^\mu{}_\mu, \quad (2.27)$$

$$E := n^\alpha n^\beta T_{\alpha\beta}, \quad (2.28)$$

which decompose the stress-energy tensor as

$$T_{\mu\nu} = E n_\mu n_\nu + S_\mu n_\nu + S_\nu n_\mu + S_{\mu\nu}. \quad (2.29)$$

By contracting the α, μ indices in equation(2.22), we can obtain

$$R_{\mu\nu} = \gamma_\mu^\alpha \gamma_\nu^\beta \left({}^{(4)}R_{\alpha\beta} + n^\rho n^\sigma {}^{(4)}R_{\alpha\rho\beta\sigma} \right) + K_{\mu\lambda} K_\nu^\lambda - K_{\mu\nu} K, \quad (2.30)$$

where $K := K^\mu_\mu$ is the trace of the extrinsic curvature. Further contracting the μ, ν indices in equation(2.30), the contracted Gauss' equation becomes

$$2n^\mu n^\nu G_{\mu\nu} = R + K^2 - K_{\mu\nu} K^{\mu\nu}, \quad (2.31)$$

Using the Einstein equation(1.1), we can obtain the Hamiltonian constraint

$$R + K^2 - K_{\mu\nu} K^{\mu\nu} = 16\pi E. \quad (2.32)$$

Similarly, by contracting α, ν indices in equation(2.23), the Codazzi equation yields

$$D_\nu K_\mu^\nu - D_\mu K = -8\pi \gamma_\mu^\alpha n^\beta T_{\alpha\beta}, \quad (2.33)$$

and thus obtain the momentum constrain equation

$$D_\nu K_\mu^\nu - D_\mu K = 8\pi S_\mu. \quad (2.34)$$

The Ricci equation (2.24) can be rewritten using equation(2.30) to

$$\mathcal{L}_n K_{\mu\nu} = R_{\mu\nu} - \gamma_\mu^\rho \gamma_\nu^\sigma {}^{(4)}R_{\rho\sigma} - 2K_{\mu\lambda} K_\nu^\lambda + K K_{\mu\nu} - \frac{1}{\alpha} D_\mu D_\nu \alpha. \quad (2.35)$$

Using the Einstein equations (1.1)

$$\begin{aligned} \gamma_\mu^\rho \gamma_\nu^\sigma {}^{(4)}R_{\rho\sigma} &= 8\pi \gamma_\mu^\rho \gamma_\nu^\sigma \left(T_{\rho\sigma} - \frac{1}{2} g_{\rho\sigma} T^\mu_\mu \right) \\ &= 8\pi \left[S_{\mu\nu} - \frac{1}{2} \gamma_{\mu\nu} (S - E) \right] \end{aligned} \quad (2.36)$$

and equation(2.20), we can finally obtain the evolution equation for $K_{\mu\nu}$ as

$$\mathcal{L}_t K_{\mu\nu} = -D_\mu D_\nu \alpha + \alpha \left(R_{\mu\nu} - 2K_{\mu\lambda} K_\nu^\lambda + K K_{\mu\nu} \right) - 8\pi \alpha \left[S_{\mu\nu} - \frac{1}{2} \gamma_{\mu\nu} (S - E) \right] + \mathcal{L}_\beta K_{\mu\nu}. \quad (2.37)$$

2.2.6 The Arnowitt, Deser and Misner equations

The Lie derivative in the evolution equations (2.21) and (2.37) can be expressed in terms of coordinate basis as

$$\mathcal{L}_t K_{\mu\nu} = \partial_t K_{\mu\nu} \quad (2.38)$$

$$\mathcal{L}_\beta K_{\mu\nu} = \beta^\lambda D_\lambda K_{\mu\nu} + K_{\mu\lambda} D_\nu \beta^\lambda + K_{\lambda\nu} D_\mu \beta^\lambda \quad (2.39)$$

As the result, the Einstein field equations (1.1) in the standard 3+1 decomposition can be decomposed into a set of constraint equations and evolution equation of (γ_{ij}, K_{ij}) in terms of coordinate basis, which are referred to as the Arnowitt, Deser and Misner (ADM) equations [1, 12]

$$R + K^2 - K_{ij} K^{ij} = 16\pi E, \quad (\text{Hamiltonian constraint}) \quad (2.40)$$

$$D_j (K^{ij} - \gamma^{ij} K) = 8\pi S^i, \quad (\text{momentum constraint}) \quad (2.41)$$

$$\partial_t \gamma_{ij} = -2\alpha K_{ij} + D_i \beta_j + D_j \beta_i, \quad (\text{spatial metric evolution}) \quad (2.42)$$

$$\begin{aligned} \partial_t K_{ij} = & -D_i D_j \alpha + \alpha (R_{ij} - 2K_{ik} K_j^k + K K_{ij}) \\ & - 8\pi \alpha \left[S_{ij} - \frac{1}{2} \gamma_{ij} (S - E) \right] \quad (\text{extrinsic curvature evolution}) \\ & + \beta^k D_k K_{ij} + K_{ik} D_j \beta^k + K_{kj} D_i \beta^k. \end{aligned} \quad (2.43)$$

2.3 Conformal Decomposition

The conformal decomposition factors out a scalar component from a spatial metric. It was first developed for initial data problems in general relativity [4, 10, 11, 13], and then used in reformulating evolution equations in the 3+1 formulation. In this section, we will discuss the conformal decomposition of spatial metric and extrinsic curvature in numerical relativity.

2.3.1 Conformal transformation of the spatial metric

We consider the conformal transformation of the spatial metric γ_{ij} as

$$\tilde{\gamma}_{ij} = \psi^{-4} \gamma_{ij}, \quad (2.44)$$

where $\tilde{\gamma}_{ij}$ is the *conformal metric* and ψ is a positive scaling factor satisfying

$$\psi := \det \left(\frac{\gamma}{f} \right), \quad \gamma := \det (\gamma_{ij}), \quad f := \det (f_{ij}) \quad (2.45)$$

for a time independent flat metric f_{ij} (i.e. $\det (\tilde{\gamma}_{ij}) = f$ by construction).

Thus, the *inverse conformal metric* is given by

$$\tilde{\gamma}^{ij} := \psi^4 \gamma^{ij}. \quad (2.46)$$

Substituting the conformal transformation (2.44) into equation(2.11), we can obtain the transformation law for 3-dimensional connection coefficient

$$\Gamma^i_{jk} = \tilde{\Gamma}^i_{jk} + 2 \left(\delta^i_j \tilde{D}_k \ln \psi + \delta^i_k \tilde{D}_j \ln \psi - \tilde{\gamma}_{jk} \tilde{\gamma}^{il} \tilde{D}_l \ln \psi \right). \quad (2.47)$$

From now on, we denote all objects associated with the conformal metric $\tilde{\gamma}^{ij}$ with a tilde symbol. Similarly, the transformation for Ricci tensor and scalar curvature are given by

$$\begin{aligned} R_{ij} = & \tilde{R}_{ij} - 2 \left(\tilde{D}_i \tilde{D}_j \ln \psi + \tilde{\gamma}_{ij} \tilde{\gamma}^{lm} \tilde{D}_l \tilde{D}_m \ln \psi \right) \\ & + 4 \left(\left(\tilde{D}_i \ln \psi \right) \left(\tilde{D}_j \ln \psi \right) - \tilde{\gamma}_{ij} \tilde{\gamma}^{lm} \tilde{D}_l \left(\tilde{D}_m \ln \psi \right) \right) \end{aligned} \quad (2.48)$$

$$R = \psi^{-4} \tilde{R} - 8 \psi^{-5} \tilde{D}^2 \psi, \quad (2.49)$$

where $\tilde{D}^2 = \tilde{\gamma}^{ij} \tilde{D}_i \tilde{D}_j$ denotes the Laplace operator associated with $\tilde{\gamma}_{ij}$. Therefore, using equation(2.49), the Hamiltonian constraint (2.40) becomes

$$8 \tilde{D}^2 \psi - \psi \tilde{R} - \psi^5 K^2 + \psi^2 K_{ij} K^{ij} = -16 \pi \psi^5 E. \quad (2.50)$$

2.3.2 Conformal transformation of the extrinsic curvature

Traceless decomposition

Before we perform the conformal transformation to the extrinsic curvature K_{ij} , it is convenient to split K_{ij} into the trace part

$$K := \gamma^{ij} K_{ij}, \quad (2.51)$$

and its traceless part

$$A_{ij} := K_{ij} - \frac{1}{3}\gamma_{ij}K, \quad \text{tr}_\gamma A_{ij} = \gamma^{ij}A_{ij} = 0. \quad (2.52)$$

Therefore, we can obtain the traceless decomposition of the extrinsic curvature

$$K_{ij} = A_{ij} + \frac{1}{3}\gamma_{ij}K, \quad K^{ij} = A^{ij} + \frac{1}{3}\gamma^{ij}K. \quad (2.53)$$

The evolution equations of the spatial metric (2.42) in conformal decomposition formulation can hence be written as

$$\partial_t \psi = \beta^i \tilde{D}_i \psi - \frac{1}{6} \psi \left(\alpha K - \tilde{D}_i \beta^i \right), \quad (\text{conformal factor evolution}) \quad (2.54)$$

$$\partial_t \tilde{\gamma}_{ij} = -2\alpha \psi^{-4} A_{ij} + \tilde{\gamma}_{jk} \tilde{D}_i \beta^k + \tilde{\gamma}_{ik} \tilde{D}_j \beta^k - \frac{1}{3} \tilde{\gamma}_{ij} \tilde{D}_k \beta^k, \quad (\text{conformal metric evolution}) \quad (2.55)$$

and the constraint equations become

$$\tilde{D}^2 \psi - \frac{1}{8} \psi \tilde{R} + \left(\frac{1}{8} A_{ij} A^{ij} - \frac{1}{12} K^2 + 2\pi E \right) \psi^5 = 0, \quad (\text{Hamiltonian constraint}) \quad (2.56)$$

$$\tilde{D}_i (\psi^{10} A^{ij}) - \frac{2}{3} \psi^6 \tilde{D}^i K = 8\pi \psi^{10} S^i. \quad (\text{momentum constraint}) \quad (2.57)$$

Conformal transformation of the traceless part

We consider the transformation

$$A^{ij} := \psi^a \tilde{A}^{ij}, \quad (2.58)$$

for some undetermined exponent α . Here, we discuss two natural choices of a : $a = -4$ and $a = -10$.

“Time-evolution” scaling: $a = -4$. This choice of scaling was considered by Nakamura in 1994 [6]. It comes naturally from the evolution equation of conformal metric (2.55), where the $\psi^{-4} A_{ij}$ term suggests the conformal transformation of A_{ij} to have the same scaling factor as the conformal spatial metric (2.46)

$$\tilde{A}^{ij} := \psi^4 A^{ij}, \quad (2.59)$$

where the indices of \tilde{A}^{ij} and \tilde{A}_{ij} are lowered and raised by the conformal metric $\tilde{\gamma}_{ij}$ and $\tilde{\gamma}^{ij}$ respectively (i.e. $\tilde{A}_{ij} = \tilde{\gamma}_{il}\tilde{\gamma}_{jm}\tilde{A}^{lm} = \psi^{-4}A_{ij}$). The evolution equations of conformal spatial metric therefore become

$$\partial_t \tilde{\gamma}_{ij} = -2\alpha \tilde{A}_{ij} + \tilde{\gamma}_{jk} \tilde{D}_i \beta^k + \tilde{\gamma}_{ik} \tilde{D}_j \beta^k - \frac{1}{3} \tilde{\gamma}_{ij} \tilde{D}_k \beta^k. \quad (2.60)$$

The Hamiltonian constraint and momentum constraint in this scaling is rewritten as

$$\tilde{D}^2 \psi = \frac{1}{8} \psi \tilde{R} - \left(2\pi E - \frac{1}{12} K^2 + \frac{1}{8} \tilde{A}_{ij} \tilde{A}^{ij} \right) \psi^5, \quad (2.61)$$

$$\tilde{D}_i \left(\psi^6 \tilde{A}^{ij} \right) - \frac{2}{3} \psi^6 \tilde{D}^i K = 8\pi \psi^{10} S^i. \quad (2.62)$$

“Momentum-constraint” scaling: $a = -10$. Another possible choice of scaling factor $a = -10$ originates from the momentum constraint equation (2.57), which was first suggested by Lichnerowicz in 1944 [4]. we define

$$\hat{A}^{ij} := \psi^{10} \tilde{A}^{ij}, \quad (2.63)$$

and thus

$$\hat{A}_{ij} = \psi^2 \tilde{A}_{ij}. \quad (2.64)$$

Here, we use hat symbol to separate the "momentum-constraint" scaling from the tilde symbol in "time-evolution" scaling. These two scaling are related by

$$\hat{A}^{ij} = \psi^6 \tilde{A}^{ij}, \quad \hat{A}_{ij} = \psi^6 \tilde{A}_{ij}, \quad \hat{A}_{ij} \hat{A}^{ij} = \psi^{12} \tilde{A}_{ij} \tilde{A}^{ij} \quad (2.65)$$

Therefore, the constraint equations can be written as

$$\tilde{D}^2 \psi = \frac{1}{8} \psi \tilde{R} - \left(2\pi E - \frac{1}{12} K^2 \right) \psi^5 - \frac{1}{8} \hat{A}_{ij} \hat{A}^{ij} \psi^{-7}, \quad (2.66)$$

$$\tilde{D}_i \hat{A}^{ij} - \frac{2}{3} \psi^6 \tilde{D}^i K = 8\pi \psi^{10} S^i. \quad (2.67)$$

Equation(2.67) is known as *Lichnerowicz equation*. Although equation (2.62) and equation (2.67) are equivalent, they have different mathematical properties if they are treated as a partial differential equation for ψ . According to the maximal principle, the local uniqueness of the solutions depends on the sign of the exponent of ψ in the quadratic extrinsic curvature A^2 term [2, 3, 7–9]. Equation (2.62) suffers from

the mathematical nonuniqueness problems due to the positive exponent (+5), while the negative exponent (−7) in equation (2.67) guarantees the local uniqueness of the solutions.

2.3.3 Conformal transverse-traceless decomposition

Using the "momentum-constraint" scaling mentioned previously, we can further decompose the symmetric, traceless tensor \hat{A}^{ij} as

$$\hat{A}^{ij} = \hat{A}_{TT}^{ij} + \hat{A}_L^{ij}, \quad (2.68)$$

where \hat{A}_{TT}^{ij} is the transverse-traceless part which is divergenceless

$$\tilde{D}_j \hat{A}_L^{ij} = 0, \quad (2.69)$$

and \hat{A}_L^{ij} is the longitudinal part satisfying

$$\hat{A}_L^{ij} = \tilde{D}^i X^j + \tilde{D}^j X^i - \frac{2}{3} \tilde{\gamma}^{ij} \tilde{D}_k X^k \equiv \left(\tilde{L} W \right)^{ij}. \quad (2.70)$$

The vector X^i here is the vector potential and \tilde{L} is the *longitudinal operator* or *conformal Killing operator* associated with $\tilde{\gamma}$ which gives a symmetric, traceless tensor. The divergence of \hat{A}^{ij} becomes

$$\tilde{D}_j \hat{A}^{ij} = \tilde{D}^2 X^i + \frac{1}{3} \tilde{D}^i \left(\tilde{D}_j X^j \right). \quad (2.71)$$

Thus, the momentum constraint in the conformal transverse-traceless (CTT) decomposition yields

$$\tilde{D}^2 X^i + \frac{1}{3} \tilde{D}^i \left(\tilde{D}_j X^j \right) - \frac{2}{3} \psi^6 \tilde{D}^i K = 8\pi \psi^{10} S^i, \quad (2.72)$$

with the Hamiltonian constraint same as equation (2.66).

2.4 Gauge Condition

2.4.1 Maximal Slicing

2.4.2 Generalized Dirac Gauge

2.5 Constrained scheme for the Einstein equations

2.5.1 The Conformal Flatness Condition

2.5.2 The Fully Constrained Formulation

Chapter 3

Formulations of the Relativistic Hydrodynamics

3.1 The 3+1 "Valencia" formulation

3.2 The reference-metric formulism

3.3 Conserved to Primitive variables conversion

Chapter 4

NUMerical Methods and Tests

4.1 Multigrid Method for elliptic equations

4.1.1 Overview

4.1.2 Smoother

4.1.3 Restriction and Prolongation

4.1.4 Multigrid cycle

4.1.5 Grid Structure

4.1.6 Ghost cells and refinement boundary

4.1.7 Numerical Tests

4.2 Numerical method for hydrodynamics

4.2.1 Finite volume methods

4.2.2 Time Discretization

4.2.3 Atmosphere Treatment

4.2.4 Numerical Tests

References

- [1] R Amowitt, S Deser, and CW Misner. “The dynamics of general relativity”. In: *Gravitation: An Introduction to Current Research* (1962), pp. 227–265.
- [2] Isabel Cordero-Carrión et al. “Improved constrained scheme for the Einstein equations: An approach to the uniqueness issue”. In: *Physical Review D* 79.2 (2009), p. 024017.
- [3] Lawrence C Evans. “Partial differential equations and Monge-Kantorovich mass transfer”. In: *Current developments in mathematics* 1997.1 (1997), pp. 65–126.
- [4] André Lichnerowicz. *L’intégration des équations de la gravitation relativiste et le problème des n-corps*. Gauthier-Villars, 1944.
- [5] Charles W Misner, Kip S Thorne, and John Archibald Wheeler. *Gravitation*. Macmillan, 1973.
- [6] Takashi Nakamura. “3D numerical relativity.” In: *Relativistic Cosmology* (1994), pp. 155–182.
- [7] Murray H Protter and Hans F Weinberger. *Maximum principles in differential equations*. Springer Science & Business Media, 2012.
- [8] Larry L Smarr. *Sources of Gravitational Radiation: Proceedings of the Battelle Seattle Workshop*. CUP Archive, 1979.
- [9] Michael Eugene Taylor. *Partial differential equations. 3, Nonlinear equations*. Springer, 1991.
- [10] James W York Jr. “Conformally invariant orthogonal decomposition of symmetric tensors on Riemannian manifolds and the initial-value problem of general relativity”. In: *Journal of Mathematical Physics* 14.4 (1973), pp. 456–464.
- [11] James W York Jr. “Gravitational degrees of freedom and the initial-value problem”. In: *Physical Review Letters* 26.26 (1971), p. 1656.
- [12] James W York Jr. “Kinematics and dynamics of general relativity”. In: *Sources of gravitational radiation* (1979), pp. 83–126.
- [13] James W York Jr. “Role of conformal three-geometry in the dynamics of gravitation”. In: *Physical review letters* 28.16 (1972), p. 1082.

Appendix A

Useful relations for implementation of constrained scheme

A.1 The elliptic equations in constrained scheme

A.2 Generalized Dirac gauge conditions

Appendix B

Reference flat metric in 3D

