

# Continuity with respect to initial conditions for a numerical approximation of Kolmogorov equations.

Francisco Delgado-Vences\*, Saul Diaz-Infante† and Alan Matsumiya‡

November 6, 2018

## Abstract

In this paper we prove that the numerical method proposed in [13] is continuous with respect to the initial condition.

## 1 Introduction

Stochastic Partial Differential Equations (SPDEs) are important tools in modeling complex phenomena, they arise in many fields of knowledge like Physics, Biology, Economy, Finance, etc.. Develop efficient numerical methods for simulating SPDEs is very important but also very difficult and challenging.

The Fokker-Planck-Kolmogorov (FPK) equation is a partial differential equation that describes the time evolution of the probability density function of the velocity of a particle under the influence of drag forces and random forces, it is a kind of continuity equation for densities. Citing [10] “parabolic equations on Hilbert spaces appear in mathematical physics to model systems with infinitely many degrees of freedom. Typical examples are provided by spin configurations in statistical mechanics and by crystals in solid state theory. Infinite-dimensional parabolic equations provide an analytic description of infinite dimensional diffusion processes in such branches of applied mathematics as population biology, fluid dynamics, and mathematical finance.”. This kind of equations have been deeply studied in the last years, see for instance [2], [12], [7] and the references therein.

This paper is organized as follows. In section 2 we review the Fokker-Planck-Kolmogorov equation associated with SPDEs in a separable Hilbert space.

## 2 Kolmogorov equations for SPDEs in Hilbert spaces

Let  $\mathcal{H}$  be a separable infinite-dimensional Hilbert space with inner product  $(\cdot, \cdot)_{\mathcal{H}}$  and norm  $\|\cdot\|_{\mathcal{H}}$ . We define a Gaussian measure  $\mu$  with mean zero and nuclear covariance operator  $\Lambda$  with  $Tr(\Lambda) < +\infty$ . We focus on the following Kolmogorov equation

$$\frac{\partial u}{\partial t} = \frac{1}{2}Tr(QD^2u) + \langle Ax, Du \rangle_{\mathcal{H}} + \langle B(x), Du \rangle_{\mathcal{H}}, \quad x \in D(A). \quad (2.1)$$

Several authors have proved results on existence and uniqueness of the solution of the Kolmogorov equations, see for instance Da Prato [7] for a survey, Da Prato-Debussche [9] for the Burgers equation, Barbu-Da Prato [1] for the 2D Navier-Stokes stochastic flow in a channel.

---

\*CONACYT-UNAM, Instituto de Matemáticas, Sede Oaxaca, México. Email addresses: delgado@im.unam.mx

†CONACYT-Universidad de Sonora, Departamento de Matemáticas. Hermosillo, Sonora, México. Email address: saul.diazinfante@unison.mx

‡Universidad de Sonora, Departamento de Matemáticas. Hermosillo, Sonora, México. Email address: @.

## 2.1 On the Ornstein-Uhlenbeck semigroup

Following [6], in  $\mathcal{H}$  we define a Gaussian measure  $\mu$  with mean zero and nuclear covariance operator  $\Lambda$  with  $Tr(\Lambda) < +\infty$  and since  $\Lambda : \mathcal{H} \mapsto \mathcal{H}$  is a positive definite, self-adjoint operator then its square-root operator  $\Lambda^{1/2}$  is a positive definite, self-adjoint Hilbert-Schmidt operator on  $\mathcal{H}$ .

Define the inner product

$$(g, h)_0 := (\Lambda^{-1/2}g, \Lambda^{-1/2}h)_{\mathcal{H}}, \quad \text{for } g, h \in \Lambda^{1/2}\mathcal{H}.$$

Let  $\mathcal{H}_0$  denote the Hilbert subspace of  $\mathcal{H}$ , which is the completion of  $\Lambda^{1/2}\mathcal{H}$  with respect to the norm  $\|g\|_0 := (g, g)_0^{1/2}$ . Then  $\mathcal{H}_0$  is dense in  $\mathcal{H}$  and the inclusion map  $i : \mathcal{H}_0 \hookrightarrow \mathcal{H}$  is compact. The triple  $(i, \mathcal{H}_0, \mathcal{H})$  forms an abstract Wiener space.

Let  $\mathbb{H} = L^2(\mathcal{H}, \mu)$  denote the Hilbert space of Borel measurable functionals on the probability space with inner product

$$[\Phi, \Psi]_{\mathbb{H}} := \int_{\mathcal{H}} \Phi(v)\Psi(v)\mu(dv), \quad \text{for } \Phi, \Psi \in \mathbb{H},$$

and norm  $\|\Phi\|_{\mathbb{H}} := [\Phi, \Phi]_{\mathbb{H}}^{1/2}$ . We choose a basis system  $\{\varphi_k\}$  for  $\mathcal{H}$ .

A functional  $\Phi : \mathcal{H} \mapsto \mathbb{R}$ , is said to be a smooth simple functional (or a cylinder functional) if there exists a  $C^\infty$ -function  $\phi$  on  $\mathbb{R}^n$  and  $n$ -continuous linear functional  $l_1, \dots, l_n$  on  $\mathcal{H}$  such that for  $h \in \mathcal{H}$

$$\Phi(h) = \phi(h_1, \dots, h_n) \quad \text{where} \quad h_i = l_i(h), \quad i = 1, \dots, n. \quad (2.2)$$

The set of all such functionals will be denoted by  $\mathcal{S}(\mathbb{H})$ .

Denote by  $P_k(x)$  the Hermite polynomial of degree  $k$  taking values in  $\mathbb{R}$ . Then,  $P_k(x)$  is given by the following formula

$$P_k(x) = \frac{(-1)^k}{(k!)^{1/2}} e^{\frac{x^2}{2}} \frac{d^k}{dx^k} e^{-\frac{x^2}{2}}$$

with  $P_0 = 1$ . It is well-known that  $\{P_k(\cdot)\}_{k \in \mathbb{N}}$  is a complete orthonormal system for  $L^2(\mathbb{R}, \mu_1(dx))$

with  $\mu_1(dx) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$ .

Define the set of infinite multi-index as

$$\mathcal{J} = \left\{ \alpha = (\alpha_i, i \geq 1) \mid \alpha_i \in \mathbb{N} \cup \{0\}, \quad |\alpha| := \sum_{i=1}^{\infty} \alpha_i < +\infty \right\}$$

For  $n \in \mathcal{J}$  define the *Hermite polynomial functionals* on  $\mathcal{H}$  by

$$H_n(h) = \prod_{i=1}^{\infty} P_{n_i}(l_i(h)), \quad h \in \mathcal{H}_0, \quad n \in \mathcal{J}, \quad (2.3)$$

and where

$$l_i(h) = \langle h, \Lambda^{-1/2}\varphi_i \rangle_{\mathcal{H}}, \quad i = 1, 2, \dots$$

where  $P_n(\xi)$  is the usual Hermite polynomial for  $\xi \in \mathbb{R}$  and  $n \in \mathbb{N}$ .

**Remark 2.1.** Notice that  $l_i(h)$  is defined only for  $h \in \mathcal{H}_0$ . However, regarding  $h$  as a  $\mu$ -random variable in  $\mathcal{H}$ , we have  $\mathbb{E}(l_i(h)) = \|\varphi_i\|^2 = 1$  and then  $l_k(h)$  can be defined  $\mu$ -a.e.  $h \in \mathcal{H}$ , similar to defining a stochastic integral.

It is possible to identify the Hermite polynomial functionals defined in (2.3), for  $h \in \mathcal{H}_0$ , as a deterministic version of the Wick polynomials defined on the canonical Wiener space. (for further details see [14] for instance).

We have the following result (See Theorems 9.1.5 and 9.1.7 in Da Prato-Zabczyk [10] or Lemma 3.1 in chapter 9 from Chow [6]).

**Lemma 2.2.** For  $h \in \mathcal{H}$  let  $l_i(h) = \langle h, \Lambda^{-1/2} \varphi_i \rangle_{\mathcal{H}}$ ,  $i = 1, 2, \dots$ . Then the set  $\{H_n\}$  of all Hermite polynomials on  $\mathcal{H}$  forms a complete orthonormal system for  $\mathbb{H}$ . Hence the set of all functionals are dense in  $\mathbb{H}$ . Moreover, we have the direct sum decomposition:

$$\mathbb{H} = \bigoplus_{j=0}^{\infty} K_j,$$

where  $K_j$  is the subspace of  $\mathbb{H}$  spanned by  $\{H_n : |n| = j\}$ .  $\square$

Let  $\Phi$  be a smooth simple functional given by (2.2). Then the Fréchet derivatives,  $D\Phi = \Phi'$  and  $D_2\Phi = \Phi''$  in  $\mathcal{H}$  can be computed as follows:

$$\begin{aligned} (D\Phi(h), v) &= \sum_{k=1}^n [\partial_k \phi(h_1, \dots, h_n)] l_k(v) \\ (D^2\Phi(h), v) &= \sum_{j,k=1}^n [\partial_j \partial_k \phi(h_1, \dots, h_n)] l_j(v) l_k(v) \end{aligned} \quad (2.4)$$

for any  $u, v \in \mathcal{H}$ , where  $\partial_k \phi = \frac{\partial}{\partial h_k} \phi$ . Similarly, for  $m > 2$ ,  $D^m \Phi(h)$  is a  $m$ -linear form on  $\mathcal{H}^m$  with inner product  $(\cdot, \cdot)_m$ . We have  $[D^m \Phi(h)](v_1, \dots, v_m) = (D^m \Phi(h), v_1 \otimes \dots \otimes v_m)_m$ , for  $h, v_1, \dots, v_m \in \mathcal{H}$ .

Consider the following linear stochastic equation

$$\begin{aligned} du_t &= Au_t dt + dW_t, \\ u_0 &= h \in \mathcal{H}. \end{aligned} \quad (2.5)$$

Where  $A : \mathcal{D}(A) \subset \mathcal{H} \rightarrow \mathcal{H}$  is the infinitesimal generator of a strongly continuous semigroup  $e^{tA}$  in  $\mathcal{H}$ .  $W_t$  is a  $Q$ -Wiener process in  $\mathcal{H}$ .

Chow in [6, Lemma 9.4.1] has shown the following result.

**Lemma 2.3.** Suppose that  $A$  and  $Q$  satisfy the following:

1.  $A : \mathcal{D}(A) \subset \mathcal{H} \rightarrow \mathcal{H}$  is self-adjoint and there is  $\beta > 0$  such that

$$\langle Av, v \rangle_{\mathcal{H}} \leq -\beta \|v\|_{\mathcal{H}}^2 \quad \forall v \in \mathcal{H}.$$

2.  $A$  commutes with  $Q$  in  $\mathcal{D}(A) \subset \mathcal{H}$ .

Then (2.5) has a unique invariant measure  $\mu$  which is a Gaussian measure on  $\mathcal{H}$  with zero mean and covariance operator  $\Lambda = \frac{1}{2}Q(-A)^{-1} = \frac{1}{2}(-A)^{-1}Q$ .  $\square$

Suppose that  $A$  and  $Q$  have the same eigenfunctions  $e_k$  with eigenvalues  $\lambda_k$  and  $\rho_k$  respectively. It is well-known (See for instance Da Prato and Zabczyk [10]) that the solution of (2.5) is a time-homogeneous Markov process with transition operator  $P_t$  defined for  $\Phi \in \mathbb{H}$  given by

$$(P_t \Phi)(h) = \int_{\mathcal{H}} \Phi(v) \mu_t^h(dv) = \mathbb{E}[\Phi(u_t^h)] \quad (2.6)$$

Let  $\Phi \in \mathcal{S}(\mathbb{H})$  be a smooth simple functional. By setting  $\varphi_k = e_k$  in (2.2), it takes the form

$$\Phi(h) = \phi(l_1(h), \dots, l_n(h)),$$

where  $l_k(h) = (h, \Lambda^{1/2} e_k)$ . Define a differential operator  $A_0$  on  $\mathcal{S}(\mathbb{H})$  by

$$A_0 \Phi(v) = \frac{1}{2} \text{Tr}[RD^2\Phi(v)] + \langle Av, D\Phi(v) \rangle, \quad v \in \mathcal{H}, \quad (2.7)$$

which is well defined, since  $D\Phi \in \mathcal{D}(A)$  and  $\langle Av, D\Phi(v) \rangle = (v, AD(v))_{\mathcal{H}}$ .

The following results have been proved in [6].

**Lemma 2.4.** *Let  $P_t$  be the transition operator as defined by (2.5). Then the following properties hold:*

1.  $P_t : \mathcal{S}(\mathbb{H}) \rightarrow \mathcal{S}(\mathbb{H})$  for  $t \geq 0$ .
2.  $\{P_t, t \geq 0\}$  is a strongly continuous semigroup on  $\mathcal{S}(\mathbb{H})$  so that, for any  $\Phi \in \mathcal{S}(\mathbb{H})$ , we have  $P_0 = I$ ,  $P_{t+s}\Phi = P_t P_s \Phi$ , for all  $t, s \geq 0$ , and  $\lim_{t \downarrow 0} P_t \Phi = \Phi$ .
3.  $\mathcal{A}_0$  is the infinitesimal generator of  $P_t$  so that, for each  $\Phi \in \mathcal{S}(\mathbb{H})$ ,

$$\lim_{t \downarrow 0} \frac{1}{t} (P_t - I) \Phi = \mathcal{A}_0 \Phi.$$

□

**Lemma 2.5.** *Let  $H_n(h)$  be a Hermite polynomial functional given by (2.3). Then the following hold:*

$$\mathcal{A}_0 H_n(h) = \lambda_n H_n(h), \quad (2.8)$$

and

$$P_t H_n(h) = \exp\{\lambda_n t\} H_n(h), \quad (2.9)$$

for any  $\mathbf{n} \in \mathcal{J}$  and  $h \in H$ , where

$$\lambda_n = \sum_{i=1}^{\infty} n_i \lambda_i.$$

□

The following Theorem is a Green formula we will need forward. Its proof can be seen, for instance, in [6], Theorem 3.3, chapter 9.

**Theorem 2.6.** *Let  $\Phi \in \mathcal{S}(\mathbb{H})$  be a smooth simple functional and let  $\mu \sim N(0, \Lambda)$  be a Gaussian measure in  $\mathcal{H}$ . Then, for any  $g, h \in \mathcal{H}$  the following formula holds*

$$\int_{\mathcal{H}} (\Lambda h, D\Phi(v))_{\mathcal{H}} \mu(dv) = \int_{\mathcal{H}} (v, h)_{\mathcal{H}} \Phi(v) \mu(dv) \quad (2.10)$$

□

**Lemma 2.7.** *Assume the conditions for Lemma 2.5 hold. Then, for any  $\Phi, \Psi \in \mathcal{S}(\mathbb{H})$ , the following Greens formula holds:*

$$\int_{\mathcal{H}} (\mathcal{A}_0 \Phi) \Psi d\mu = \int_{\mathcal{H}} \Phi (\mathcal{A}_0 \Psi) d\mu = -\frac{1}{2} \int_{\mathcal{H}} (QD\Phi, D\Psi) d\mu \quad (2.11)$$

□

By Lemma 2.2, for  $\Phi \in \mathbb{H}$ , it can be represented as

$$\Phi(v) = \sum_{n=0}^{\infty} \phi_n H_n(v), \quad (2.12)$$

where  $n = |\mathbf{n}|$  and  $\mathbf{n} \in \mathcal{J}$ . Notice that we can think in  $\mathbf{n}$  as a vector of  $r$  dimension, i.e.  $\mathbf{n} = (n_1, \dots, n_r)$ . Let  $\alpha_{\mathbf{n}} = \alpha_{n_1} \cdots \alpha_{n_r}$  be a sequence of positive numbers with  $\alpha_{\mathbf{n}} > 0$ , such that  $\alpha_{\mathbf{n}} \rightarrow \infty$  as  $n \rightarrow \infty$ . Define

$$\begin{aligned} |||\Phi|||_{k,\alpha} &= \left[ \sum_{\mathbf{n}} (1 + \alpha_{\mathbf{n}})^k |\phi_n|^2 \right]^{1/2} \\ |||\Phi|||_{0,\alpha} &= |||\Phi||| = \left[ \sum_{\mathbf{n}} |\phi_n|^2 \right]^{1/2} \end{aligned}$$

which is  $L^2(\mu)$ -norm of  $\Phi$ . For the given sequence  $\alpha = \{\alpha_n\}$ , let  $\mathbb{H}_{k,\alpha}$  denote the completion of  $S(\mathbb{H})$  with respect to the norm  $||| \cdot |||_{k,\alpha}$ . Then  $\mathbb{H}_{k,\alpha}$  is called a GaussSobolev space of order  $k$  with parameter  $\alpha$ . The dual space of  $\mathbb{H}_{k,\alpha}$  is  $\mathbb{H}_{-k,\alpha}$ . From now on, we will fix the sequence  $\alpha_n = \lambda_n$ , where  $\lambda_n$  is given in Lemma 2.5. We shall simply denote  $\mathbb{H}_{k,\alpha}$  by  $\mathbb{H}_k$  and  $|||\Phi|||_{k,\alpha}$  by  $|||\Phi|||_k$ .

The following results ensure the existence of an extension for the operator  $\mathcal{A}_0$  to a domain containing  $\mathbb{H}_2$ . Their proofs can be found in [6] for instance.

**Theorem 2.8.** *Let the conditions on  $A$  and  $Q$  in Lemma 2.3 hold. Then  $P_t : \mathbb{H} \rightarrow \mathbb{H}$ , for  $t \geq 0$ , is a contraction semigroup with the infinitesimal generator  $\tilde{A}$ . The domain of  $\tilde{A}$  contains  $\mathbb{H}_2$  and we have  $\tilde{A} = \mathcal{A}_0$  in  $S(\mathbb{H})$ .  $\square$*

**Theorem 2.9.** *Let the conditions for Theorem 2.8 hold true. The differential operator  $\mathcal{A}_0$  defined by (2.7) in  $S(\mathbb{H})$  can be extended to be a self-adjoint linear operator  $A$  in  $\mathbb{H}$  with domain  $\mathbb{H}_2$ .  $\square$*

Since both  $\tilde{A}$  and  $A$  are extensions of  $\mathcal{A}_0$  to a domain containing  $\mathbb{H}_2$ , they must coincide there.

Given the Gauss-Sobolev space  $\mathbb{H}_k$  with norm  $||| \cdot |||_k$  we denote its dual space by  $\mathbb{H}_{-k}$  with norm  $||| \cdot |||_{-k}$ . Thus, we have the following inclusions,

$$\mathbb{H}_k \subset \mathbb{H} \subset \mathbb{H}_{-k}.$$

We denote the duality between  $\mathbb{H}_k$  and  $\mathbb{H}_{-k}$  by

$$\langle \langle \Psi, \Phi \rangle \rangle_k, \quad \Phi \in \mathbb{H}_k, \quad \Psi \in \mathbb{H}_{-k}$$

We also set  $\mathbb{H}_0 = \mathbb{H}$ , with  $||| \cdot |||_0 = ||| \cdot |||$  and  $\langle \langle \cdot, \cdot \rangle \rangle_1 = \langle \langle \cdot, \cdot \rangle \rangle$ ,  $\langle \langle \cdot, \cdot \rangle \rangle_0 = [\cdot, \cdot]$ .

### 2.2 A non linear Kolmogorov equation

Consider the following Kolmogorov equation,

$$\begin{aligned} \frac{\partial}{\partial t} \Psi(v, t) &= \mathcal{A} \Psi(v, t) + \langle B(v), D \Psi(v, t) \rangle_{\mathcal{H}}, \quad \text{a.e. } v \in \mathbb{H}_2 \\ \Psi(v, 0) &= \phi(v) \end{aligned} \quad (2.13)$$

where, as defined in Theorem 2.8,  $\mathcal{A} : \mathbb{H}_2 \rightarrow \mathbb{H}$  is given by

$$\mathcal{A} \Phi = \frac{1}{2} \text{Tr}[R D^2 \Phi(v)] + \langle A v, D \Phi(v) \rangle \quad (2.14)$$

Hypothesis on  $B$  will be specified latter. For now, we will consider that it is a locally Lipschitz function. The additional term  $\langle B(v), D \Psi(v, t) \rangle_{\mathcal{H}}$  is defined  $\mu$ -a.e.  $v \in \mathbb{H}_2$ . We will allow the initial datum  $\phi$  will be in  $\mathbb{H}$ .

We will study a mild solution of the equation (2.13). Let  $\lambda > 0$  be a parameter. By changing  $\Psi$  to  $e^{\lambda t} \Psi$  in (2.13) we get the following equation:

$$\begin{aligned} \frac{\partial}{\partial t} \Psi(v, t) &= \mathcal{A}_\lambda \Psi(v, t) + \langle B(v), D \Psi(v, t) \rangle_{\mathcal{H}}, \quad \text{a.e. } v \in \mathbb{H}_2 \\ \Psi(v, 0) &= \phi(v) \end{aligned} \quad (2.15)$$

where  $\mathcal{A}_\lambda = \mathcal{A} - \lambda I$ , with  $I$  the identity operator in  $\mathbb{H}$ . Clearly, the problems (2.13) and (2.15) are equivalent, as far for the existence and uniqueness questions are concerned. We will work on the problem (2.15).

Denote by  $P_t$  the semigroup with infinitesimal generator  $\mathcal{A}_\lambda$ . The existence of  $P_t$  is ensured by the Theorem 2.8. Then, we can rewrite the equation (2.15) in an integral form by using the semigroup  $P_t$

$$\Psi(v, t) = e^{-\lambda t} (P_t \phi)(v) + \int_0^t e^{-\lambda(t-s)} [P_{t-s}(B, D \Psi_s)](v) ds, \quad (2.16)$$

where we denote  $\phi = \phi(\cdot)$  and  $\Psi_s = \Psi(\cdot, s)$ .

Chow [6] had proved the following lemma.

**Lemma 2.10.** *Let  $\Psi \in L^2((0, T); \mathbb{H})$  for some  $T > 0$ . Then, for any  $\lambda > 0$  there exists  $C_\lambda > 0$  such that*

$$||| \int_0^t e^{-\lambda(t-s)} P_{t-s} \Psi_s ds |||^2 \leq C_\lambda \int_0^t ||| \Psi_s |||_{-1}^2 ds \quad 0 < t \leq T \quad (2.17)$$

□

We now prove the following theorem on existence and uniqueness of a mild solution to (2.15).

**Theorem 2.11.** *Suppose that  $B : \mathcal{H} \rightarrow \mathcal{H}_0$  satisfies  $(B, D\Phi) \in L^2((0, T); \mathbb{H})$  for any  $\Phi \in \mathbb{H}$  and*

$$\sup_{v \in \mathcal{H}} ||| \Lambda^{-1/2} B(v) |||_{\mathcal{H}} < +\infty. \quad (2.18)$$

*Then,  $B$  satisfies*

$$||| (B(v), D\Phi(v)) |||_{-1}^2 \leq C ||| \Phi(v) |||^2 \quad \text{for any } \Phi \in \mathbb{H}, \quad v \in \mathbb{H}_2 \quad (2.19)$$

*for some  $C > 0$ . Moreover, for  $\Phi \in \mathbb{H}$ , the initial-value problem (2.15) has a unique mild solution  $\Psi \in C((0, T); \mathbb{H})$ .*

For the part of the existence and uniqueness of the solution we will adapt the proof of the Theorem 5.2 in chapter 9 from [6].

*Proof.* First we will prove (2.19). We have

$$||| (B(v), D\Phi(v)) |||_{-1}^2 = \sum_{\mathbf{n}} (1 + \lambda_{\mathbf{n}})^{-1} |\phi_n|^2$$

with

$$\phi_n = \left( (B(v), D\Phi(v))_{\mathcal{H}}, H_{\mathbf{n}}(v) \right)_{\mathbb{H}} = \int_{\mathcal{H}} (B(v), D\Phi(v))_{\mathcal{H}} H_{\mathbf{n}}(v) \mu(dv). \quad (2.20)$$

By the Theorem 2.6, in particular (2.10), we have

$$\int_{\mathcal{H}} (\Lambda h, D\Phi(v))_{\mathcal{H}} \mu(dv) = \int_{\mathcal{H}} (v, h)_{\mathcal{H}} \Phi(v) \mu(dv)$$

for all  $\Phi \in \mathcal{S}(\mathbb{H})$ ,  $g, h \in \mathcal{H}$  and  $\mu \sim N(0, \Lambda)$ . Then, in particular, in each direction  $H_{\mathbf{n}}$  this formula is still true, so we have

$$\int_{\mathcal{H}} (\Lambda h, D\Phi(v))_{\mathcal{H}} H_{\mathbf{n}}(v) \mu(dv) = \int_{\mathcal{H}} (v, h)_{\mathcal{H}} \Phi(v) H_{\mathbf{n}}(v) \mu(dv)$$

Then, applying this last equality to (2.20) we get

$$\begin{aligned} \phi_n &= \int_{\mathcal{H}} (\Lambda [\Lambda^{-1} B(v)], D\Phi(v))_{\mathcal{H}} H_{\mathbf{n}}(v) \mu(dv) \\ &= \int_{\mathcal{H}} (\Lambda^{-1} B(v), v)_{\mathcal{H}} \Phi(v) H_{\mathbf{n}}(v) \mu(dv) \\ &= \int_{\mathcal{H}} (\Lambda^{-1/2} B(v), \Lambda^{1/2} v)_{\mathcal{H}} \Phi(v) H_{\mathbf{n}}(v) \mu(dv) \end{aligned}$$

Thus,

$$\begin{aligned} |\phi_n|^2 &= \left| \int_{\mathcal{H}} (\Lambda^{-1/2} B(v), \Lambda^{1/2} v)_{\mathcal{H}} \Phi(v) H_{\mathbf{n}}(v) \mu(dv) \right|^2 \\ &\leq \int_{\mathcal{H}} |(\Lambda^{-1/2} B(v), \Lambda^{1/2} v)_{\mathcal{H}}|^2 |H_{\mathbf{n}}(v)|^2 \mu(dv) \int_{\mathcal{H}} |\Phi(v)|^2 \mu(dv) \end{aligned} \quad (2.21)$$

We now focus on the first integral. Let  $I_1$  be the first integral of (2.21). Then,

$$\begin{aligned} I_1 &\leq \int_{\mathcal{H}} \|\Lambda^{-1/2} B(v)\|_{\mathcal{H}}^2 \|\Lambda^{1/2} v\|_{\mathcal{H}}^2 |H_{\mathbf{n}}(v)|^2 \mu(dv) \\ &\leq \sup_{v \in \mathcal{H}} \|\Lambda^{-1/2} B(v)\|_{\mathcal{H}}^2 \int_{\mathcal{H}} \|\Lambda^{1/2} v\|_{\mathcal{H}}^2 |H_{\mathbf{n}}(v)|^2 \mu(dv) \\ &\leq C \int_{\mathcal{H}} \|v\|_{\mathcal{H}}^2 |H_{\mathbf{n}}(v)|^2 \mu(dv) \\ &\leq C \end{aligned}$$

The last inequality follows by using proposition 3.11 in page 64 from [8]. Then, by using this bound on (2.21) we have.

$$\begin{aligned} |\phi_n|^2 &\leq C \int_{\mathcal{H}} |\Phi(v)|^2 \mu(dv) \\ &\leq C \|\Phi(v)\|^2 \end{aligned}$$

Thus,

$$\|(B(v), D\Phi(v))\|_{-1}^2 \leq C \|\Phi(v)\|^2 \sum_{\mathbf{n}} (1 + \lambda_{\mathbf{n}})^{-1} \leq C \|\Phi(v)\|^2$$

which proves (2.19).

We now prove the existence and uniqueness of a solution to the initial-value problem (2.15). Let  $\mathbb{X}_T$  denote the Banach space  $\mathcal{C}([0, T]; \mathbb{H})$  with the sup-norm

$$\|\Psi\|_T := \sup_{0 \leq t \leq T} \|\Psi(t)\|$$

In  $\mathbb{X}_T$  define the linear operator  $\mathbb{Q}$  as

$$\mathbb{Q}\Psi = e^{-\lambda t} P_t \Phi + \int_0^t e^{-\lambda(t-s)} P_{t-s}(B, D\Psi_s) ds, \quad \text{for any } \Psi \in \mathbb{X}_T.$$

By Theorem 2.8  $P_t$  is a contraction semigroup, then using this fact and Lemma 2.10 we have

$$\begin{aligned} \|\mathbb{Q}\Psi\|^2 &\leq 2 \left[ \|e^{-\lambda t} P_t \Phi\|^2 + \left\| \int_0^t e^{-\lambda(t-s)} P_{t-s}(B, D\Psi_s) ds \right\|^2 \right] \\ &\leq 2 \left[ \|\Phi\|^2 + C_{\lambda} \int_0^t \|(B, D\Psi_s)\|_{-1}^2 ds \right] \\ &\leq 2\|\Phi\|^2 + C_1 \int_0^t \|\Psi_s\|^2 ds, \end{aligned}$$

for some  $C_1 > 0$ . Hence,

$$\|\mathbb{Q}\Psi\|_T \leq C(1 + \|\Psi\|_T),$$

with  $C = C(\Phi, \lambda, T)$ . Then, the map  $\mathbb{Q} : \mathbb{X}_T \rightarrow \mathbb{X}_T$  is well defined. We now show that is a contraction for a small  $t$ . Let  $\Psi, \Psi' \in \mathbb{X}_T$ . Then

$$\begin{aligned} \|\mathbb{Q}\Psi - \mathbb{Q}\Psi'\|^2 &= \left\| \int_0^t e^{-\lambda(t-s)} P_{t-s}[(B, D\Psi_s) - (B, D\Psi'_s)] ds \right\|^2 \\ &\leq C_{\lambda} \int_0^t \|(B, D\Psi_s - D\Psi'_s)\|_{-1}^2 ds \\ &\leq C_2 \int_0^t \|\Psi_s - \Psi'_s\|^2 ds, \end{aligned}$$

### 3 CONTINUITY WITH RESPECT TO THE INITIAL CONDITIONS OF A NUMERICAL APPROXIMATIONS

For some  $C_2 > 0$ . It follows that

$$|||\mathbb{Q}\Psi - \mathbb{Q}\Psi'\|||_T \leq \sqrt{C_2 T} |||\Psi_s - \Psi'\|||_T.$$

Then, for small  $T$ ,  $\mathbb{Q}$  is a contraction on  $\mathbb{X}_T$ . Hence the Cauchy problem (2.15) has a unique mild solution.  $\square$

We now prove a theorem on the dependence on initial conditions for the mild solution of (2.15).

**Theorem 2.12.** *Suppose that  $B : \mathcal{H} \rightarrow \mathcal{H}_0$  satisfies  $(B, D\Phi) \in L^2((0, T); \mathbb{H})$  for any  $\Phi \in \mathbb{H}$  and*

$$\sup_{v \in \mathcal{H}} |||\Lambda^{-1/2} B(v)|||_{\mathcal{H}} < +\infty. \quad (2.22)$$

*Then, the unique mild solution  $\Psi \in C((0, T); \mathbb{H})$  for (2.15) depends continuously on the initial conditions.*

*Proof.* We know, with the assumption (2.22), that the existence of a unique mild solution for (2.15) is guaranteed by the Theorem 2.11. We will denote by  $\Psi_t^\varphi$  its mild solution at time  $t$  with initial condition  $\varphi$ :

$$\Psi_t^\varphi = e^{-\lambda t} P_t \varphi + \int_0^t e^{-\lambda(t-s)} P_{t-s}(B, D\Psi_s) ds$$

Then,

$$\begin{aligned} \Psi_t^\varphi - \Phi_t^\psi &= e^{-\lambda t} P_t \varphi - e^{-\lambda t} P_t \psi + \int_0^t e^{-\lambda(t-s)} P_{t-s}(B, D\Psi_s^\varphi - D\Phi_s^\psi) ds \\ &= e^{-\lambda t} P_t(\varphi - \psi) + \int_0^t e^{-\lambda(t-s)} P_{t-s}(B, D\Psi_s^\varphi - D\Phi_s^\psi) ds. \end{aligned}$$

From this expression we get

$$\begin{aligned} |||\Psi_t^\varphi - \Phi_t^\psi|||^2 &\leq |||e^{-\lambda t} P_t(\varphi - \psi)|||^2 + |||\int_0^t e^{-\lambda(t-s)} P_{t-s}(B, D\Psi_s^\varphi - D\Phi_s^\psi)|||^2 ds \\ &\leq |||\varphi - \psi|||^2 + C_\lambda \int_0^t |||(B, D\Psi_s^\varphi - D\Phi_s^\psi)|||_{-1}^2 ds \\ &\leq |||\varphi - \psi|||^2 + C_2 \int_0^t |||\Psi_s^\varphi - \Phi_s^\psi|||^2 ds. \end{aligned}$$

Thus, by Gronwall's inequality we obtain

$$|||\Psi_t^\varphi - \Phi_t^\psi|||^2 \leq \exp(C_2 t) |||\varphi - \psi|||^2 \quad (2.23)$$

which implies,

$$|||\Psi_t^\varphi - \Phi_t^\psi||| \leq \exp(Ct) |||\varphi - \psi|||$$

This completes the proof.  $\square$

### 3 Continuity with respect to the initial conditions of a numerical approximations

In this section we will prove the continuity with respect to the initial conditions for a numerical approximations of the Kolmogorov equations associated to SPDE's.



### 3 CONTINUITY WITH RESPECT TO THE INITIAL CONDITIONS OF A NUMERICAL APPROXIMATIONS

---

We use Lemma 2.2 to write the solution  $\Psi_t^\varphi$  as in a Fourier-Hermite decomposition:

$$\Psi_t^\varphi = \sum_{n \in \mathcal{J}} u_n(t) H_n(x), \quad x \in \mathcal{H}, \quad t \in [0, T] \quad (3.1)$$

Notice that the time-dependent coefficients  $u_n(t)$  depend on the functional and on the initial condition but it is not a function of the initial condition.

We now prove a previous result before proving the result on the dependence of the initial conditions.

**Lemma 3.1.** *Set  $\{P_k(\xi)\}_{k \in \mathbb{N}}$  the family of normalized Hermite polynomials in  $\mathbb{R}$ . For every  $k \in \mathbb{N}$  and  $\xi, \eta \in \mathbb{R}$  such that  $\eta < \xi$  we have that*

$$P_k(\xi) - P_k(\eta) = C(k) Pe_{k+1}(\gamma) \cdot (\xi - \eta), \quad (3.2)$$

where  $\gamma \in (\eta, \xi)$  and  $C(k) = \frac{(-1)^k}{(k+1)(k!)^{1/2}}$ . Moreover,  $Pe_k(x)$  is the unnormalized Hermite polynomial of  $k$  degree.

*Proof.* We know that

$$P_k(\xi) = \frac{(-1)^k}{(k!)^{1/2}} e^{\xi^2/2} \frac{d}{d\xi^k} e^{-\xi^2/2}.$$

Set  $c(k) = (-1)^k (k!)^{-1/2}$ , then

$$\begin{aligned} P_k(\xi) - P_k(\eta) &= c(k) \left[ e^{\xi^2/2} \frac{d}{d\xi^k} e^{-\xi^2/2} - e^{\eta^2/2} \frac{d}{d\eta^k} e^{-\eta^2/2} \right] \\ &= c(k) \left[ e^{x^2/2} \frac{d}{dx^k} e^{-x^2/2} \Big|_{x=\eta}^\xi \right] \\ &= c(k) \int_\eta^\xi F_k(x) dx \end{aligned}$$

where  $F_k$  is a continuous function such that  $F'_k(x) = e^{x^2/2} \frac{d}{dx^k} e^{-x^2/2}$ . In fact, denoting by  $Pe_k(x)$  the unnormalized Hermite polynomial of  $k$  degree, then

$$F'_k(x) = e^{x^2/2} \frac{d}{dx^k} e^{-x^2/2} = Pe_k(x)$$

and since the Hermite polynomials constitute an Appell sequence we have that

$$F'_k(x) = Pe_k(x) = \frac{1}{k+1} Pe'_{k+1}(x)$$

which implies that  $F_k(x) = \frac{1}{k+1} Pe_{k+1}(x)$ . Now, since  $F_k(x)$  is a continuous function, then there exists  $\gamma \in (\eta, \xi)$  such that

$$\int_\eta^\xi F_k(x) dx = F_k(\gamma) \cdot (\xi - \eta).$$

All these implies that

$$P_k(\xi) - P_k(\eta) = c(k) F_k(\gamma) \cdot (\xi - \eta).$$

From this expression the lemma follows immediately.  $\square$

Consider the stochastic differential equation in  $\mathcal{H}$

$$dX_t = AX_t dt + B(X_t) dt + \sqrt{Q} dW_t, \quad (3.3)$$

where the operator  $A : \mathcal{D}(A) \subset \mathcal{H} \rightarrow \mathcal{H}$  is the infinitesimal generator of a strongly continuous semigroup  $e^{tA}$  in  $\mathcal{H}$ ,  $Q$  is a bounded operator from another Hilbert space  $\mathcal{U}$  to  $\mathcal{H}$  and  $B : \mathcal{D}(B) \subset \mathcal{H} \rightarrow \mathcal{H}$  is a nonlinear mapping.

### 3 CONTINUITY WITH RESPECT TO THE INITIAL CONDITIONS OF A NUMERICAL APPROXIMATIONS

---

The equation (3.3) can be associated to a Kolmogorov equation in the next way, we define

$$u(t, x) = \mathbb{E}[\varphi(X_t^x)], \quad (3.4)$$

where  $\varphi : \mathcal{H} \rightarrow \mathbb{R}$  and  $X_t^x$  is the solution to (3.3) with initial conditions  $X_0 = x$  where  $x \in \mathcal{H}$ . Then  $u$  satisfies the Kolmogorov equation (2.1). We will use some technical results on the SPDE to prove the following result.

We present the main result of this section.

**Theorem 3.2.** *Assume that the eigenvalues of  $\Lambda$ , satisfies that for every  $k \in \mathbb{N}$ ,  $\lambda_k < \lambda_{k+1} \rightarrow \infty$ . Assume that the functional  $\varphi$  is Borel. Then, The numeric approximation  $\Psi_t^\varphi$  (given by (3.1)) to the solution of the Kolmogorov equation  $\Psi \in C((0, T); \mathbb{H})$  also depends continuously on the initial conditions.*

*Proof.* Let  $x, y \in H$  be two different initial values. We want to estimate  $\Psi_t^x - \Psi_t^y$ . By definition,

$$\Psi_t^x = \sum_{\bar{n} \in \mathcal{J}} u_{\bar{n}}^x(t) H_{\bar{n}}(x) \quad (3.5)$$

Thus,

$$\begin{aligned} \Psi_t^x - \Psi_t^y &= \sum_{\bar{n} \in \mathcal{J}} u_{\bar{n}}^x(t) H_{\bar{n}}(x) - \sum_{\bar{n} \in \mathcal{J}} u_{\bar{n}}^y(t) H_{\bar{n}}(y) \\ &= \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t)] H_{\bar{n}}(x) + \sum_{\bar{n} \in \mathcal{J}} u_{\bar{n}}^y(t) [H_{\bar{n}}(x) - H_{\bar{n}}(y)] \end{aligned} \quad (3.6)$$

We focus on the first term in (3.6). From the definition of the initial condition we obtain the following expression for the time-dependent coefficient

$$u_{\bar{n}}^x(t) = \int_{\mathcal{H}} H_{\bar{n}}(x) \mathbb{E}[\varphi(X_t^x)] \mu(dx)$$

From this we get

$$\begin{aligned} u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t) &= \int_{\mathcal{H}} H_{\bar{n}}(x) \mathbb{E}[\varphi(X_t^x)] \mu(dx) - \int_{\mathcal{H}} H_{\bar{n}}(y) \mathbb{E}[\varphi(X_t^y)] \mu(dy) \\ &= \int_{\mathcal{H} \times \mathcal{H}} H_{\bar{n}}(x) \mathbb{E}[\varphi(X_t^x)] \mu(dx) \mu(dy) - \int_{\mathcal{H} \times \mathcal{H}} H_{\bar{n}}(y) \mathbb{E}[\varphi(X_t^y)] \mu(dx) \mu(dy) \\ &= \int_{\mathcal{H} \times \mathcal{H}} H_{\bar{n}}(x) \left( \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right) \mu(dx) \mu(dy) \\ &\quad + \int_{\mathcal{H} \times \mathcal{H}} \left( H_{\bar{n}}(x) - H_{\bar{n}}(y) \right) \mathbb{E}[\varphi(X_t^y)] \mu(dx) \mu(dy) \end{aligned}$$

Then, by using Cauchy-Schwartz we obtain

$$\begin{aligned} |u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t)|^2 &\leq \left| \int_{\mathcal{H} \times \mathcal{H}} H_{\bar{n}}(x) \left( \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right) \mu(dx) \mu(dy) \right|^2 \\ &\quad + \left| \int_{\mathcal{H} \times \mathcal{H}} \left( H_{\bar{n}}(x) - H_{\bar{n}}(y) \right) \mathbb{E}[\varphi(X_t^y)] \mu(dx) \mu(dy) \right|^2 \\ &\leq \int_{\mathcal{H} \times \mathcal{H}} H_{\bar{n}}^2(x) \mu(dx) \mu(dy) \int_{\mathcal{H} \times \mathcal{H}} \left| \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right|^2 \mu(dx) \mu(dy) \\ &\quad + \int_{\mathcal{H} \times \mathcal{H}} \mathbb{E}^2[\varphi(X_t^y)] \mu(dx) \mu(dy) \int_{\mathcal{H} \times \mathcal{H}} \left| H_{\bar{n}}(x) - H_{\bar{n}}(y) \right|^2 \mu(dx) \mu(dy) \end{aligned}$$

### 3 CONTINUITY WITH RESPECT TO THE INITIAL CONDITIONS OF A NUMERICAL APPROXIMATIONS

---

$$\begin{aligned}
&= \int_{\mathcal{H} \times \mathcal{H}} \left| \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right|^2 \mu(dx) \mu(dy) \\
&\quad + \int_{\mathcal{H} \times \mathcal{H}} \mathbb{E}^2[\varphi(X_t^y)] \mu(dx) \mu(dy) \int_{\mathcal{H} \times \mathcal{H}} \left| H_{\bar{n}}(x) - H_{\bar{n}}(y) \right|^2 \mu(dx) \mu(dy) \quad (3.7)
\end{aligned}$$

We now estimate the norm of the expression (3.6) with the help of (3.7).

$$\begin{aligned}
\|\Psi_t^x - \Psi_t^y\|_{(L^2(\mathcal{H}, \mu))^2}^2 &= \int_{\mathcal{H} \times \mathcal{H}} |\Psi_t^x - \Psi_t^y|^2 \mu(dx) \mu(dy) \\
&\leq \int_{\mathcal{H} \times \mathcal{H}} \left| \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t)] H_{\bar{n}}(x) \right|^2 \mu(dx) \mu(dy) \\
&\quad + \int_{\mathcal{H} \times \mathcal{H}} \left| \sum_{\bar{n} \in \mathcal{J}} u_{\bar{n}}^y(t) [H_{\bar{n}}(x) - H_{\bar{n}}(y)] \right|^2 \mu(dx) \mu(dy) \\
&\leq \int_{\mathcal{H} \times \mathcal{H}} \sum_{\bar{n} \in \mathcal{J}} |u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t)|^2 H_{\bar{n}}^2(x) \mu(dx) \mu(dy) \\
&\quad + \int_{\mathcal{H} \times \mathcal{H}} \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^y(t)]^2 \sum_{\bar{n} \in \mathcal{J}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \\
&= \sum_{\bar{n} \in \mathcal{J}} |u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t)|^2 \int_{\mathcal{H} \times \mathcal{H}} H_{\bar{n}}^2(x) \mu(dx) \mu(dy) \\
&\quad + \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^y(t)]^2 \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \\
&= \sum_{\bar{n} \in \mathcal{J}} |u_{\bar{n}}^x(t) - u_{\bar{n}}^y(t)|^2 + \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^y(t)]^2 \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \\
&= \int_{\mathcal{H} \times \mathcal{H}} \left| \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right|^2 \mu(dx) \mu(dy) \\
&\quad + \int_{\mathcal{H} \times \mathcal{H}} \mathbb{E}^2[\varphi(X_t^y)] \mu(dx) \mu(dy) \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \\
&\quad + \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^y(t)]^2 \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \quad (3.8)
\end{aligned}$$

Notice that  $\mathbb{E}^2[\varphi(X_t^y)] = u^2(t, x) \in L^2(\mathcal{H}, \mu)$  therefore the first integral in the second term is a continuous bounded function of  $t$ . Moreover,  $\sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^y(t)]^2$  is the  $L^2(\mathcal{H}, \mu)$ -norm of the function  $u(t, x)$  then the series converges and it is also a continuous bounded function of  $t$ . Thus, from (3.8) we get

$$\begin{aligned}
\|\Psi_t^x - \Psi_t^y\|_{(L^2(\mathcal{H}, \mu))^2}^2 &\leq \int_{\mathcal{H} \times \mathcal{H}} \left| \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right|^2 \mu(dx) \mu(dy) \\
&\quad + f(t) \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \quad (3.9)
\end{aligned}$$

where  $f(t) = \sum_{\bar{n} \in \mathcal{J}} [u_{\bar{n}}^y(t)]^2$ .

From the proof of Theorem 2.12 (see (2.23)) we know that

$$\begin{aligned}
\|\Psi_t^\varphi - \Phi_t^\psi\|^2 &= \int_{\mathcal{H} \times \mathcal{H}} \left| \mathbb{E}[\varphi(X_t^x)] - \mathbb{E}[\varphi(X_t^y)] \right|^2 \mu(dx) \mu(dy) \\
&\leq \exp(Ct) \int_{\mathcal{H} \times \mathcal{H}} \|x - y\|_{\mathcal{H}}^2 \mu(dx) \mu(dy)
\end{aligned}$$

### 3 CONTINUITY WITH RESPECT TO THE INITIAL CONDITIONS OF A NUMERICAL APPROXIMATIONS

$$= \exp(Ct) \|x - y\|^2. \quad (3.10)$$

Therefore the first term in the right side of (3.9) is bounded by (3.10).

We now focus on the second term in the last inequality. Notice that for every  $\bar{n} \in \mathcal{J}$  we have

$$H_{\bar{n}}(x) - H_{\bar{n}}(y) = \prod_{i=1}^{\infty} [P_{n_i}(\xi_i) - P_{n_i}(\eta_i)] \quad (3.11)$$

where  $\xi_i = \langle x, \Lambda^{-1/2} e_i \rangle_{\mathcal{H}}$  and  $\eta_i = \langle y, \Lambda^{-1/2} e_i \rangle_{\mathcal{H}}$  (see (2.3) and lines after that for the definition). By applying Lemma 3.1 to equation (3.11) we have that

$$\begin{aligned} H_{\bar{n}}(x) - H_{\bar{n}}(y) &= \prod_{i=1}^{\infty} C(i) P_{e_{i+1}}(\gamma_i) \cdot (\xi_i - \eta_i) \\ &= \prod_{i=1}^{\infty} C(i) P_{e_{i+1}}(\gamma_i) \langle x - y, \Lambda^{-1/2} e_i \rangle_{\mathcal{H}} \end{aligned} \quad (3.12)$$

here  $\gamma_i \in (\xi_i \wedge \eta_i, \xi_i \vee \eta_i)$  for every  $i \in \mathbb{N}$ . Then

$$\begin{aligned} \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) &= \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} \left| \prod_{i=1}^{\infty} C(i) P_{e_{i+1}}(\gamma_i) \langle x - y, \Lambda^{-1/2} e_i \rangle_{\mathcal{H}} \right|^2 \mu(dx) \mu(dy) \\ &= \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} \prod_{i=1}^{\infty} [C(i) P_{e_{i+1}}(\gamma_i)]^2 \left| \langle x - y, \Lambda^{-1/2} e_i \rangle_{\mathcal{H}} \right|^2 \mu(dx) \mu(dy) \\ &\leq \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} \prod_{i=1}^{\infty} [C(i) P_{e_{i+1}}(\gamma_i)]^2 \|x - y\|_{\mathcal{H}}^2 \|\Lambda^{-1/2} e_i\|_{\mathcal{H}}^2 \mu(dx) \mu(dy) \\ &= \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} \prod_{i=1}^{\infty} [C(i) P_{e_{i+1}}(\gamma_i)]^2 \|x - y\|_{\mathcal{H}}^2 \lambda_i^{-1} \|e_i\|_{\mathcal{H}}^2 \mu(dx) \mu(dy) \\ &= \|x - y\|_{\mathcal{H}}^2 \sum_{\bar{n} \in \mathcal{J}} \prod_{i=1}^{\infty} [C(i)]^2 \lambda_i^{-1} \int_{\mathcal{H} \times \mathcal{H}} [P_{e_{i+1}}(\gamma_i)]^2 \mu(dx) \mu(dy) \end{aligned} \quad (3.13)$$

Recall that for every  $i \in \mathbb{N}$  we have that  $\gamma_i \in (\xi_i \wedge \eta_i, \xi_i \vee \eta_i)$ , set  $\hat{\gamma}_i \in (\xi_i \wedge \eta_i, \xi_i \vee \eta_i)$  such that  $P_{e_i}^2(\gamma_i) \leq P_{e_{i+1}}^2(\hat{\gamma}_i)$  for every  $\gamma_i \in (\xi_i \wedge \eta_i, \xi_i \vee \eta_i)$ , notice that the existence of  $\hat{\gamma}_i$  is guaranteed since  $P_{e_{i+1}}^2(\cdot)$  is a continuous function. Then, from (3.13) we get

$$\begin{aligned} \sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) &\leq \|x - y\|_{\mathcal{H}}^2 \sum_{\bar{n} \in \mathcal{J}} \prod_{i=1}^{\infty} [C(i)]^2 \lambda_i^{-1} [P_{e_{i+1}}(\hat{\gamma}_i)]^2 \int_{\mathcal{H}} \int_{\mathcal{H}} \mu(dx) \mu(dy) \\ &= \|x - y\|_{\mathcal{H}}^2 \sum_{\bar{n} \in \mathcal{J}} \prod_{i=1}^{\infty} [C(i)]^2 \lambda_i^{-1} [P_{e_{i+1}}(\hat{\gamma}_i)]^2 \end{aligned} \quad (3.14)$$

Here, we recall that  $C(i) = \frac{(-1)^i}{(i+1)(i!)^{1/2}}$  then  $\frac{(-1)^i}{[(i+1)!]^{1/2}} P_{e_{i+1}}(\hat{\gamma}_i)$  is the normalized Hermite polynomial of  $i + 1$  degree evaluated on  $\hat{\gamma}_i$  which is bounded by a constant  $C$  for every  $i \in \mathbb{N}$ . Moreover, since

## REFERENCES

$\lambda_k < \lambda_{k+1} \rightarrow \infty$  then this implies that

$$\sum_{\bar{n} \in \mathcal{J}} \prod_{i=1}^{\infty} \left[ C(i) \right]^2 \lambda_i^{-1} \left[ P e_{i+1}(\hat{\gamma}_i) \right]^2 \leq C \sum_{\bar{n} \in \mathcal{J}} \prod_{i=1}^{\infty} \lambda_i^{-1} (i+1)^{-1} \leq C \quad (3.15)$$

where  $C$  is a finite constant. Putting together (3.13) and (3.15) we get that

$$\sum_{\bar{n} \in \mathcal{J}} \int_{\mathcal{H} \times \mathcal{H}} |H_{\bar{n}}(x) - H_{\bar{n}}(y)|^2 \mu(dx) \mu(dy) \leq C \|x - y\|_{\mathcal{H}} \quad (3.16)$$

Putting together inequalities (3.9), (3.10) and (3.16) we obtain

$$\|\Psi_t^x - \Psi_t^y\|_{(L^2(\mathcal{H}, \mu))^2}^2 \leq \exp(Ct) \int_{\mathcal{H} \times \mathcal{H}} \|x - y\|_{\mathcal{H}}^2 \mu(dx) \mu(dy) + f(t) \|x - y\|_{\mathcal{H}} \quad (3.17)$$

Now, if  $\|x - y\|_{\mathcal{H}} \leq \delta$  then from (3.17) we get

$$\|\Psi_t^x - \Psi_t^y\|_{(L^2(\mathcal{H}, \mu))^2}^2 \leq G(t) \epsilon \quad (3.18)$$

for some  $\epsilon > 0$ , which implies the theorem. The proof is complete.  $\square$

**Remark 3.3.** If we consider in addition the supremum norm on  $t$ , then from (3.17) we get

$$\begin{aligned} \sup_{0 \leq t \leq T} \|\Psi_t^x - \Psi_t^y\|_{(L^2(\mathcal{H}, \mu))^2}^2 &\leq C \|x - y\|_{\mathcal{H}}^2 \sup_{0 \leq t \leq T} f(t) \\ &\quad + \exp(CT) \int_{\mathcal{H} \times \mathcal{H}} \|x - y\|_{\mathcal{H}}^2 \mu(dx) \mu(dy). \end{aligned} \quad (3.19)$$

Notice that  $f(t)$  is differentiable and continuous, then  $\sup_{0 \leq t \leq T} f(t) \leq C$ , then from (3.19) we obtain

$$\sup_{0 \leq t \leq T} \|\Psi_t^x - \Psi_t^y\|_{(L^2(\mathcal{H}, \mu))^2}^2 \leq C \|x - y\|_{\mathcal{H}} + \exp(CT) \int_{\mathcal{H} \times \mathcal{H}} \|x - y\|_{\mathcal{H}}^2 \mu(dx) \mu(dy). \quad (3.20)$$

From this inequality it is possible to show the continuous dependence on the initial conditions for this norm.

## 4 Numerical experiments

### References

- [1] V. Barbu, G. Da Prato: *The Kolmogorov equation for a 2D Navier-Stokes stochastic flow in a channel*. nonlinear Analysis. 69, pp. 940-949. (2008).
- [2] V. Bogachev, G. Da Prato, M. Röeckner: *Existence results for Fokker-Planck equations in Hilbert space*. Seminar on Stochastic Analysis, Random Fields and Applications VI. Progress in Probability Volume 63, 2011, pp 23-35 (2010)
- [3] Cameron R.H., Martin W. T.: *The orthogonal development of non-linear functionals in series of Fourier-Hermite functionals*. Annals of Mathematics 48 (2): 385-392. (1947).
- [4] P.-L. Chow: *Infinte-dimensional Kolmogorov equations in Gauss-Sobolev spaces*. Stoch. Anal. Applic. 14, 257-282. (1996).
- [5] P.-L. Chow: *Infinite-dimensional parabolic equations in Gauss-Sobolev spaces*. Comm. Stoch. Anal. 1, 71-86. (2007).

## REFERENCES

---

- [6] P.-L. Chow: Stochastic partial differential equations. Chapman and Hall/CRC. (2007)
- [7] G. Da Prato: Kolmogorov equations for stochastic partial differential equations. Advanced Courses in Mathematics - CRM Barcelona. Birkhäuser. (2004).
- [8] G. Da Prato: An introduction to Kolmogorov equations in Hilbert spaces. Lecture notes. May 2011. (2011).
- [9] G. Da Prato, A. Debussche: *m-Dissipativity of Kolmogorov Operators Corresponding to Burgers Equations with Space-time White Noise*. Potential Analysis. Volume 26, Issue 1, pp 31-55. (2007).
- [10] G. Da Prato, J. Zabczyk: Second order partial differential equations in Hilbert spaces. Cambridge University Press, (2002).
- [11] G. Da Prato, J. Zabczyk: Stochastic equations in infinite dimensions. Cambridge University Press, (1992).
- [12] G. Da Prato, F. Flandoli, M. Röckner: *Fokker-Planck Equations for SPDE with Non-trace-class Noise*. Communications in Mathematics and Statistics. Volume 1, Issue 3, pp 281-304, (2013).
- [13] F.J. Delgado-Vences and F. Flandoli: *A spectral-based numerical method for Kolmogorov equations in Hilbert spaces*. Infinite Dimension analysis and Quantum Probability. Vol. 19, No. 3 (2016) 1650020 (37 pages) (DOI: 10.1142/S021902571650020X)
- [14] P. Imkeller: Malliavin's calculus and applications in stochastic control and finance IMPAN Lecture Notes, Vol. 1, Warsaw (2008).
- [15] J. L. Lions: *Quelques methodes de resolution des problemes aux limites non lineaires*, Dunod, Paris (1969).
- [16] A. Pazy: Semigroups of Linear Operators and Applications to Partial Differential Equations Springer-Verlag, New York/Berlin (1983)
- [17] M. Renardy, R.C Rogers: An Introduction to Partial Differential Equations. second edition. Springer. (2004).