

## **Ultimate Nutrition**

### **Beta Methoxy Caps**

5-Methy-7-Methoxyisoflavone, commonly referred to as Methoxyisoflavone or Methoxy, has shown the ability to decrease the effects of muscle breakdown by decreasing the bodies normal production of Cortisol after an intense workout. Methoxy boosts the bodies protein synthesis leading to a shorter post-workout recovery period. Additionally, Methoxy acts as a fuel source for the body and provides energy during workouts. The combination of Methoxy's protein synthesis and Cortisol reduction make it the ideal supplement for the athlete looking to reduce fat while increasing lean body mass.

Recent research studies for Methoxyisoflavone and Beta-Ecdysterone indicate:

- Increased protein synthesis
- Increased lean muscle mass
- Increased nitrogen retention
- Improved Nitrogen balance
- Increased working capacity

Ecdysterone is an anabolic compound from Russia that is derived from *Rhaponticum carhamoides*. This remarkable plant extract has been proven in numerous clinical studies to be more anabolic than methandrostenolone (Dianabol), with no androgenic or other undesirable side effects. Ecdysterone is a non-hormonal dietary supplement that will cause a huge increase in anabolism but will not effect hormone levels.

The Russian ingredients in Ecdysterone have also been clinically proven to increase in strength and endurance. This is the same exact ingredient that the Russian government gave to its athletes prior to competitions in order to optimize performance while coming off anabolic steroids to beat doping tests. Strangely enough, many Russian athletes performed better in competition while using Ecdysterone than they did in preparation while on anabolic steroids.

### **SELECTED REFERENCES**

Batkai L, Nogradi M, Farkas L, Feuer L, Horvath I. Related Articles Demethylation of 7-methoxyisoflavone by *Penicillium cyclopium*. *Arch Mikrobiol.* 1973 Mar 26;90(2):165-6.

Dobens L, Rudolph K, and Berger E Ecdysterone regulatory elements function as both transcriptional activators and repressors. *Mol. Cell. Biol.*, Apr 1991; 11(4): 1846-53.

Kraft C, Jenett-Siems K, Siems K, Gupta M, Bienzle U, and Eich E Antiplasmodial activity of isoflavones from *Andira inermis*. *J Ethnopharmacol*, Nov 2000; 73(1-2): 131-5.

Vafopoulou-Mandalos X and Laufer H Regulation of hemoglobin synthesis by

ecdysterone and juvenile hormone during development of *Chironomus thummi* (Diptera). *Differentiation*, Jan 1984; 27(2): 94-105.

Webster S Effects of exogenous ecdysterone upon moulting, proecdysial development, and limb regeneration in the prawn *Palaemon elegans*. *Gen Comp Endocrinol*, Mar 1983; 49(3): 459-69.

## FREQUENTLY ASKED QUESTIONS

### 1 What is Methoxyisoflavone & Ecdysterone?

5-Methyl-7-Methoxy-Isoflavone or Methoxyisoflavone-7-Isopropoxyisoflavone was developed in the early 1970's by a Hungarian company who originally used the compound as an animal feed additive to increase the lean body mass in cattle. Ecdysterone is a non-hormonal anabolic supplement whose ingredients have been proven in clinical studies to be as anabolic as actual steroids without exhibiting any undesirable side effects.

### 2 What are the benefits of taking Methoxyisoflavone & Ecdysterone?

World class athletes have reported excellent results in gaining massive amounts of lean body mass, increased recovery times and superior strength gains that correlate to the increased muscle mass. Methoxyisoflavone and Ecdysterone are directly responsible for an increase in nitrogen retention and increased muscle growth. Finally, research indicates that this combination produces very strong Cortisol-suppressing effects.

### 3 Has research provided any negative effects that are associated with similar anabolic compounds?

The active anabolic ingredient in Ecdysten is Ecdysterone. This remarkable plant extract has been proven in numerous clinical studies to be more anabolic than methandrostenolone (Dianabol), with no androgenic or other undesirable side effects. Ecdysterone is a non-hormonal dietary supplement that will cause a huge increase in anabolism but will not effect hormone levels.

### 4 How is Methoxyisoflavone and Ecdysterone taken?

According to research, Methoxyisoflavone and Ecdysterone are most effective when taken together and can be found in powder form, capsules, tablets or liquids. Studies performed on Methoxyisoflavone conclude that anabolic activity is present when taking between 300 - 500 mg per day. Subjects showed improvement in recuperation, with subsequent gains in strength. For maximum absorption into the body, the recommended dosage is approximately 40 mg of Ecdysterone plus 200 of Methoxyisoflavone twice per day.