

Analytic Wellfield Model

ALAN LEWIS¹

January 25, 2018

¹www.github.com/geouke/awfm

Contents

1	Mathematical Model	3
1.1	Forward Model	3
1.1.1	Recoverable Water Level	3
1.1.2	Aquifer Drawdown Model	3
1.1.3	Well-Loss Model	3
1.2	Parameter Estimation	3
1.2.1	Aquifer Parameters	3
1.2.2	Recoverable Water Level(s) and Well-Loss Coefficient(s)	3
2	Using the Graphical Interface	5
2.1	Creating a New Model	5
2.2	Handling Units	5
2.3	Data Importing	5
2.3.1	CSV	5
2.3.2	Excel	5
2.3.3	SQLite	5
2.4	Timeseries Processing	5
2.4.1	Scale	5
2.4.2	Translate	6
2.4.3	Erroneous/Missing Data	6
2.4.4	Project Onto Line	6
2.4.5	Range Constraints	6
2.4.6	Data Reduction	6
3	Examples	7
3.1	Simple Pumping Test	7
3.2	Analysis With Realistic Data	7

List of Figures

List of Tables

Introduction

“This is a quote and I don’t know who said this.”

– Author’s name, *Source of this quote*

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis risus ante, auctor et pulvinar non, posuere ac lacus. Praesent egestas nisi id metus rhoncus ac lobortis sem hendrerit. Etiam et sapien eget lectus interdum posuere sit amet ac urna.

1

Mathematical Model

1.1 Forward Model

Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat.

Duis aute irure dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

Lorem ipsum list:

1.1.1 Recoverable Water Level

1.1.2 Aquifer Drawdown Model

1.1.3 Well-Loss Model

1.2 Parameter Estimation

1.2.1 Aquifer Parameters

1.2.2 Recoverable Water Level(s) and Well-Loss Coefficient(s)

2

Using the Graphical Interface

2.1 Creating a New Model

2.2 Handling Units

The units dialog is used for specifying three unit types in the model: length, time, and discharge. The discharge unit applies to pumping rates ¹, while length and time units apply to everything else.

2.3 Data Importing

2.3.1 CSV

2.3.2 Excel

2.3.3 SQLite

2.4 Timeseries Processing

The timeseries dialog, which is used for importing and viewing pumping rates and observed water levels, has a handful of useful functions detailed below.

2.4.1 Scale

Scale here is used in the mathematical sense of multiplying a value by a scalar. This function is most useful for converting between units, but could

¹In reality, discharge may be expressed in units of length and time. Here, discharge units are specified separately to allow plotting in units familiar to the user.

also be used to model what-if scenarios. For example, it is possible to see the effect of increasing or decreasing discharge volumes by a percent.

2.4.2 Translate

Translation is also used in the mathematical sense of adding some scalar to a value. This function is useful if observed water levels at multiple wells were all taken relative to their own piezometer elevations and need to be converted to a consistent elevation.

Another application of the translation function is to adjust t_0 . Data imported from Excel may have a t_0 in the early 1900s, whereas it may be more useful to have a t_0 when production began at the wellfield.

2.4.3 Erroneous/Missing Data

The data import utility will set missing row values to -9999. These need to be dealt with before the forward model can produce meaningful results.

One option is to simply remove missing data. In the context of pumping rates, this means that the value at the missing data point is effectively equal to the value at the previously measured data point:

<insert table>

Another option is to perform a linear interpolation through the missing data point, which may be desirable for approximating missing water levels.

2.4.4 Project Onto Line

This operation performs a piece-wise linear interpolation, which is useful for data reduction or simply for obtaining a data set with a constant time step for input into a numerical model.

2.4.5 Range Constraints

2.4.6 Data Reduction

3

Examples

3.1 Simple Pumping Test

3.2 Analysis With Realistic Data