

Why LD Helps Us Find Selective Sweeps

Alan R. Rogers

March 13, 2020

LD helps us detect ongoing selective sweeps


This is curious, because we have known for over 50 years that selection at a single locus does not cause LD (Felsenstein 1965).

This lecture will explain why selective sweeps are *associated* with LD even though they don't cause it.


We begin with a graphical argument that shows why selection at a single locus doesn't cause LD.

Linkage equilibrium \iff shaded fractions equal

A-gametes



a-gametes




LE: Neither locus predicts other


Here, *B* is equally common among *a*-gametes and *A*-gametes.

Suppose allele A is favored; B/b are neutral

A -gametes



a -gametes



Selection inflates left circle.

Shrinks right one.

Shaded wedges unchanged.

System remains in LE.

If selection doesn't *cause* LD, why does LD help is detect selection?

D , the conventional measure of LD, doesn't help us here, because it *is* affected by selection—but not in a way that is helpful.

To explain this puzzle, I introduce another measure, which does not change in response to selection at a single locus.

Conditional allele frequencies

Gamete type	AB	Ab	aB	ab
Frequency	x_1	x_2	x_3	x_4

Freq of B among A -gametes

$$p_{B|A} = \frac{x_1}{x_1 + x_2}$$

The size of one pie slice.

Freq of B among a -gametes

$$p_{B|a} = \frac{x_3}{x_3 + x_4}$$

The size of the other slice.

Neither is affected by selection on A/a .

The Nei-Li measure of linkage disequilibrium

$$d = p_{B|A} - p_{B|a}$$

(Nei & Li, 1980)

Difference in frequency of B between A - and a -gametes.

Selection on A/a affects neither $p_{B|A}$, $p_{B|a}$, nor d .

Calculating D and d

Gamete	Locus	
	1	2
1	A	B
2	A	B
3	A	B
4	A	B
5	A	B
6	A	b
7	a	B
8	a	B
9	a	b
10	a	b

	A	a	
B	5	2	7
b	1	2	3
	6	4	10

$$D = \frac{5}{10} \cdot \frac{2}{10} - \frac{1}{10} \cdot \frac{2}{10} = \frac{2}{25}$$

$$d = p_{B|A} - p_{B|a} = \frac{5}{6} - \frac{1}{2} = \frac{1}{3}$$

D depends on heterozygosity at locus A .

$$\begin{aligned}d &= \frac{x_1}{x_1 + x_2} - \frac{x_3}{x_3 + x_4} \\&= \frac{D}{p_A(1 - p_A)}\end{aligned}$$

$$D = dp_A(1 - p_A)$$

Selection at locus A affects $p_A(1 - p_A)$ and therefore D .

LD of a new mutant


When allele A first arises by mutation, it will be rare, so $p_A(1 - p_A) \approx 0$ and $D \approx 0$.

But d may be large: it is either $1 - p_B$ or $-p_B$, depending on whether the mutant gamete is AB or Ab .


The value of d stays large if selection outruns recombination and drift.

This is why LD helps us detect selection.


Selective sweep of A , with $s = 0.02$, $c = 0.001$, and $2N = 50,000$



Neutral drift to fixation, $c = 0.001$, and $2N = 50,000$



Selective sweep in small population ($s = 0.02$, $c = 0.001$, $2N = 5000$)



Summary

- ▶ LD decays gradually in response to recombination.
- ▶ This is more obvious for d than for D , because d is insensitive to p_A .
- ▶ Advantageous alleles increase rapidly: there is little time for LD to decay.
- ▶ Neutral alleles increase slowly; plenty of time for LD to decay.