

Population Structure with Genomic Data

Alan R. Rogers

March 30, 2020


Shared blocks of identity by descent (IBD)

- With molecular data, we can identify shared IBD blocks.
- Close relatives tend to share long blocks.
- Distant relatives share short blocks.


1 / 12

2 / 12

IBD sharing between my mother and daughter



IBD sharing with French-speaking Swiss and UK



Each dot an individual. Color shows individual's origin. (FR, France; IT, Italy; EL, Greece; TR, Turkey, CY, Cyprus)

Those who share ancestry with Swiss also share ancestry with UK.


Presumably reflects immigration from ancestors of Swiss and British.

Ralph and Coop (2013)

3 / 12

4 / 12

Sharing with Irish and Germans




Each dot an individual. Pink, Germans; Green, British; Blue, Irish.

Brits with lots of German ancestry have little Irish ancestry.

British population is a mixture of Celts and Germans.

If we focused on shorter IBD blocks, we'd be studying a different time scale, and the pattern might be different.


Geographic decay of recent relatedness




Genetic similarity versus geographic distance. Small dots are pairs of individuals. E, Eastern Europe; W, Western Europe; N, Northern Europe; I, Italy & Iberia; TC, Turkey & Cyprus. (Ralph & Coop 2013)

5 / 12


6 / 12




7 / 12




8 / 12



9 / 12



10 / 12



11 / 12

Summary

- By distinguishing between long and short IBD blocks, we can examine geographic population structure on different time scales.

12 / 12