






1
point

1. Consider the following 2D dataset:



Which of the following figures correspond to possible values that PCA may return for $u^{(1)}$ (the first eigenvector / first principal component)? Check all that apply (you may have to check more than one figure).

☐

1
point

2. Which of the following is a reasonable way to select the number of principal components k ?

(Recall that n is the dimensionality of the input data and m is the number of input examples.)

- ☒ Choose k to be the smallest value so that at least 99% of the variance is retained.
- ☐ Choose k to be the smallest value so that at least 95% of the variance is retained.
- ☐ Choose k to be 99% of n (i.e., $k = 0.99 * n$, rounded to the nearest integer).
- ☐ Choose the value of k that minimizes the approximation error $\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}^{(i)}\|^2$.

Principal Component Analysis

Quiz, 5 questions

1
point

3. Suppose someone tells you that they ran PCA in such a way that "95% of the variance was retained." What is an equivalent statement to this?

- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}^{(i)}\|^2} \geq 0.05$
- ☒ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}^{(i)}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \leq 0.05$
- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}^{(i)}\|^2} \leq 0.95$
- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}^{(i)}\|^2} \leq 0.05$

1
point

4. Which of the following statements are true? Check all that apply.

- ☒ If the input features are on very different scales, it is a good idea to perform feature scaling before applying PCA.
- ☐ PCA can be used only to reduce the dimensionality of data by 1 (such as 3D to 2D, or 2D to 1D).

- ☒ Given an input $x \in \mathbb{R}^n$, PCA compresses it to a lower-dimensional vector $z \in \mathbb{R}^k$.

- ☐ Feature scaling is not useful for PCA, since the eigenvector calculation (such as using Octave's `svd(Sigma)` routine) takes care of this automatically.

1
point

5. Which of the following are recommended applications of PCA? Select all that apply.

- ☐ Clustering: To automatically group examples into coherent groups.
- ☒ Data compression: Reduce the dimension of your input data $x^{(i)}$, which will be used in a supervised learning algorithm (i.e., use PCA so that your supervised learning algorithm runs faster).
- ☒ Data visualization: Reduce data to 2D (or 3D) so that it can be plotted.
- ☐ To get more features to feed into a learning algorithm.

- ☒ I understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account.
Learn more about Coursera's Honor Code

Alan Ross

Submit Quiz

