



Taylor & Francis
Taylor & Francis Group

A GIANT EARLY PLEISTOCENE BIRD FROM EASTERN EUROPE

Author(s): NIKITA V. ZELENKOV, ALEXANDER V. LAVROV, DMITRY B. STARTSEV, INNESSA A. VISLOBOKOVA and ALEXEY V. LOPATIN

Source: *Journal of Vertebrate Paleontology*, March 2019, Vol. 39, No. 2 (March 2019), pp. 1-7

Published by: Taylor & Francis, Ltd. on behalf of The Society of Vertebrate Paleontology

Stable URL: <https://www.jstor.org/stable/10.2307/26776730>

REFERENCES

Linked references are available on JSTOR for this article:

https://www.jstor.org/stable/10.2307/26776730?seq=1&cid=pdf-reference#references_tab_contents

You may need to log in to JSTOR to access the linked references.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

The Society of Vertebrate Paleontology and Taylor & Francis, Ltd. are collaborating with JSTOR to digitize, preserve and extend access to *Journal of Vertebrate Paleontology*

A GIANT EARLY PLEISTOCENE BIRD FROM EASTERN EUROPE: UNEXPECTED COMPONENT OF TERRESTRIAL FAUNAS AT THE TIME OF EARLY *HOMO* ARRIVAL

NIKITA V. ZELENKOV,^{*,1} ALEXANDER V. LAVROV,¹ DMITRY B. STARTSEV,² INNESSA A. VISLOBOKOVA,¹ and ALEXEY V. LOPATIN^{1,3}

¹Borissiak Paleontological Institute of Russian Academy of Sciences, Profsoyuznaya 123, 117647 Moscow, Russian Federation, nzelen@paleo.ru;

²Vernadsky Crimean Federal University, Akademika Vernadskogo 4, 95000 Simferopol, Crimea;

³Lomonosov Moscow State University, Leninskie Gory 1, 119991 Moscow, Russian Federation

ABSTRACT—Giant birds, comparable in size to elephant birds and moa, have never been reported from Europe. Here, we describe a femur from the lower Pleistocene of the north Black Sea area (Crimea) that is referred to *Pachystruthio dmanisensis*, comb. nov., a giant bird with an estimated body mass of about 450 kg. This value makes this extinct bird one of the largest known avians (comparable to *Aepyornis maximus*) and the only bird of such giant size in Europe and the Northern Hemisphere in general. In contrast to very large insular birds, *Pachystruthio dmanisensis* was a good runner, which may be explained by its coexistence with large carnivorous mammals. *Pachystruthio dmanisensis* and associated assemblage of fossil mammals are shared with the Dmanisi locality in Georgia (~1.8–1.7 Ma); thus, this giant bird was likely a typical component of eastern European faunas at the time of early hominin arrival. We suggest that *Pachystruthio dmanisensis*, together with early *Homo* and a variety of mammals, reached the northern Black Sea region via the southern Caucasus and Anatolia, because the older (Pliocene) finds of this fauna are known from Georgia and Turkey.

Citation for this article: Zelenkov, N. V., A. V. Lavrov, D. B. Startsev, I. A. Vislobokova, and A. V. Lopatin. 2019. A giant early Pleistocene bird from eastern Europe: unexpected component of terrestrial faunas at the time of early *Homo* arrival. Journal of Vertebrate Paleontology. DOI: 10.1080/02724634.2019.1605521.

INTRODUCTION

The Cenozoic diversity of flightless, large-bodied ground birds was higher than at present, and several now extinct groups include notably large representatives (Angst and Buffetaut, 2017). The true giants of the avian world were the largest species of mihirungs (Dromornithidae) from the Miocene of Australia and elephant birds (Aepyornithidae) from the late Quaternary of Madagascar, with an estimated body mass of up to 700 kg (Murray and Vickers-Rich, 2004; Hansford and Turvey, 2018; but see below). The size of these birds greatly exceeded that of the largest living bird, the common ostrich (*Struthio camelus*), which weighs up to 150 kg (Folch, 1992).

Fossil birds of such an impressively giant size have never been documented from Europe or the Northern Hemisphere in general (Angst and Buffetaut, 2017). The North American late Quaternary giant bird *Titanis walleri* Brodkorb, 1963, weighed about 150 kg (Baskin, 1995), and the Paleogene Gastornithidae most probably weighed no more than 200 kg (Angst and Buffetaut, 2017). Although several large ostriches (see, however, the discussion concerning their taxonomic affinities) have been described from Neogene to Quaternary sediments of Eurasia (Boev and Spassov, 2009; Mayr, 2017), they were never considered to be as large as elephant birds or the New Zealand moa. Here, we report a femur of a very large bird from the early Pleistocene of the Crimean Peninsula (north Black Sea region). The estimated body mass of this bird, calculated in accordance with the formulae widely used for fossil giant birds

(Campbell and Marcus, 1992; Worthy and Holdaway, 2002; Field et al., 2013; Handley et al., 2016), is about 450 kg. This calculated value is comparable to that of *Aepyornis maximus* and thus makes this eastern European giant bird one of the largest (heaviest) known avians, being smaller only than the late Miocene Australian *Dromornis stirtoni* and the Quaternary aepyornithid *Vorombe titan* (Murray and Vickers-Rich, 2004; Handley et al., 2016; Hansford and Turvey, 2018). We further propose a revision of the previously named late Cenozoic ostrich-like birds from eastern Europe and highlight a previously unrecognized wide distribution of giant birds in this region at the time of the arrival of early *Homo* in the early Pleistocene.

Institutional Abbreviations—**D**, collection of the Institute of Paleobiology of the National Museum of Georgia, Tbilisi, Georgia; **PI**, Borissiak Paleontological Institute of the Russian Academy of Sciences, Moscow, Russia.

LOCALITY

The specimen comes from the recently discovered Taurida Cave in the Crimean Peninsula, north Black Sea region (Fig. 1; see Lopatin et al., 2019, for details on geology and fauna). The associated late Villafranchian fauna of the locality comprises a variety of mammals, including representatives of the genera *Archidiskodon*, *Arvernoceros*, *Bison* (*Eobison*), *Canis*, *Elasmotherium*, *Equus*, *Leptobos*, *Gazellospira*, *Homotherium*, *Hypolagus*, *Hystrix* (*Acanthion*), *Pachycrocuta*, *Paracamelus*, *Pontoceros*, and *Stephanorhinus* (see Lopatin et al., 2019). Associated bird remains include the extinct Villafranchian bustard *Tetrax kalmani* Jánossy, 1972.

The association of *Leptobos* and *Eobison* indicates an age of 1.8–1.5 Ma. The assemblage of fossil mammals (and birds, e.g.,

*Corresponding author.

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/ujvp.




FIGURE 1. Map showing the geographical location of the fossil locality and the cave plan.

Pachystruthio), however, is substantially similar to that of the Dmanisi locality, which indicates an age close to 1.8–1.7 Ma (Lopatin et al., 2019).

SYSTEMATIC PALEONTOLOGY

Order ?STRUTHIONIFORMES

Family INCERTAE SEDIS

Genus *PACHYSTRUTHIO* Kretzoi, 1954

?*PACHYSTRUTHIO* *DMANISENSIS* (Burchak-Abramovich and Vekua, 1990), comb. nov.
(Fig. 2A, C, E, F)

Holotype—D70, a right femur (Vekua, 2013).

Type Locality—Dmanisi, Georgia; lower Pleistocene (1.85–1.78 Ma; Ferring et al., 2011).

Referred Specimen—PIN 5644/56, a nearly complete left femur from the Taurida Cave, Crimean Peninsula (Fig. 1); early Pleistocene (late Villafranchian).

Remarks—The distinctive morphology of PIN 5644/56 (see description below) precludes assigning this giant bird to the genus *Struthio*. Furthermore, PIN 5644/56 corresponds well with '*Struthio*' *dmanisensis* Burchak-Abramovich and Vekua, 1990, from the early Pleistocene of Georgia and can be referred to this extinct species. In addition to its very large size, the highly distinctive feature shared by the new specimen and '*S.*' *dmanisensis* is the exceptionally robust femoral shaft. The robustness index (minimal lateromedial shaft width divided by total length, expressed as a percentage) is 20% in the new specimen and '*S.*' *dmanisensis* from Dmanisi (Burchak-Abramovich and Vekua, 1990). This index is 12.8–16.4 ($n=11$) in modern *Struthio camelus* (Burchak-Abramovich, 1953; Burchak-Abramovich and Vekua, 1990; Vekua, 2013; this study). In the very large extinct ostriches *S. oldawayi* Lowe, 1933, and *S. karatheodori* Forsyth Major, 1888, this index also does not exceed 16 (Burchak-Abramovich and Vekua, 1990). Referral to '*S.*'

dmanisensis is further supported by the general faunal similarity between the localities of Dmanisi and Taurida (see above).

Other named taxa of very large birds, which show remarkable genus-level osteological differences from modern ostriches, are the late Pliocene '*Struthio*' *transcausicus* Burchak-Abramovich and Vekua, 1971, and the early Pleistocene *Struthio* (*Pachystruthio*) *pannonicus* Kretzoi, 1954. The latter species was described based on a remarkably large and morphologically distinct pedal phalanx from the Kisláng locality in Hungary (Kretzoi, 1954). The fossil assemblage from Kisláng has a mixture of Pliocene–early Pleistocene taxa, dated 3.5–1.7 Ma, but the absolute majority of small mammals belong to biozones MN17–MQ1, 2.0–1.6 Ma (Mayhew, 2012). This date corresponds well with the age of the Dmanisi locality (1.85–1.78 Ma; Ferring et al., 2011). The older '*S.*' *transcausicus* was described based on a pelvis from the late Pliocene (MN16b; 3.1–3.0 Ma; Bukhsianidze and Kohiava, 2018) of Georgia (Burchak-Abramovich and Vekua, 1971). The pelvis of '*S.*' *transcausicus* also differs morphologically from that of ostriches (Burchak-Abramovich and Vekua, 1971). These three late Cenozoic eastern European taxa are thus giant birds that are notably different from modern and extinct *Struthio* in morphology. Hence, given the close ages of the localities, absolute size similarities, and generally low diversity of giant flightless birds in continental deposits (and in Europe in particular), it is plausible to classify '*S.*' *dmanisensis*, '*S.*' *transcausicus*, and '*S.*' (*P.*) *pannonicus* within the genus *Pachystruthio*, although the specific identity of these taxa remains to be confirmed. Importantly, the morphological distinctiveness of these eastern European Pleistocene giant birds cannot be explained simply by their large size, because the very large *S. oldawayi* from Tanzania is morphologically similar to modern *Struthio* and has a considerably thinner femoral shaft (Table 1; Lowe, 1933; Burchak-Abramovich and Vekua, 1990). Affinities of *Pachystruthio* are unclear, and its referral to the order Struthioniformes remains to be confirmed.

The separate generic status of *Pachystruthio* is supported by the presence of a morphologically distinct and very thick (2.9–3.3 mm) aepeyornithoid eggshell in the late Pliocene–early

Pleistocene of Azerbaijan (Burchak-Abramovich, 1953; Mikhailov and Kurochkin, 1988), which was associated with bone remains of ‘*S.*’ *transcaucasicus* (see Burchak-Abramovich and Vekua, 1971). Eggshell associated with the holotype of *P. pannonicus* is also much thicker (2.6–3.4 mm) than in modern ostriches and, unlike ostriches, has pore canals, which form irregular longitudinal channels (Kretzoi, 1954).

The evolutionary lineage of these giant early Pleistocene birds can probably be traced back to the early Pliocene and even Miocene of Anatolia: a very large femur from the early Pliocene of the Çalta locality in Turkey (Jannoo and Sen, 1998) may belong to *Pachystruthio*. The extinct species *Struthio karatheodoris* was described based on a slightly smaller, but still very large femur from the late Miocene (Maeotian) of Samos Island (Martin, 1903), which was a part of the Anatolian mainland at that time (Popov et al., 2004). This specimen notably differs in proportions from modern and extinct ostriches (Martin, 1903) and thus could belong to this lineage as well. Sauer (1979) further mentioned thick aepyornithoid eggshells in the middle Miocene Çandır locality in Turkey, which he attributed to the same evolutionary lineage as *P. pannonicus*.

In addition to *Pachystruthio*, ostriches similar to members of the modern genus *Struthio* are also present in the latest Cenozoic (Plio-Pleistocene) fossil record of eastern Europe (Table 2). They are represented by an unnamed species from the early Pliocene (MN15) of the Odessa catacombs in southern Ukraine (Burchak-Abramovich, 1953) and an ostrich from the early Pleistocene (MN17) of the Liventsovka locality (Sea of Azov area, Russia), which was referred to *Struthio asiaticus* Milne-Edwards, 1871, by Kurochkin and Lungu (1970). These forms are smaller than the species of *Pachystruthio* (Table 1), although they are somewhat larger than the living *S. camelus* and the Pliocene Siwalik *S. asiaticus*. The specimens from the Odessa catacombs and Liventsovka may represent one species; Mikhailov (1988) had referred a large egg from the Neogene (probably late Pliocene or early Pleistocene; Burchak-Abramovich and Vekua, 1971) of southern Ukraine to *Struthiolithus chersonensis* (Brandt, 1873). The eggshell thickness in *S. chersonensis* ranges from 2.6 to 2.9 mm, which is comparable to that of the Odessan ostrich (2.7–3.3 mm; Burchak-Abramovich, 1953), but the specimens’ referral to one species is unclear. The early–middle Villafranchian European ostrich has not been recorded in the younger deposits of the north Black sea region and thus most likely was replaced in the late Villafranchian by the larger, southern migrant *Pachystruthio*.

DESCRIPTION

Specimen PIN 5644/56 (Fig. 2A, C, E, F) has a completely preserved shaft and partly damaged proximal and distal ends. The general proportions of the specimen are elongate as in modern *Struthio* and smaller taxa of moa and Phorusrhacidae, but in contrast to larger species of Aepyornithidae, Dromornithidae, and Emeidae, which have very robust femora with a strongly concave medial margin of the shaft. Such a concavity of the medial margin is poorly pronounced in PIN 5644/56, but it is still better expressed than in modern *Struthio*, which have subparallel medial and lateral margins of the femoral shaft.

A large part of the facies articularis antitrochanterica is preserved, indicating that the trochanter femoris was not prominent proximally, as in modern *Struthio* and in contrast to Aepyornithidae. The pneumatic foramen on the caudal surface of the proximal end is present but small (as in ‘*S.*’ *dmanisensis*), not contacting the articular surface of the caput femoris. On the caudal surface of the shaft, there is a well-developed linea intermuscularis caudalis, which bifurcates proximally, as in *Struthio*. In contrast to modern ostriches, this line is more centrally located (more medially in *Struthio*) and becomes indiscernible

TABLE 1. Basic measurements (in mm) of the femur in *Pachystruthio pannonicus* and selected species of *Struthio*.

Dimension	<i>?Pachystruthio dmanisensis</i> , PIN 5644/56, early Pleistocene, Taurida Cave	<i>?Pachystruthio dmanisensis</i> (n = 2), early Pleistocene, Dmanisi; after Burchak-Abramovich and Vekua (1990) and Vekua (2013)	<i>Struthio oldawayi</i> , early Pleistocene, Olduvai Gorge; after Burchak-Abramovich and Vekua (1990)	<i>Struthio karatheodoris</i> , late Miocene, Samos; after Martin (1903)	<i>Struthio</i> sp., early Pleistocene (Gelasian), Liventsovka; this study	<i>Struthio</i> sp., early Pliocene, Odessa Catacombs; after Burchak-Abramovich (1953)	<i>Struthio camelus</i> (n = 11), Recent; after Burchak-Abramovich (1953), Burchak-Abramovich and Vekua (1990), and our data	<i>Struthio molybdophanes</i> , Recent; after Burchak-Abramovich (1953)
Greatest (lateral) length	~390	~380, 385	~400	360	—	>342	294–337	279
Minimal shaft height (oblique craniocaudal)	74.7	76, 76	~65	—	63.5	67	42–57	46
Minimal shaft width (mediolateral)	59.3	54	—	55	—	49	33–47	33

TABLE 2. Described or figured bone remains of fossil *Struthio* and ostrich-like birds from the Pliocene and the Pleistocene of Europe, Turkey, and Transcaucasia.

Original assignment	Geography	Locality	Age	Osteological material	Reference	Current taxonomic assignment (this paper)
? <i>Pachystruthio dmanisensis</i>	Crimea	Taurida	Early Pleistocene (1.8–1.5 Ma)	Femur	This paper	? <i>Pachystruthio dmanisensis</i>
<i>Struthio dmanisensis</i>	Georgia	Dmanisi	Early Pleistocene (1.85–1.78 Ma)	Several femora	Burchak-Abramovich and Vekua, 1990; Vekua, 2013; Zelenkov and Kurochkin, 2015	? <i>Pachystruthio dmanisensis</i> , comb. nov.
<i>Struthio</i> (<i>Pachystruthio</i>) <i>pannonicus</i>	Hungary	Kisláng	Early Pleistocene (2.0–1.6 Ma)	Pedal phalanx	Kretzoi, 1954	<i>Pachystruthio pannonicus</i>
<i>Struthio asiaticus</i>	Russia, Rostov-on-Don region	Liventsovka	Early Pleistocene (~2.6–2.0 Ma)	Various leg bones	Kurochkin and Lungu, 1970; Baigusheva, 1971; Zelenkov and Kurochkin, 2015	<i>Struthio</i> sp.
<i>Struthio transcaucasicus</i>	Georgia	Kvabebi	Late Pliocene (3.2–2.5 Ma)	Pelvis	Burchak-Abramovich and Vekua, 1971; Zelenkov and Kurochkin, 2015	? <i>Pachystruthio transcaucasicus</i> , comb. nov.
<i>Struthio</i> sp.	Turkey	Çalta	Early Pliocene (4.2–3.6 Ma)	Various leg bones (femora, tibiotarsus, tarsometatarsi)	Janoo and Sen, 1998	? <i>Pachystruthio</i> sp.
<i>Struthio</i> sp.	Ukraine	Odessa catacombs	Early Pliocene (3.8–3.7 Ma)	Many postcranial bones	Burchak-Abramovich, 1952; Zelenkov and Kurochkin, 2015	<i>Struthio</i> sp.




FIGURE 2. Fossil femora of an extinct giant bird from the Crimean Peninsula (eastern Europe), with that of an average Recent ostrich for comparison. **A, C, E, F**, *Pachystruthio dmanisensis*, comb. nov., specimen PIN 5644/56, from Taurida Cave, Crimean Peninsula (early Pleistocene); **B, D**, *Struthio camelus*, osteological collection of PIN 1741-1. **A, B**, cranial view; **C, D**, caudal view; **E**, lateral view; **F**, proximal view. **Abbreviations:** cf, caput femoris; cl, condylus lateralis; cm, condylus medialis; ct, crista trochanteris; faa, facies articularis antitrochanterica; fop, fossa poplitea; fp, foramen pneumaticum; li, linea intermuscularis; sic, sulcus intercondylaris; sp, sulcus patellaris; tfi, trochlea fibularis.

FIGURE 3. Map showing distribution of bony remains of the giant species of *Pachystruthio* (large ostrich silhouettes) and smaller *Struthio* ostriches (small ostrich silhouettes) in the Black Sea region in the Pliocene (5.3–2.6 Ma; blue), Gelasian (2.6–1.8 Ma; green), and Calabrian (1.8–0.8 Ma; brown). *Pachystruthio* from Hungary (Kretzoi, 1954) is not shown. Localities: 1, Odessa catacombs (early Pliocene); 2, Kvabebi (late Pliocene); 3, Liventsovka (Gelasian); 4, Taurida Cave (Calabrian); 5, Dmanisi (Calabrian).



distally, near the mid-length of the bone. In *Struthio*, this line bifurcates distally, thus producing a prominent lateral branch, which extends distolaterally. In the fossil specimen, the lateral branch is indistinct.

The linea intermuscularis cranialis, connecting the distal part of the crista trochanteris and the proximal part of the lateral crest of the sulcus patellaris, is placed more medially than in modern *Struthio* and has a medial curvature near the mid-shaft that is absent in modern *Struthio*.

Distally, the fossa poplitea is notably shallower than in *Struthio* and does not extend far proximally. The proximal margin of the condylus medialis, bordering this sulcus caudally, is more steeply inclined than in modern *Struthio*. The sulcus intercondylaris, which is preserved between the damaged condylae, is more obliquely oriented. The lateral side of the sulcus curves distally steeply, which may indicate a strongly distally protruding condylus lateralis, as in Aepyornithidae. A proximal extension of the trochlea fibularis, which is well pronounced in modern *Struthio* and protrudes more proximally than the condylus lateralis, is poorly developed in the fossil. This crest is positioned approximately level with the proximal margin of the condylus lateralis. The lateral surface of the distal end, just proximal and dorsal to the epicondylus lateralis, is nearly flat and is not concave as in *Struthio*. In dorsal view, the distal end protrudes only slightly laterally relative to the shaft, in contrast to *Struthio*.

WEIGHT ESTIMATION

The body mass of a bird may be reconstructed based on femoral measurements using several empirical formulae, including the following ones that are widely used (see discussion in Handley et al., 2016) for large avian taxa: $\log_{10}M = 2.411 \times \log_{10}LCF - 0.065$ (Campbell and Marcus, 1992) and $\ln M = 2.4 \times \ln Fc - 0.11$ (Field et al., 2013), in which M is body mass and LCF and Fc are the least shaft circumference and the central shaft circumference, respectively. These measurements equal 240 mm in PIN 5644/56 and are actually the same in this specimen, because the mid-shaft is the thinnest part of the femur. The body mass of PIN 5644/56, calculated with these formulae, equals 462 and 453 kg, respectively.

The regressions used are based on modern avian diversity, which does not include such large-bodied forms; thus, the resulting calculations should be taken as an approximation. These

formulae would likely provide overestimated results when applied to morphologically strongly modified taxa with unusual limb proportions, for example, large species of moa (see Brassey et al., 2013) and aepyornithids. Accordingly, we note that the femur of *Pachystruthio* is generally similar to that of *Struthio* in overall morphology; thus, applying linear regressions to this taxon would likely provide a more reliable estimation.

DISCUSSION

The new fossil and mass estimations show that eastern Europe (Hungary through northern Caucasus) was inhabited in the early Pleistocene by some of the largest known giant birds, albeit of unclear taxonomic affinities. We tentatively assign them to the extinct genus *Pachystruthio*, although *P. pannonicus* from Hungary and ‘*S.*’ *dmanisensis* from Georgia and Crimea are based on different skeletal elements and thus direct comparisons of these two species is not yet possible. However, these similarly sized taxa are known from roughly contemporaneous localities and thus might even represent one species, which in this case would be the only bird of such a formidable size in the Northern Hemisphere.

Although similar in size to elephant birds, *Pachystruthio* differs from *Aepyornis maximus* and other large birds in having a relatively gracile and elongate femur, comparable to (although notably larger than) that of modern ostriches and smaller species of moa, aepyornithids, and phorusrhacids. The overall similarity to the modern ostrich indicates that *Pachystruthio* was a better runner than the graviportal giant species of aepyornithids. This distinctive morphological feature may be explained by the continental distribution of *Pachystruthio*. In contrast to other giant birds, *Pachystruthio* lived with a variety of highly specialized large carnivorous mammals, such as the giant cheetah (*Actionyx pardinensis*), giant hyenas (*Pachycrocuta brevirostris*), and saber-toothed cats (*Homotherium* spp., *Meganteron* spp.). It has been shown that large animals, including such massive forms as young *Mammuthus*, constituted a significant part of the prey of some early Pleistocene carnivorous mammals, especially *Homotherium* (Palmqvist et al., 2003). However, the great body mass of *Pachystruthio* suggests that it might not have been as good a runner as modern ostriches. Because of their huge size, adult individuals of *Pachystruthio* were obviously

less vulnerable than young individuals and smaller *Struthio* ostriches.

The appearance of the giant *Pachystruthio* birds in the late Villafranchian of eastern Europe coincides with a notable reorganization of the Eurasian terrestrial mammal faunas, which occurred at around the Olduvai subchron (former Pliocene–Pleistocene boundary; Rook and Martínez-Navarro, 2010; Vislobokova and Tesakov, 2013). In the northern Caucasus, the late Villafranchian (early Apscheronian) stage is characterized by an increased aridity and further expansion of open environments, such as steppes and semideserts, which first became widespread in this region in the middle Villafranchian (Naidina and Richards, 2016). In the late Villafranchian, several southern faunal elements, such as *Pachystruthio* and a number of extinct mammals, reached the north Black Sea region from Transcaucasia, as evident from the faunal similarity between the Taurida and Dmanisi localities (Lopatin et al., 2019), from where the oldest European skeletal remains of *Homo* are known (Ferring et al., 2011; see also Spassov, 2016). The presence of *Pachystruthio* in several early Pleistocene sites in eastern Europe (Fig. 3) indicates that these giant birds were a characteristic, although previously neglected, component of the terrestrial faunas at the time of early *Homo* arrival. These large birds might have been a source of meat, bones, feathers, and eggshell for early hominin populations.

It has been shown that a larger body mass is energetically advantageous in utilizing tougher, low-nutrition food due to decreased mass-specific metabolic demands in larger animals ('Jarman-Bell Principle'; see Müller et al., 2013). Murray and Vickers-Rich (2004) relate the evolutionary increase of body mass in mihirungs (Dromornithidae) to an increasing aridity in Australia during the Neogene–Pleistocene. This concept may also be applied to the large flightless birds of continental northern Eurasia, such as *Pachystruthio* and the Eocene–Pliocene Eogruidae (Ergilornithidae). The latter also became progressively larger as aridity increased in the Neogene of eastern Europe and Central Asia (Kurochkin, 1981; Zelenkov et al., 2016). The very large body mass (more than 250 kg; Buffetaut and Angst, 2017) of the late Pleistocene Chinese ostrich *S. anderssoni* may also be explained by an increase in aridity in Inner Asia during the cold climatic phases of the glacial periods (Lu et al., 2013).

ACKNOWLEDGMENTS

The authors thank K. K. Tarasenko for his assistance during field work. The authors are further indebted to T. Worthy, E. Buffetaut, Z. Boev, and editor V. De Pietri for valuable comments on the early version of this paper. Research was supported by the Russian Science Foundation, project 18-74-10081.

LITERATURE CITED

- Angst, D., and E. Buffetaut. 2017. Paleobiology of Giant Flightless Birds. ISTE Press–Elsevier, London, 281 pp.
- Baigusheva, V. S. 1971. Fossil theriofauna of the Liventzovka sand-pit; pp. 5–29 in N. K. Vereschagin (ed.), Materials on the Anthropogene faunas of USSR. Transactions of the Zoological institute of Russian Academy of Sciences. Volume 49. Nauka, Leningrad.
- Baskin, J. A. 1995. The giant flightless bird *Titanis walleri* (Aves: Phorusrhacidae) from the Pleistocene coastal plain of south Texas. Journal of Vertebrate Paleontology 15:842–844.
- Boev, Z. N., and N. Spassov. 2009. First record of ostriches (Aves, Struthioniformes, Struthionidae) from the late Miocene of Bulgaria with taxonomic and zoogeographic discussion. Geodiversitas 31:493–507.
- Brandt, A. 1873. Über ein grosses fossiles Vogelei aus der Umgegend von Cherson. Bulletin de l'Académie Impériale de Sciences de St. Petersburg 18:158–161.
- Brassey, C. A., R. N. Holdaway, A. G. Packham, J. Anné, P. L. Manning, and W. I. Sellers. 2013. More than one way of being a Moa: differences in leg bone robustness map divergent evolutionary trajectories in Dinornithidae and Emeidae (Dinornithiformes). PLoS ONE 8:e82668.
- Brodtkorb, P. 1963. A giant flightless bird from the Pleistocene of Florida. The Auk 80:111–115.
- Buffetaut, E., and D. Angst. 2017. How large was the giant ostrich of China? Evolução, Revista de Geostória e Pré-História 2:6–8.
- Bukhsianidze, M., and K. Koiava. 2018. Synopsis of the terrestrial vertebrate faunas from the Middle Kura Basin (Eastern Georgia and Western Azerbaijan, South Caucasus). Acta Palaeontologica Polonica 63:441–461.
- Burchak-Abramovich, N. I. 1953. Fossil Ostriches of Caucasus and Southern Ukraine. Azerbaijan SSR Academy of Science, Baku, Azerbaijan, 206 pp. [Russian]
- Burchak-Abramovich, N. I., and A. K. Vekua. 1971. The fossil ostrich from the Akchagil layers of Georgia. Acta Zoologica Cracoviensia 16:1–26.
- Burchak-Abramovich, N. I., and A. K. Vekua. 1990. The fossil ostrich *Struthio dmanisensis* sp. n. from the Lower Pleistocene of eastern Georgia. Acta Zoologica Cracoviensia 33:121–132.
- Campbell, K. E., and L. Marcus. 1992. The relationships of hindlimb bone dimensions to body weight in birds. Natural History Museum of Los Angeles County, Sciences Series 36:395–412.
- Ferring, R., O. Oms, J. Agustí, F. Berna, M. Nioradze, T. Shelia, M. Tappen, A. Vekua, D. Zhvania, and D. Lordkipanidze. 2011. Earliest human occupations at Dmanisi (Georgian Caucasus) dated to 1.85–1.78 Ma. Proceedings of the National Academy of Sciences of the United States of America 108:10432–10436.
- Field, D. J., C. Lynner, C. Brown, and S. A. F. Darroch. 2013. Skeletal correlates for body mass estimation in modern and fossil flying birds. PLoS ONE 8:e82000.
- Folch, A. 1992. Order Struthioniformes; pp. 76–110 in J. del Hoyo, A. Elliott, and J. Sargatal (eds.), Handbook of the Birds of the World, Volume 1. Lynx Edicions, Barcelona, Spain.
- Forsyth Major, C. J. 1888. Sur un gisement d'ossements fossiles dans l'île de Samos, contemporains de l'âge Pikermi. Comptes Rendus de l'Académie des Sciences 107:1178–1181.
- Handley, W. D., A. Chinsamy, A. M. Yates, and T. H. Worthy. 2016. Sexual dimorphism in the late Miocene mihirung *Dromornis stirtoni* (Aves: Dromornithidae) from the Alcoota Local Fauna of central Australia. Journal of Vertebrate Paleontology. doi: 10.1080/02724634.2016.1180298.
- Hansford, J. P., and S. T. Turvey. 2018. Unexpected diversity within the extinct elephant birds (Aves: Aepyornithidae) and a new identity for the world's largest bird. Royal Society Open Science 5:181295.
- Janoo, A., and S. Sen. 1998. Pliocene vertebrate locality of Çalta, Ankara, Turkey. 2. Aves: Struthionidae. Geodiversitas 20:339–351.
- Jánossy, D. 1972. Die mittelpleistozäne Vogelfauna der Stránská skála. Anthropos (Brno) 20:35–64.
- Kretzoi, M. 1954. Ostrich and camel remains from the Central Danube basin. Acta Geologica 2:231–242.
- Kurochkin, E. N. 1981. New representatives and evolution of two archaic gruiform families in Eurasia. Transaction of the Joint Soviet-Mongolian Paleontological Expedition 15:59–86.
- Kurochkin, E. N., and A. N. Lungu. 1970. [A new ostrich from the middle Sarmatian of Moldavia.] Paleontologicheskij Zhurnal 1970:118–126. [Russian]
- Lopatin, A. V., I. A. Vislobokova, A. V. Lavrov, D. B. Startsev, D. O. Gimranov, N. V. Zelenkov, E. N. Maschenko, M. V. Sotnikova, K. K. Tarasenko, and V. V. Titov. In press. Taurida Cave, a new locality of the early Pleistocene vertebrates in Crimea. Doklady Biological Sciences 485:40–43.
- Lowe, P. R. 1933. On some struthious remains: 1. Description of some pelvic remains of a large fossil ostrich, *Struthio oldawayi*, sp. n., from the lower Pleistocene of Oldaway (Tanganyika Territory); 2. Egg-shell fragments referable to *Psammornis* and other Struthionides collected by Mr. St. John Philby in Southern Arabia. Ibis 75:652–658.
- Lu, H.-Y., S.-W. Yi, Z.-W. Xu, Y.-L. Zhou, L. Zeng, F.-Y. Zhu, H. Feng, L.-N. Dong, H.-X. Zhuo, K.-F. Yu, J. Mason, X.-Y. Wang, Y.-Y. Chen, Q. Lu, B. Wu, Z.-B. Dong, J.-J. Qu, X.-M. Wang, and Z.-T. Guo. 2013. Chinese deserts and sand fields in Last Glacial Maximum and Holocene Optimum. Chinese Science Bulletin 58:2775–2783.
- Martin, R. 1903. Note on some remains of *Struthio karatheodoris* Maj. of the island of Samos. Proceedings of the Zoological Society of London 1903:203–210.

- Mayhew, D. F. 2012. Revision of the fossil vole assemblage (Mammalia, Rodentia, Arvicolidae) from Pleistocene deposits at Kisláng, Hungary. *Palaeontology* 55:11–29.
- Mayr, G. 2017. Avian Evolution. The Fossil Record of Birds and Its Paleobiological Significance. John Wiley & Sons, Chichester, U.K., 293 pp.
- Mikhailov, K. E. 1988. Comparison of eggshell of East European and Asian Pliocene ostriches. *Transaction of the Joint Soviet-Mongolian Paleontological Expedition* 34:65–72.
- Mikhailov, K. E., and E. N. Kurochkin. 1988. The eggshells of Struthioniformes from the Palearctic and its position in the system of views on Ratitae evolution. *Transaction of the Joint Soviet-Mongolian Paleontological Expedition* 34:43–64.
- Milne-Edwards, A. 1869–1871. *Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France*, Volume 2. Librairie de G. Masson, Paris, 632 pp.
- Müller, D. W., D. Codron, C. Meloro, A. Munn, A. Schwarm, J. Hummel, and M. Clauss. 2013. Assessing the Jarman-Bell Principle: scaling of intake, digestibility, retention time and gut fill with body mass in mammalian herbivores. *Comparative Biochemistry and Physiology A* 164:129–140.
- Murray, P. F., and P. Vickers-Rich. 2004. *Magnificent Mihirungs: The Colossal Flightless Birds of the Australian Dreamtime*. Indiana University Press, Bloomington, Indianapolis, 410 pp.
- Naidina, O. D., and K. Richards. 2016. Pollen evidence for Late Pliocene–Early Pleistocene vegetation and climate change in the North Caucasus, North-Western Caspian Region. *Quaternary International* 409:50–60.
- Palmqvist, P., D. R. Gröcke, A. Arribas, and R. A. Fariña. 2003. Paleoeological reconstruction of a lower Pleistocene large mammal community using biogeochemical ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$, $\delta^{18}\text{O}$, Sr:Zn) and ecomorphological approaches. *Paleobiology* 29:205–229.
- Popov, S. V., F. Rögl, F. F. Steininger, I. G. Shcherba, and M. Kovac (eds.). 2004. Lithological-Paleogeographical maps of Paratethys. 10 maps from Late Eocene to Pliocene. *Courier Forschungsinstitut Senckenberg* 250:1–46.
- Rook, L., and B. Martínez-Navarro. 2010. Villafranchian: the long story of a Plio-Pleistocene European large mammal biochronologic unit. *Quaternary International* 219:134–144.
- Sauer, F. G. F. 1979. A Miocene ostrich from Anatolia. *Ibis* 121:494–501.
- Spassov, N. 2016. Southeastern Europe as a route for the earliest dispersal of *Homo* toward Europe: ecological conditions and the timing of the first human occupation of Europe; pp. 281–290 in K. Harvati and M. Roksandic (eds.), *Paleoanthropology of the Balkans and Anatolia: Human Evolution and Its Context*. Springer Science + Business Media, Dordrecht, The Netherlands.
- Vekua, A. 2013. Giant ostrich in Dmanisi fauna. *Bulletin of the Georgian National Academy of Sciences* 7:143–148.
- Vislobokova, I., and A. Tesakov. 2013. Early and Middle Pleistocene of Northern Eurasia. *The Encyclopedia of Quaternary Science* 4:605–614.
- Worthy, T. H., and R. N. Holdaway. 2002. *The Lost World of the Moa. Prehistoric Life of New Zealand*. Indiana University Press, Bloomington, Indiana, 718 pp.
- Zelenkov, N. V., and E. N. Kurochkin. 2015. Class Aves; pp. 86–290 in E. N. Kurochkin, A. V. Lopatin, and N. V. Zelenkov (eds.), *Fossil Vertebrates of Russia and Adjacent Countries: Fossil Reptiles and Birds*, Part 3. GEOS, Moscow, Russia.
- Zelenkov, N. V., Z. Boev, and G. Lazaridis. 2016. A large ergilornithine (Aves, Gruiformes) from the Late Miocene of the Balkan Peninsula. *Paläontologische Zeitschrift* 90:145–151.

Submitted January 23, 2019; revisions received March 15, 2019; accepted April 2, 2019.

Handling editor: Vanesa De Pietri.