



Taylor & Francis  
Taylor & Francis Group

---

## Implications of Body-Mass Estimates for Dinosaurs

Author(s): Jan Peczakis

Source: *Journal of Vertebrate Paleontology*, Feb. 15, 1995, Vol. 14, No. 4 (Feb. 15, 1995), pp. 520-533

Published by: Taylor & Francis, Ltd. on behalf of The Society of Vertebrate Paleontology

Stable URL: <http://www.jstor.com/stable/4523591>

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

The Society of Vertebrate Paleontology and Taylor & Francis, Ltd. are collaborating with JSTOR to digitize, preserve and extend access to *Journal of Vertebrate Paleontology*

## IMPLICATIONS OF BODY-MASS ESTIMATES FOR DINOSAURS

JAN PECZKIS

Department of Earth Science, Northeastern Illinois University, Chicago, Illinois, 60625

**ABSTRACT**—Body-mass estimates have been made for 220 of the over 300 generally accepted dinosaur genera. The 1–10 ton body-mass category is the modal one for all dinosaur genera, dinosaurs on nearly every continent, dinosaurs during most stages of the Mesozoic, and dinosaurs in two of the three peak historical periods of dinosaur discoveries. Carnivorous dinosaurs were much smaller than herbivores during the Late Jurassic and again in the Late Cretaceous; at other times the two were roughly equivalent in mass. In terms of discovery of dinosaur genera over time, there has been a simultaneous increase in proportion of very small (under 10 kg) dinosaurs and a relative decline in giant dinosaurs (10–100 tons). This suggests that early researchers tended to collect giant dinosaurs.

### INTRODUCTION

Despite the existence of small dinosaurs (see Callison and Quimby, 1984), the dinosaurs' general large size has caused them to capture both the public and scientists' imagination. The purpose of this work is to analyze the entire dinosaur fossil record with respect to body masses.

There has been much research in recent years on the biology of body size (Calder, 1984; Schmidt-Nielsen, 1984). Size is important in the study of reproductive rates, speed of movement, distance travelled, etc., to say nothing of anatomy, physiology, and ecology (Wing and Sues, 1992). Body-mass estimates for many dinosaur genera are useful for paleoecological studies of dinosaur populations, such as those done by Farlow (1976) and Beland and Russell (1979).

Sixteen papers have been published, between 1905 and 1986, regarding the determination of dinosaur body mass (for a bibliographical listing, see Chure and McIntosh, 1989:216). Among early works, Colbert's (1962) is the most comprehensive. He provided body-mass estimates for 14 well-known dinosaur genera. Since his study, a number of authors (e.g., Bakker, 1980) have followed his lead, using volumetric displacement of scale models to calculate body masses. Hotton (1980) showed the distribution of over 50 unspecified dinosaur genera in terms of body mass. It was based on comparative allometric relationships (cubic scaling of body length, etc.) with the specimens upon which Colbert's (1962) scale models had been based.

The purpose of this work is to: 1) provide a compendium of masses (see Appendix) for as many valid dinosaurian genera as possible using only reliable methods (see below), 2) determine whether the body-mass clustering of genera in dinosaurian taxa resembles that of modern mammals, 3) compare and contrast the body-mass distribution of dinosaurian genera with that of mammals, 4) reexamine the biogeographic and temporal distribution of dinosaurian genera in the light of

body masses, 5) compare and contrast the body-mass distribution of carnivorous and herbivorous dinosaurs, 6) compare and contrast the dates of discovery of small and large dinosaurs.

Because the rate at which dinosaurian genera are described (6 per year on average: Dodson and Dawson, 1991) is so great, a work like this is out of date before it is published. It is current as of the compilation by Weishampel et al. (1992), with the addition of the following more recently described genera: *Andesaurus* (Calvo and Bonaparte, 1991), *Amargasaurus* (Salgado and Bonaparte, 1991), *Dyslocosaurus* (McIntosh et al., 1992), *Eoraptor* (Sereni et al., 1993), and *Utahraptor* (Kirkland et al., 1993).

### METHODS

Almost every skeletal feature in linear dimension correlates positively with body mass. However, an accurate estimate of body mass can only be made through the use of skeletal features that have been tested and found to have a close allometric relationship between some linear variable and body mass.

Anderson et al. (1985) have shown that there is an unexpectedly close correlation between femoral and/or humeral circumference and body mass for land vertebrates. This correlation is quite independent of the magnitude of body mass, lower-level taxonomy, or thermal physiology. A separate regression is necessary for bipeds and quadrupeds, and both were used by Anderson et al. (1985) to estimate the body masses of several dinosaurian genera. Inasmuch as limb-bone circumference is directly related to the bearing of body mass whereas pelvic height of scale models (e.g., Colbert, 1962) is only incidentally related to mass, mid-shaft limb-bone circumference is probably the most accurate method available for estimating the body masses of extinct tetrapods.

Whenever limb bones were known from skeletal material of a given genus of dinosaur, I measured, from

the cited literature, the diameter of the mid-shaft of the femur and/or humerus from various longitudinal perspectives. After averaging the values, I multiplied it by pi in order to convert it into circumference, and then used the regressions in Anderson et al. (1985) to estimate the body masses of the specimen of the genus. Because body mass estimates differ significantly when different methods are used on even the same specimen (Anderson et al., 1985), I did not quote specific values for body mass but placed genera into order-of-magnitude categories (e.g., 1–10 kg, 10–100 kg, 100 kg–1 ton), each of which I subdivided into thirds (e.g., 1–4 tons, 4–7 tons, and 7–10 tons for the 1–10 ton category).

If a body mass estimate had been made for a given genus by a previous researcher (e.g., Colbert, 1962; Bakker, 1980), I incorporated it into the database and marked it with an asterisk (\*). Whenever I used pelvic heights for my own body mass estimates, I used the reference models in Colbert (1962), but calibrated them according to the Anderson et al. (1985) data (which had suggested that several of the Colbert models had been too ponderous). In a few cases in the compendium (i.e., Giffin, 1989), I used cranial capacity (Jerison, 1973) to estimate body mass for dinosaurs that have no known postcranial remains. While this cannot, of course, be used for precise body-mass estimates, it can be used to place genera into different order-of-magnitude body-mass categories.

Disappointingly, many putatively valid genera are based on fragmentary remains. In such cases body-mass estimates were not attempted unless enough of the skeleton is known for the cited author to have made an estimate of the dinosaur's size. From this, a pelvic height (and hence a body mass) was estimated. Again, this cannot be very accurate, but it is suitable for placing genera into body-mass categories.

There is a large range of body masses for adult (i.e., sexually mature) members of a given dinosaurian genus, particularly if dinosaurs grew indeterminately as do modern reptiles. This is another reason why body-mass estimates were expressed as body mass categories and not specific values of body mass. The placing of a dinosaur genus in a given mass category means that most adult members of the genus fitted in that category and does not imply that all did, nor that adult members of the genus actually ranged in mass through all the values of the category.

Measurements from specimens identified as juveniles were not used in the database except in rare cases (and then labeled as such) when the only known usable specimen of a genus was a juvenile (i.e., *Mussaurus* and three other genera). These were included in the database for the sake of completeness but were not, of course, included in any of the histograms or calculations in this work.

The variation in adult body mass among congeneric dinosaurian individuals is significant. However, where there is a large range of sizes for a given genus in a collection (Dodson et al., 1980; Russell and McIntosh,




FIGURE 1. Weight-category distribution of all 216 adult dinosaur genera in the compendium (Appendix), and all 72 carnivore genera of same.

1980), the modal value tends to be near the maximum value. For this reason, whenever there was a range of values for body mass available for a genus, I used the largest value. However, in most cases, the values in the database are necessarily based on a single specimen, as nearly half of all dinosaur genera are known from only one specimen (Dodson, 1990).

Once the data (genera) were assembled, the database was expressed as a histogram (Fig. 1), with numbers included above the bars of the histogram. Next, the database was divided into geographic areas (Fig. 2), and then subject to an analysis according to the three historical periods of greatest dinosaur diversity (Fig. 3). This was followed by a stage-by-stage analysis (Fig. 4) with very brief stages combined together with longer stages (as in Dodson, 1990). In order to determine if there have been any obvious mass-related biases of dinosaur collection over time, the three peak historical periods of dinosaur description (Dodson and Dawson, 1991) were each expressed in terms of the body masses of the dinosaurs discovered (Fig. 5). For the biogeographic analysis (Fig. 2), Australia and Antarctica were omitted because of the paucity of their dinosaurian remains. For the carnivore/herbivore analysis, only the staurikosaurids, herrerasaurids, and theropods were accepted as carnivorous. For the stage-by-stage analysis (Fig. 4), the 100 g–1 kg category was omitted because only two adult dinosaur genera occur in that category.

## ANALYSIS

The database (Appendix) reveals that dinosaurs ranged in mass over six orders of magnitude, with the




FIGURE 2. The biogeographic distribution of dinosaurs sorted by body weight.

1–10 ton category being the modal weight range for dinosaurs. The distribution of the 216 adult dinosaur genera by body mass (Fig. 1) is similar to that of the 50 unspecified genera in Hotton (1980). This suggests that the relative number of genera in body mass categories is well established for dinosaurs and will not change substantially as a result of future discoveries.

Dinosaurs show a different relationship of body mass versus taxonomy than do mammals. Among mammalian orders, 51% of genera, on average, fall in the modal body-mass category of each order, with a range from 34% to 62% (Peczkis, 1988). To compare this with dinosaurs, I used dinosaur suborders (because there are only 2 commonly-recognized dinosaur orders). The modal body-mass category (in terms of per-

centages of total genera) varies greatly by suborder: 31% (Theropoda), 53% (Ceratopsia), 58% (Sauropodomorpha), 58% (Ornithopoda), 78% (Pachycephalosauria), 79% (Ankylosauria), and 100% (Stegosauria). Although the order is more inclusive than the suborder, it is the dinosaurian suborders that are more heterogeneous in terms of body mass than are mammalian orders. It thus appears that dinosaurs were less conservative in terms of evolution of a variety of body masses per equivalent taxon, than were mammals.

The modal body-mass category for modern mammals is 10–100 g (Hotton, 1980), whereas that of the rapsids was 1–10 kg (Peczkis, unpubl. data). By contrast, only a small percentage of dinosaur genera were under 10 kg (Fig. 1). At the same time, only a minority

of dinosaurs were in the giant (10–100 ton) range, meaning that most dinosaurs were no heavier than a large modern elephant.

Dinosaurs do not appear to show any great differences in terms of body mass relative to biogeography (Fig. 2). Except for Africa and South America (and then barely), the 1–10 ton category is the modal category for dinosaur genera on all continents. Africa has a larger fraction (nearly half) of its dinosaurs being giant (10–100 tons) than any other continent. However, the significance of this is uncertain, as the sample is small and Africa has not been well explored for dinosaurs. A clue to this distribution is the fact that 12 of the 27 African dinosaur genera (Fig. 2e) are Jurassic, a time of high sauropod diversity (see below).

The dinosaurs were very unevenly distributed during Mesozoic time, with most known genera appearing during three small intervals of the Mesozoic (Dodson, 1990). The first peak of diversity was the Norian-Sinemurian interval (Fig. 3a), followed by a larger peak during the Kimmeridgian-Tithonian (Fig. 3b), and finally by the largest peak in the Campanian-Maastrichtian (Fig. 3c). The first peak (Fig. 3a) was relatively homogenous in terms of numbers of genera in the various body mass categories. The second peak (Fig. 3b) was characterized by the dominance of large sauropods (10–100 tons), whereas the final peak (Fig. 3c) was marked by the overwhelming dominance of genera of the 1–10 ton category, especially herbivores. The significance of the disparity between large herbivores and carnivores is discussed below.

Since most dinosaurian genera existed for only one Mesozoic stage (Dodson, 1990), the foregoing analysis (Fig. 3a–c) involves the amalgamation of genera not all of which were contemporaries of each other. For this reason, the entire Mesozoic was subject to a stage-by-stage analysis (Fig. 4) in order to show the number of herbivorous and carnivorous genera, relative to body mass, in existence during each stage.

As a result, several trends can be seen. During all but a few of the Mesozoic stages, dinosaurs in the 1–10 ton category were the most numerous (Fig. 4). The larger sauropods (10–100 tons) were rare in the earliest stages of dinosaur evolution, peaked in the Late Jurassic, and then had a smaller peak in the Late Cretaceous (when dinosaurs peaked as a whole). Very small dinosaurs (1–10 kg) appeared infrequently and without any obvious trend throughout the Mesozoic. Dinosaurs in the 100 kg–1 ton and 1–10 ton categories appear closely to reflect overall dinosaur diversity trends throughout the Mesozoic. But since these two categories comprise the majority of dinosaur genera, they determine the diversity trend for all dinosaurs.

Carnivores comprise a greater fraction of dinosaurs in the 10–100 kg and 100 kg–1 ton categories than in any other. In fact, three-fourths of all carnivore genera were under 1 ton. This demonstrates that carnivores as a whole were significantly smaller than herbivores. However, there appear to be different evolutionary ten-




FIGURE 3. The body-weight distribution of dinosaur genera during the three greatest periods of dinosaur diversity in the Mesozoic.







FIGURE 4. Body-weight categories of dinosaur herbivores (white bars) and carnivores (black bars) during stages of the Mesozoic.

dencies, with regards to this, in the Mesozoic (Fig. 4). In the Late Triassic and Early Jurassic, there appears to be a rough equivalence in size between herbivores and carnivores. In the Late Jurassic, carnivores are markedly smaller than herbivores, especially because of the large numbers of giant sauropods. In the Early Cretaceous, carnivores and herbivores are once again roughly equivalent in mass. Finally, in the Late Cretaceous, carnivores are again significantly smaller than herbivores, though the disparity was not as extreme as in the Late Jurassic.


Carnivores are absent in the 10–100 ton category (Fig. 1). This suggests that the giant size of sauropods was an evolutionary strategy to avoid predation. If so, it is analogous to modern mammals. The largest land mammals alive today (rhinoceri, hippopotami, and elephants) are herbivores with no comparatively-sized carnivores in existence. These large mammals, at least when adult, are almost never attacked by predators

(Schaller, 1972). The largest known therapsids were also entirely herbivorous for possibly the same evolutionary reasons (Peczis, unpubl. data).


Dodson and Dawson (1991) have pointed out that there are three historical periods when dinosaur genera were discovered at greater rates than at any other time. They are: 1870–1900, 1900–1940, and 1970–present. The analysis of the body masses of dinosaurs discovered during these three historical periods (Fig. 5) show several trends. The 1–10 ton category is overwhelmingly dominant in the two most recent periods of dinosaur discovery, thus following the previously-discussed tendencies for this category to dominate. Very small dinosaurs (under 10 kg) increase from none in the first period, to 6% in the middle period, and amount to 17% of the dinosaur genera named during the most recent (i.e., current) period. Giant sauropods (10–100 tons) show nearly the opposite trend. They account for 36% of the genera in the earliest period of discovery,



1970-1993



1900-1940



1870-1900

FIGURE 5. Body-weight categories of dinosaurs discovered during the three historical periods of maximal dinosaur-searching activity.

decline to 12% during the second period, and are 14% of the current period. These trends may indicate that early dinosaur researchers were more likely to collect giant dinosaurs, perhaps because museums sought large dinosaur specimens for display.

It can be concluded that dinosaurs repeatedly evolved to a large range of sizes, and exploited a variety of niches. The human factor in dinosaur exploration (Dodson and Dawson, 1991) has also found an expression in the sizes of dinosaurs that have been found over time.

#### ACKNOWLEDGMENTS

I thank Drs. Peter Dodson and Paul Sereno for reviewing earlier versions of this manuscript. I am also grateful to Drs. Mike Brett-Surman and John McIntosh for supplying me with copies of papers from obscure publications.

#### LITERATURE CITED

- Anderson, J. F., A. Hall-Martin, and D. A. Russell. 1985. Long-bone circumference and weight in mammals, birds, and dinosaurs. *Journal of Zoology (London)* 207:53–61.
- Bakker, R. T. 1980. Dinosaur heresy—dinosaur renaissance; pp. 351–462 in R. D. K. Thomas and E. C. Olson (eds.), *A Cold Look at the Warm-blooded Dinosaurs*. AAAS Selected Symposia Series No. 28.
- . 1988. Review of the Late Cretaceous nodosaurid *Dinosauria*. *Denversaurus schlessmani*, a new armored dinosaur from the latest Cretaceous of South Dakota, the last survivor of the nodosaurians, with comments on stegosaur–nodosaur relationships. *Hunteria* 1:1–23.
- , P. M. Galton, J. Siegwarth, and J. Filla. 1990. A new latest Jurassic vertebrate fauna, from the highest levels of the Morrison Formation at Como Bluff, Wyoming. Part IV. The dinosaurs: a new *Othnielia*-like hypsilophodontoid. *Hunteria* 2:8–13.
- , M. Williams, and P. Currie. 1988. *Nanotyrannus*, a new genus of pygmy tyrannosaur, from the latest Cretaceous of Montana. *Hunteria* 1:1–30.
- Barsbold, R., T. Maryanska, and H. Olsomolska. 1992. Oviraptorosauria; pp. 249–258 in D. B. Weishampel, P. Dodson, and H. Osmolska (eds.), *The Dinosauria*. University of California Press, Berkeley.
- Bartholomai, A., and R. E. Molnar. 1981. *Muttaborrasaurus*, a new iguanodontid (Ornithischia: Ornithopoda) dinosaur from the Lower Cretaceous of Queensland. *Memoir of the Queensland Museum* 20:319–349.
- Beland, P., and D. A. Russell. 1979. Ectothermy in dinosaurs: paleoecological evidence from Dinosaur Provincial Park, Alberta. *Canadian Journal of Earth Sciences* 16:25–255.
- Bonaparte, J. F. 1976. *Pisanosaurus mertii* Casamiquela and the origin of the Ornithischia. *Journal of Paleontology* 50:808–820.
- . 1979. Dinosaurs: a Jurassic assemblage from Patagonia. *Science* 205:1377–1379.
- . 1984. Locomotion in rauisuchid thecodonts. *Journal of Vertebrate Paleontology* 3:210–218.
- . 1991. Los vertebrados fosiles de la Formacion Rio Colorado. de la ciudad de Neuquen v Cercanias. Cre-

- tatico superior, Argentina. *Revista del Museo Argentino de Ciencias Naturales "Bernardino Rivadavia" Instituto Nacional de Investigación de la Ciencias Naturales—Paleontología* 4:15–123.
- , F. E. Novas, and R. A. Coria. 1990. *Carnotaurus sastrei* Bonaparte, the horned, lightly built carnosaur from the Middle Cretaceous of Patagonia. *Contributions in Science (Natural History Museum of Los Angeles County)* 416:1–42.
- and J. E. Powell. 1980. A continental assemblage of tetrapods from the Upper Cretaceous bed of El Brete, northwest Argentina (Sauropoda-Coelurosauria-Carnosauria-Aves). *Societe Geologique de France Memoire* 139:19–28.
- and M. Vince. 1979. El hallazgo del primer nido de Dinosaurios Triasicos (Saurischia, Prosauropoda), Triasico superior de Patagonia, Argentina. *Ameghiniana* 16:173–182.
- Brett-Surman, M. K. 1980. The Iren Dabasu fauna from Mongolia; p. 19 in G. Olshevsky (ed.), *Mesozoic Vertebrate Life*. Stevens Publishing Co., Illinois.
- Brinkman, D. B., and H.-D. Sues. 1987. A staurikosaurid dinosaur from the Upper Triassic Ischigualasto Formation of Argentina and the relationships of the Staurikosauridae. *Palaeontology* 30:493–503.
- Brown, B., and E. M. Schlaikjer. 1942. The skeleton of *Leptoceratops* with the description of a new species. *American Museum Novitates* 1169:1–15.
- Calder, W. A. 1984. *Size, Function, and Life History*. Harvard University Press, Cambridge, London, 431 pp.
- Callison, G. 1987. Small problems: biological implications of tiny dinosaurs; pp. 70–80 in S. J. Czerkas and E. C. Olson (eds.) *Dinosaurs Past and Present*, Vol. I. University of Washington Press, Seattle.
- and H. M. Quimby. 1984. Tiny dinosaurs: are they fully grown? *Journal of Vertebrate Paleontology* 3:200–209.
- Calvo, J. O., and J. F. Bonaparte. 1991. *Andesaurus delgadoi* gen. et. sp. nov. (Saurischia-Sauropoda) dinosaurio Titanosauridae de la Formacion Rio Limay (Albiano-Cenomaniano), Neuquen, Argentina. *Ameghiniana* 28:303–310.
- Carpenter, K. 1990. Ankylosaur systematics: example using *Panoplosaurus* and *Edmontonia* (Ankylosauria: Nodosauridae); pp. 281–298 in K. Carpenter and P. J. Currie (eds.), *Dinosaur Systematics*. Cambridge University Press, Cambridge and New York.
- Chapman, R. E., and M. K. Brett-Surman. 1990. Morphometric observations on hadrosaurid ornithomorphs; pp. 163–178 in K. Carpenter and P. J. Currie (eds.), *Dinosaur Systematics*. Cambridge University Press, New York.
- Charig, A. J., and A. C. Milner. 1986. *Baryonyx*, a remarkable new theropod dinosaur. *Nature* 324:359–361.
- Chatterjee, S. 1978. *Indosuchus* and *Indosaurus*, Cretaceous carnosaur from India. *Journal of Paleontology* 52:570–580.
- 1987. A new theropod dinosaur from India with remarks on the Gondwana-Laurasia connection in the Late Triassic; pp. 183–189 in G. D. McKenzie (ed.), *Gondwana 6: Stratigraphy, Sedimentology, and Paleontology*. Geophysical Monograph 41.
- Chure, D. J., and J. S. McIntosh. 1989. A bibliography of the Dinosauria. *Museum of Western Colorado Paleontological Series* 1, 226 pp.
- Colbert, E. H. 1962. The weights of dinosaurs. *American Museum Novitates* 2076:1–16.
- Coombs, W. P., Jr. 1978a. The families of the ornithischian dinosaur order Ankylosauria. *Palaeontology* 21:143–170.
- 1978b. Theoretical aspects of cursorial adaptations in dinosaurs. *Quarterly Review of Biology* 53:393–418.
- Cooper, M. R. 1984. A reassessment of *Vulcanodon karibaensis* Raath (Dinosauria: Saurischia) and the origin of the Sauropoda. *Palaeontologia Africana* 25:203–231.
- 1989. The first record of the prosauropod dinosaur *Euskelosaurus* from Zimbabwe. *Arnoldia* 9:1–17.
- Cope, E. D. 1977. On *Amphicoelias* a genus of saurian from the Dakota epoch of Colorado. *Proceedings of the American Philosophical Society* 17:242–246.
- Corro, G. del. 1975. Un nuevo sauropodo del Cretacico superior. *Chubutisaurus insignis* gen. et sp. nov. (Saurischia-Chubutisauridae nov.) del Cretacico superior (Chubutiano), Chubut, Argentina. *Actas del Primer Congreso Argentino de Paleontologia y Biostratigrafia* 2:229–240.
- Currie, P. J., and D. A. Russell. 1988. Osteology and relationships of *Chirostenotes pergracilis* (Saurischia, Theropoda) from the Judith River (Oldman) Formation of Alberta, Canada. *Canadian Journal of Earth Sciences* 25:972–986.
- Dodson, P. 1983. A faunal review of the Judith River (Oldman) Formation, Dinosaur Provincial Park, Alberta. *The Mosasaur* 1:89–121.
- 1986. *Avaceratops lammersi*: a new ceratopsid from the Judith River Formation of Montana. *Proceedings of the Academy of Natural Sciences of Philadelphia* 138:305–317.
- 1990. Counting dinosaurs: how many kinds were there? *Proceedings of the Natural Academy of Sciences* 87:7608–7612.
- Dodson, P., A. K. Behrensmeyer, R. T. Bakker, and J. S. McIntosh. 1980. Taphonomy and palaeoecology of the dinosaur beds of the Jurassic Morrison Formation. *Paleobiology* 6:208–232.
- and S. D. Dawson. 1991. Making the fossil record of dinosaurs. *Modern Geology* 16:3–15.
- Dong, Z. 1977. On the dinosaurian remains from Turpan, Xinjiang. *Vertebrata Palasiatica* 15:59–66.
- 1984. A new theropod dinosaur from the Middle Jurassic of Sichuan Basin. *Vertebrata Palasiatica* 22:213–218.
- 1989. On a small ornithomorph (*Gongbasaurus wucuiwenensis* sp. nov.) from Kelamaili, Junggar Basin, Xinjiang, China. *Vertebrata Palasiatica* 27:140–147.
- 1990a. On remains of the sauropods from Kelamaili region, Junggar Basin, Xinjiang, China. *Vertebrata Palasiatica* 28:43–58.
- 1990b. Stegosaurids of Asia; pp. 255–268 in K. Carpenter and P. J. Currie (eds.), *Dinosaur Systematics*. Cambridge University Press, New York.
- , G. Peng, and D. Huang. 1989. The discovery of the bony tail club of sauropods. *Vertebrata Palasiatica* 27:219–224.
- and Z. Tang. 1984. Note on a New Mid-Jurassic Sauropod (*Datousaurus bashenensis* Gen. et. Sp. Nov.) from Sichuan Basin, China. *Vertebrata Palasiatica* 22:69–75.
- and ———. 1985. A new mid-Jurassic theropod (*Gasosaurus constructus*) Gen. et. Sp. Nov. from Das-



- hangpu, Zigong, Sichuan Province, China. *Vertebrata Palasiatica* 23:77–83.
- , S. Zhou, and Y. Zhang. 1983. The dinosaurian remains from Sichuan Basin, China. *Palaeontologica Sinica* 162:1–145.
- Dunham, A. E., K. L. Overall, W. P. Porter, and C. A. Forster. 1989. Implications of ecological energetics and biophysical and developmental constraints for life-history variation in dinosaurs; pp. 1–21 in J. O. Farlow (ed.), *Paleobiology of the Dinosaurs*. Geological Society of America Special Paper 238.
- Eaton, T. H. Jr. 1960. A new armored dinosaur from the Cretaceous of Kansas. *University of Kansas Paleontological Contributions* 25:1–24.
- Farlow, J. O. 1976. A consideration of the trophic dynamics of a Late Cretaceous large-dinosaur community (Oldham Formation). *Ecology* 57:841–857.
- Forster, C. A. 1990. The postcranial skeleton of the ornithomimid dinosaur *Tenontosaurus tilleti*. *Journal of Vertebrate Paleontology* 10:273–294.
- Fraas, E. 1908. Ostafrikanische dinosaurier. *Palaeontographica* 55:105–144.
- Galton, P. M. 1971. The prosauropod dinosaur *Ammosaurus*, the crocodile *Protosuchus*, and their bearing on the age of the Navajo Sandstone of northeastern Arizona. *Journal of Paleontology* 45:781–795.
- . 1977. On *Staurikosaurus pricei*, an early saurischian dinosaur from the Triassic of Brazil, with notes on the Herrerasauridae and Poposauridae. *Paläontologische Zeitschrift* 51:234–245.
- . 1978. Remains of ornithomimid dinosaurs from the Lower Cretaceous of North America. *Brigham Young University Studies in Geology* 25:1–10.
- . 1981. *Dryosaurus*, a hypsilophodontid dinosaur from the Upper Jurassic of North America and Africa: postcranial skeleton. *Paläontologische Zeitschrift* 55:271–312.
- . 1982. *Elaphrosaurus*, an ornithomimid dinosaur from the Upper Jurassic of North America and Africa. *Paläontologische Zeitschrift* 56:265–275.
- . 1983. Armored dinosaurs (Ornithischia: Ankylosauria) from the Middle and Upper Jurassic of Europe. *Palaeontographica* A182:1–25.
- . 1985a. The poposaurid thecodontian *Teratosaurus suevicus* v. Meyer, plus referred specimens mostly based on prosauropod dinosaurs, from the Middle Stubensandstein (Upper Triassic) of Nordwürttemberg. *Stuttgarter Beiträge zur Naturkunde* B116:1–29.
- . 1985b. British plated dinosaurs (Ornithischia, Stegosauridae). *Journal of Vertebrate Paleontology* 5:211–254.
- . 1985c. Notes on the Melanosauridae, a family of large prosauropod dinosaurs (Saurischia: Sauropodomorpha). *Géobios* 18:671–676.
- . 1992. Stegosauria; pp. 435–455 in D. B. Weishampel, P. Dodson, and H. Osmolska (eds.), *The Dinosauria*. University of California Press, Berkeley.
- and M. A. Cluver. 1976. *Anchisaurus capensis* (Broom) and a revision of the Anchisauridae (Reptilia, Saurischia). *Annals of the South African Museum* 69:121–159.
- and J. A. Jensen. 1975. *Hypsilophodon* and *Iguanodon* from the Lower Cretaceous of North America. *Nature* 257:668–669.
- and ———. 1978. Remains of ornithomimid dinosaurs from the Lower Cretaceous of North America. *Brigham Young University Studies in Geology* 25:1–10.
- and ———. 1979. A new large theropod from the Upper Jurassic Colorado. *Brigham Young University Studies in Geology* 26:1–11.
- and P. Taquet. 1982. *Valdosaurus*, a hypsilophodontid dinosaur from the Lower Cretaceous of Europe and Africa. *Géobios* 15:147–159.
- Giffin, E. B. 1989. Pachycephalosaur paleoneurology (Archosauria: Ornithischia). *Journal of Vertebrate Paleontology* 9:67–77.
- Gillette, D. D. 1991. *Seismosaurus halli*, gen et sp. nov., a new sauropod dinosaur from the Morrison Formation (Upper Jurassic/Lower Cretaceous) of New Mexico, USA. *Journal of Vertebrate Paleontology* 11:417–433.
- Gilmore, C. W. 1924. On *Troodon validus*, an orthopodous dinosaur from the Belly River Cretaceous of Alberta, Canada. *Bulletin of the Department of Geology, University of Alberta* 1:1–43.
- . 1933. On the dinosaurian fauna of the Iren Dabasu Formation. *Bulletin of the American Museum of Natural History* 67:23–78.
- Hatcher, J. B. 1903. Osteology of *Haplocanthosaurus*, with description of a new species, and remarks on the probable habits of the Sauropoda and the age and origin of the *Atlantosaurus* beds. *Memoir of the Carnegie Museum* 2:1–72.
- Haubold, H. 1991. Ein neuer Dinosaurier (Ornithischia, Thyreophora) aus dem unteren Jura des nördlichen Mitteleuropa. *Révue Paléobiologie* 9:149–177.
- He, X. 1979. A newly discovered ornithomimid dinosaur—*Yandusaurus* from Zigong, Sichuan; pp. 116–123 in *Contribution to International Exchange in Geology. Part 2. Stratigraphy and Paleontology*. Geological Publishing House, Beijing.
- Hotton, N., III. 1980. An alternative to dinosaur endothermy; pp. 311–350 in R. D. K. Thomas and E. C. Olson (eds.), *A Cold Look at the Warm-blooded Dinosaurs*. AAAS Selected Symposia Series No. 28.
- Hu, S. 1964. Carnosaurian remains from Alashan, Inner Mongolia. *Vertebrata Palasiatica* 8:42–63.
- Huene, F. von. 1923. Carnivorous Saurischia in Europe since the Triassic. *Bulletin of the Geological Society of America* 34:449–458.
- . 1926a. On several known and unknown reptiles of the order Saurischia from England and France. *Annals and Magazine of Natural History (Series 9)* 17:473–489.
- . 1926b. The carnivorous Saurischia in the Jura and Cretaceous Formations, principally in Europe. *Revista de Museum de La Plata* 29:35–167.
- . 1932. Die fossile Reptil-Ordnung Saurischia, ihre Entwicklung und Geschichte. *Monographien zur Geologie und Palaeontologie, Serie 1*, 4:1–361.
- . 1934. Ein neuer Coelurosaurier in der thüringischen Trias. *Palaeontologische Zeitschrift* 16:145–170.
- and C. A. Matley. 1933. The Cretaceous Saurischia and Ornithischia of the central provinces of India. *Memoir of the Geological Survey of India* 21:1–74.
- Hulke, J. W. 1874. Note on a very large saurian limb-bone adapted for progression on land, from the Kimmeridge Clay of Weymouth, Dorset. *Quarterly Journal of the Geological Society of London* 30:16–17.
- Jain, S. L., T. S. Kutty, T. Roy-Chowdhury, and S. Chatterjee. 1979. Some characteristics of *Baraposaurus tagorei*, a sauropod dinosaur from the Lower Jurassic of

- Deccan, India. Proceedings of the Fourth International Gondwana, Symposium, Calcutta, Volume 1:204–216.
- Janensch, W. 1925. Die Coelurosaurier und Theropoden der Tendaguru-Schichten Deutsch-Ostafrikas. *Palaeontographica* (Supplement 7)1:1–99.
- . 1961. Die Gliedmassen und Gliedmaszengürtel der Sauropoden der Tendaguru-Schichten. *Palaeontographica* (Supplement 7)3:177–235.
- Jensen, J. A. 1987. New Brachiosaur material from the Late Jurassic of Utah and Colorado. *Great Basin Naturalist* 47:592–608.
- Jerison, H. J. 1973. *Evolution of the Brain and Intelligence*. Academic Press, New York and London, 482 pp.
- Kirkland, J. I., D. Burge, and R. Gaston. 1993. A large dromaeosaur (Theropoda) from the Lower Cretaceous of eastern Utah. *Hunteria* 2:1–16.
- Langston, W., Jr., 1960. The vertebrate fauna of the Selma Formation of Alabama. Part IV. The dinosaurs. *Fieldiana, Geological Memoir* 3:313–361.
- . 1975. The ceratopsian dinosaurs and associated lower vertebrates from the St. Mary River Formation (Maastrichtian) at Scabby Butte, Southern Alberta. *Canadian Journal of the Earth Sciences* 12:1576–1608.
- Lapparent, A. F. de. 1943. Les dinosauriens Jurassiques de Damparis (Jura). *Mémoire de la Société Géologique de France* 47:5–20.
- . 1947. Les dinosauriens du Cretace superieur du Midi de la France. *Mémoire de la Société Géologique de France* 56:1–54.
- . 1960. Les dinosauriens du “Continental intercalaire” du Sahara central. *Mémoire de la Société Géologique de France* 88A:1–57.
- and G. Zbyszewski. 1957. Les dinosauriens du Portugal. *Mémoire Services Géologiques du Portugal* 2:1–63.
- Lehman, T. M., and K. Carpenter. 1990. A partial skeleton of the tyrannosaurid dinosaur *Aublysodon* from the Upper Cretaceous of New Mexico. *Journal of Paleontology* 64:1026–1032.
- LeLouff, J., and E. Buffetaut. 1991. *Tarascosaurus salluvicus* nov. gen., nov. sp., dinosaure théropode du Cretace superieur de Sud de la France. *Géobios* 25:585–594.
- Lull, R. S. 1933. A revision of the Ceratopsia or horned dinosaurs. *Peabody Museum of Natural History Bulletin* 3:1–175.
- and N. E. Wright. 1942. Hadrosaurian dinosaurs of North America. *Geological Society of America Special Paper* 40:1–242.
- Lydekker, R. 1893. Contributions to the study of the fossil vertebrates of Argentina. I. The dinosaurs of Patagonia. *Annals of the Museum of La Plata, Paleontology Section* 2:1–14.
- McIntosh, J. S., W. P. Coombs, and D. A. Russell. 1992. A new Diplodocid sauropod (Dinosauria) from Wyoming, USA. *Journal of Vertebrate Paleontology* 12:158–167.
- Maleev, E. A. 1974. Gigantskie karnazavry semeistva Tyrannosauridae. *Sovmestnaia Sovetsko-Mongol'skaia Paleontologicheskaja Ekspeditsiia, Trudy* 1:132–191.
- Mantell, G. A. 1841. Memoir on a portion of the lower jaw of the *Iguanodon*, and on the remains of the *Hylaeosaurus* and other saurians, discovered in the strata of Tilgate Forest in Sussex. *Philosophical Transactions of the Royal Society, Part II*, 151 pp.
- . 1850. On the *Pelorosaurus*; an undescribed gigantic terrestrial reptile. *Philosophical Transactions of the Royal Society of London* 140:379–390.
- Marsh, O. C. 1877. Notice of a new gigantic dinosaur. *American Journal of Science* (series 3) 14:87–88.
- . 1889. Notice of gigantic horned Dinosauria from the Cretaceous. *American Journal of Science* (series 3) 38:173–175.
- . 1896. The Dinosaurs of North America. United States Geological Survey 16th Annual Report 1894 5:133–244.
- Martinez, R., O. Gimenez, J. Rodriguez, and G. Bochaty. 1986. *Xenotarsosaurus bonapartei* nov. gen et sp. (Carnosauria, Abelisauridae), un nuevo Theropoda de la Formacion Bajo Barreal, Chubut, Argentina. *Actas IV Congreso Argentino de Paleontologia y Biostratigrafia* 2:23–31.
- Maryanska, T. 1977. Ankylosauridae (Dinosauria) from Mongolia. *Palaeontologia Polonica* 37:85–151.
- and H. Osmolska. 1974. Pachycephalosauria, a new suborder of ornithischian dinosaurs. *Palaeontologia Polonica* 30:45–102.
- and H. Osmolska. 1975. Proceratopsidae (Dinosauria) of Asia. *Palaeontologia Polonica* 33:133–182.
- Molnar, R. E. 1974. A distinctive theropod dinosaur from the Upper Cretaceous of Baja California (Mexico). *Journal of Paleontology* 48:1009–1017.
- Morris, W. J. 1981. A new species of hadrosaurian dinosaur from the Upper Cretaceous of Baja California—? *Lambeosaurus laticaudus*. *Journal of Paleontology* 55:453–462.
- Osborn, H. F. 1924. Three new Theropoda, *Protoceratops* zone, central Mongolia. *American Museum Novitates* 144:1–12.
- Osmolska, H., and E. Roniewicz. 1970. Deinocheiridae, a new family of theropod dinosaurs. *Palaeontologia Polonica* 21:5–19.
- and R. Barsbold. 1972. A new dinosaur, *Gallimimus bullatus* n. gen., n. sp. (Ornithomimidae) from the Upper Cretaceous of Mongolia. *Palaeontologia Polonica* 27:103–143.
- Ostrom, J. H. 1969. Osteology of *Deinonychus antirrhopus*, an unusual theropod from the Lower Cretaceous of Montana. *Peabody Museum of Natural History Bulletin* 30:1–165.
- . 1970. Stratigraphy and paleontology of the Cloverly Formation (Lower Cretaceous) of the Bighorn Basin area, Wyoming and Montana. *Peabody Museum of Natural History Bulletin* 35:1–234.
- . 1976. On a new specimen of the Lower Cretaceous theropod dinosaur *Deinonychus antirrhopus*. *Breviora* 439:1–21.
- Peczkis, J. 1988. Predicting body-weight distribution of mammalian genera in families and orders. *Journal of Theoretical Biology* 132:509–510.
- Perle, A. 1979. Segnosauridae—Novoe semeistvo teropod iz pozdnovo mela Mongolii. *Sovmestnaia Sovetsko-Mongol'skaia Paleontologicheskaja Ekspeditsiia, Trudy* 8:45–55.
- , T. Maryanska, and H. Osmolska. 1982. *Goyocephale laticauda* gen. et sp. n., a new flat-headed pachycephalosaur (Ornithischia, Dinosauria) from the Upper Cretaceous of Mongolia. *Acta Palaeontologica Polonica* 27:115–127.
- Powell, J. E. 1979. Sobre una asociacion de dinosaurios y otras evidencias de vertebrados del Cretacico superior

- del la region de La Candleria, Province de Salta, Argentina. *Ameghiniana* 16:191–204.
- Raath, M. A. 1990. Morphological variation in small theropods and its meaning in systematics: evidence from *Syntarsus rhodesiensis*; pp. 91–105 in K. Carpenter and P. J. Currie (eds.), *Dinosaur Systematics*. Cambridge University Press, New York.
- and J. S. McIntosh. 1987. Sauropod dinosaurs from the central Zambezi Valley, Zimbabwe, and the age of the Kadzi Formation. *South African Journal of Geology* 90:107–119.
- Reig, O. A. 1963. La presencia de dinosaurios saurisquios en los “Estratos de Ischigualasto” (Mesotriasico superior) de la Provincias de San Juan y La Rioja (Republica Argentina). *Ameghiniana* 3:3–20.
- Rich, T. H. V., and P. V. Rich. 1989. Polar dinosaurs and biotas of the Early Cretaceous of southeastern Australia. *National Geographic Research* 5:15–53.
- Russell, D. A. 1970. Tyrannosaurs from the Late Cretaceous of western Canada. *National Museum of Natural Science, Publications in Palaeontology* 1:1–34.
- 1972. Ostrich dinosaurs from the Late Cretaceous of Western Canada. *Canadian Journal of the Earth Sciences* 9:375–402.
- and P. Béland. 1976. Running dinosaurs. *Nature* 264:486.
- and J. S. McIntosh. 1980. Paleoeecology of the dinosaurs of Tendaguru (Tanzania). *Mémoire de la Société Géologique de France, N. S.*, 139:169–175.
- Salgado, L., and J. F. Bonaparte. 1991. Un nuevo sauro-podo Dicraeosauridae, *Amargosaurus cazaui* gen. et. sp. nov., de La Formacion la Amarga, Neocomiano de la Provincia del Neuquen, Argentina. *Ameghiniana* 28:333–346.
- Santa Luca, A. P., A. W. Crompton, and A. W. Charig. 1976. A complete skeleton of the Late Triassic ornithischian *Heterodontosaurus tucki*. *Nature* 264:324–328.
- Schaller, G. B. 1972. *The Serengeti Lion*. University of Chicago Press, Chicago, 480 pp.
- Schmidt-Nielsen, K. 1984. *Scaling: Why is Animal Size So Important*. Cambridge University Press, Cambridge, 241 pp.
- Sereno, P. C., and C. Shichin. 1988. *Psittacosaurus xinjiangensis* (Ornithischia: Ceratopsia), a new psittacosaur from the Lower Cretaceous of Northwestern China. *Journal of Vertebrate Paleontology* 8:353–365.
- Sereno, P. C., C. A. Forster, R. R. Rogers, and A. M. Monetta. 1993. Primitive dinosaur skeleton from Argentina and the early evolution of Dinosauria. *Nature* 361:64–66.
- Stovall, J. W., and W. Langston, Jr. 1950. *Acrocanthosaurus atokensis*, a new genus and species of Lower Cretaceous Theropoda from Oklahoma. *American Midland Naturalist* 43:696–728.
- Stromer, E. 1932. *Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wüsten Ägyptens. II. Wirbeltierreste der Baharije-Stufe (unterstes Cenoman). 11. Sauropoda*. Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung 10:1–21.
- 1934. Wirbeltierreste der Baharije-Stufe (unterstes Cenoman). 13. Dinosauria. Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung 10:1–21.
- 1934. Wirbeltierreste der Baharije-Stufe (unterstes Cenoman). 13. Dinosauria. Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Abteilung 22:1–79.
- Taquet, P. 1976. *Géologie et paléontologie du gisement de Gadoufaoua (Aptien du Niger)*. Cahiers de Paléontologie, C.N.R.S. Paris, 191 pp.
- Walker, A. D. 1964. Triassic reptiles from the Elgin area: *Ornithosuchus* and the origin of carnosaurs. *Philosophical Transactions of the Royal Society of London B248*: 53–134.
- Weishampel, D. B., P. Dodson, and H. Osmolska (eds). 1992. *The Dinosauria*. University of California Press, Berkeley, 733 pp.
- Welles, S. P. 1984. *Dilophosaurus wetherelli* (Dinosauria, Theropoda). Osteology and comparisons. *Palaeontographica A* 185:85–180.
- Wild, R. 1991. *Janenschia* n. g. *robusta* (Fraas 1908) pro *Tornieria robusta* (E. Fraas 1908) (Reptilia, Saurischia, Sauropodomorpha). *Stuttgarter Beiträge für Naturkunde, Serie B*, 173:1–4.
- Wiman, C. 1929. Die Kreide-Dinosaurier aus Shantung. *Palaeontologia Sinica, C*, 6:1–67.
- Wing, S. L., and H.-D. Sues. 1992. Mesozoic and early Cenozoic terrestrial ecosystems; pp. 327–416 in A. K. Behrensmeyer, J. D. Damuth, W. A. DiMichelle, R. Potts, H.-D. Sues, and S. L. Wing (eds.), *Terrestrial Ecosystems through Time*. University of Chicago Press, Chicago.
- Yadagiri, P. 1988. A new sauropod *Kotasaurus yumanpalliensis* from Lower Jurassic Kota Formation of India. *Records of the Geological Survey of India* 11:102–127.
- Young, C. C. 1937. A new dinosaurian from Sinkiang. *Palaeontologia Sinica, C*, 2:1–25.
- 1942. *Yunnanosaurus huangi* (gen. et sp. nov.), a new prosauropod from the Red Beds at Lufeng, Yunnan. *Bulletin of the Geological Society of China* 22:63–104.
- 1947. On *Lufengosaurus magnus* (sp. nov.) and additional finds of *Lufengosaurus huenei* Young. *Palaeontologia Sinica* 12:1–53.
- 1954. On a new sauropod from Yiping, Szechuan, China. *Acta Palaeontologica Sinica* 2:355–369.
- 1958. The dinosaurian remains of Laiyang, Shantung. *Palaeontologia Sinica, C*, 16:1–138.
- 1959. On a new Stegosauria from Szechuan, China. *Vertebrata Palasiatica* 3:1–8.

Received 19 August 1992; accepted 9 August 1993.

APPENDIX

Body weight categories of all dinosaur genera known from sufficient skeletal material for a body mass estimate. Abbreviations are as follows: body mass categories calculated from: M = scale model, P = pelvic height determination, F = femur diameter, f = femur length, H = humerus diameter, E = endocast volume. The asterisk (\*) denotes body mass already estimated by cited author; all others are my estimates using the indicated information from the cited author(s). For the convenience of the reader, the genera are listed in the same order as they appear in Weishampel et al. (1992).

Genus	Higher taxon	Method	Weight	Reference
<i>Staurikosaurus</i>	Staurikosauridae	F	10–40 kg	Galton, 1977a
<i>Herrerasaurus</i>	Herrerasauridae	F	100–400 kg	Galton, 1977a
<i>Ischisaurus</i>	Herrerasauridae	F	40–70 kg	Reig, 1963
<i>Aliwalia</i>	(Dinosauria)	F	1–4 tons	Galton, 1985a
<i>Ceratosaurus</i>	(Ceratosauria)	F(*)	400–700 kg	Anderson et al., 1985
<i>Sarcosaurus</i>	(Ceratosauria)	F	40–70 kg	Huene, 1932
<i>Segisaurus</i>	(Ceratosauria)	P	4–7 kg	Callison, 1987
<i>Dilophosaurus</i>	(Ceratosauria)	F, P	100–400 kg	Welles, 1984
<i>Liliensternus</i>	(Ceratosauria)	F	100–400 kg	Huene, 1934
<i>Coelophysis</i>	(Ceratosauria)	F	10–40 kg	Raath, 1990
<i>Syntarsus</i>	(Ceratosauria)	F	10–40 kg	Raath, 1990
<i>Acrocantosaur</i>	Allosauridae	F	1–4 tons	Stovall and Langston, 1950
<i>Allosaurus</i>	Allosauridae	F(*)	1–4 tons	Anderson et al., 1985
<i>Chilantaisaurus</i>	Allosauridae	F	1–4 tons	Hu, 1964
<i>Piatnitzkysaurus</i>	Allosauridae	F	100–400 kg	Bonaparte, 1984
<i>Szechuanosaurus</i>	Allosauridae	F	7–10 tons	Dong et al., 1983
<i>Albertosaurus</i>	Tyrannosauridae	M(*)	1–4 tons	Bakker, 1980
<i>Alectrosaurus</i>	Tyrannosauridae	P	0.7–1 ton	Brett-Surman, 1980
<i>Daspletosaurus</i>	Tyrannosauridae	F	7–10 tons	Russell, 1970
<i>Nanotyrannus</i>	Tyrannosauridae	P	400–700 kg	Bakker et al., 1988
<i>Tarbosaurus</i>	Tyrannosauridae	F, P	1–4 tons	Maleev, 1974
<i>Tyrannosaurus</i>	Tyrannosauridae	F(*)	4–7 tons	Anderson et al., 1985
<i>Bahariasaurus</i>	(Carnosauria)	F	1–4 tons	Stromer, 1934
<i>Aublysodon</i>	(Carnosauria)	F	1–4 tons	Lehman and Carpenter, 1990
<i>Eustreptospondylus</i>	(Carnosauria)	H	100–400 kg	Galton, 1982
<i>Gasosaurus</i>	(Carnosauria)	F	100–400 kg	Dong and Tang, 1985
<i>Labocania</i>	(Carnosauria)	P	1–4 tons	Molnar, 1974
<i>Magnasaurus</i>	(Carnosauria)	F	1–4 tons	Huene, 1926b
<i>Megalosaurus</i>	(Carnosauria)	H	1–4 tons	Galton and Jensen, 1979
<i>Spinosaurus</i>	(Carnosauria)	F	1–4 tons	Stromer, 1934
<i>Stokesosaurus</i>	(Carnosauria)	F	70–100 kg	Dodson et al., 1980
<i>Yangchuanosaurus</i>	(Carnosauria)	F	1–4 tons	Dong et al., 1983
<i>Ornithomimus</i>	Ornithomimidae	F, P	100–400 kg	Russell, 1972
<i>Struthiomimus</i>	Ornithomimidae	F(*)	100–400 kg	Anderson et al., 1985
<i>Dromiceiomimus</i>	Ornithomimidae	M(*)	100–400 kg	Russell and Beland, 1976
<i>Archaeornithomimus</i>	Ornithomimidae	P	40–70 kg	Brett-Surman, 1980
<i>Gallimimus</i>	Ornithomimidae	F	10–40 kg	Osmolska et al., 1972
<i>Elaphrosaurus</i>	(Ornithomimosauria)	F	100–400 kg	Janensch, 1925
<i>Chirostenotes</i>	Elmisauridae	F	40–70 kg	Currie and Russell, 1988
<i>Oviraptor</i>	Oviraptoridae	P	10–40 kg	Dodson, 1983
<i>Ingenia</i>	Oviraptoridae	F	40–70 kg	Barsbold et al., 1992
<i>Caenagnathus</i>	Caenagnathidae	P	10–40 kg	Dodson, 1983
<i>Troodon</i>	Troodontidae	P	10–40 kg	Osborn, 1924
<i>Saurnithoides</i>	Troodontidae	P	40–70 kg	Dodson, 1983
<i>Deinonychus</i>	Dromaeosauridae	F, P	70–100 kg	Ostrom, 1976
<i>Dromaeosaurus</i>	Dromaeosauridae	P	40–70 kg	Dodson, 1983
<i>Saurornitholestes</i>	Dromaeosauridae	P	10–40 kg	Dodson, 1983
<i>Utahraptor</i>	Dromaeosauridae	P	0.7–1 ton	Kirkland et al., 1993
<i>Velociraptor</i>	Dromaeosauridae	F	40–70 kg	Ostrom, 1969
<i>Avimimus</i>	(Theropoda)	P	1–4 kg	Callison, 1987
<i>Coelurus</i>	(Theropoda)	f(*)	10–40 kg	Dodson et al., 1980
<i>Compsognathus</i>	(Theropoda)	P(*)	400–700 kg	Callison, 1987
<i>Deinocheirus</i>	(Theropoda)	H	7–10 tons	Osmolska and Roniewicz, 1970
<i>Ornitholestes</i>	(Theropoda)	F, P	10–40 kg	Huene, 1932
<i>Alvarezsaurus</i>	(Theropoda)	F	4–7 kg	Bonaparte, 1991
<i>Eoraptor</i>	(Theropoda)	P	7–10 kg	Sereno et al., 1993



## APPENDIX

*Continued.*

Genus	Higher taxon	Method	Weight	Reference
<i>Microvenator</i>	(Theropoda)	F	1–4 kg	Ostrom, 1970
<i>Podokesaurus</i>	(Theropoda)	P	10–40 kg	Huene, 1932
<i>Walkeria</i>	(Theropoda)	F	1–4 kg	Chatterjee, 1987
<i>Baryonyx</i>	(Theropoda)	H	1–4 tons	Charig and Milner, 1986
<i>Carnotaurus</i>	(Theropoda)	F, P	1–4 tons	Bonaparte et al., 1990
<i>Dryptosaurus</i>	(Theropoda)	F	1–4 tons	Marsh, 1877
<i>Erectopus</i>	(Theropoda)	P	1–4 tons	Huene, 1923
<i>Frenguellisaurus</i>	(Theropoda)	F	100–400 kg	Brinkman and Sues, 1987
<i>Indosaurus</i>	(Theropoda)	F	400–700 kg	Huene and Matley, 1933; Chatterjee, 1978
<i>Indosuchus</i>	(Theropoda)	F	0.7–1 ton	Huene and Matley, 1933; Chatterjee, 1978
<i>Marshosaurus</i>	(Theropoda)	f	100–400 kg	Dodson et al., 1980
<i>Metriacanthosaurus</i>	(Theropoda)	F	0.7–1 ton	Huene, 1926a, b; Walker, 1964
<i>Shanshanosaurus</i>	(Theropoda)	F	100–400 kg	Dong, 1977
<i>Tarascosaurus</i>	(Theropoda)	F	7–10 kg	LeLoeuff and Buffetaut, 1991
<i>Xenotarsosaurus</i>	(Theropoda)	F	0.7–1 ton	Martinez et al., 1986
<i>Xuanhanosaurus</i>	(Theropoda)	P	0.7–1 ton	Dong, 1984b
<i>Thecodontosaurus</i>	Thecodosauridae	F, P	70–100 kg	Huene, 1932
<i>Anchisaurus</i>	Anchisauridae	F, P	10–40 kg	Galton and Cluver, 1976
<i>Massospondylus</i>	Massospondylidae	F, P	100–400 kg	Galton and Cluver, 1976
<i>Yunnanosaurus</i>	Yunnanosauridae	F	0.7–1 ton	Young, 1942
<i>Ammosaurus</i>	Plateosauridae	F, P	10–40 kg	Galton, 1971
<i>Mussaurus</i> (juv.)	Plateosauridae	F, H	100–400 g	Bonaparte and Vince, 1979
<i>Plateosaurus</i>	Plateosauridae	F	1–4 tons	Huene, 1932
<i>Sellosaurus</i>	Plateosauridae	F, P	100–400 kg	Galton and Cluver, 1976
<i>Camelotia</i>	Melanorosauridae	F	1–4 tons	Galton, 1985c
<i>Euskelosaurus</i>	Melanorosauridae	F	1–4 tons	Cooper, 1989
<i>Lufengosaurus</i>	Melanorosauridae	F	1–4 tons	Young, 1947
<i>Melanorosaurus</i>	Melanorosauridae	F	1–4 tons	Galton, 1985c
<i>Riojasaurus</i>	Melanorosauridae	F	1–4 tons	Galton, 1985c
<i>Vulcanodon</i>	Vulcanodontidae	F	10–40 tons	Cooper, 1984
<i>Barapasaurus</i>	Vulcanodontidae	F	10–40 tons	Jain et al., 1979
<i>Kotasaurus</i>	Vulcanodontidae	F	10–40 tons	Yadagiri, 1988
<i>Cetiosaurus</i>	Cetiosauridae	F, P	10–40 tons	Huene, 1932
<i>Bellusaurus</i> (juv.)	Vulcanodontidae	F	4–7 tons	Dong, 1990a
<i>Haplocanthosaurus</i>	Cetiosauridae	F	10–40 tons	Hatcher, 1903
<i>Patagosaurus</i>	Cetiosauridae	P	10–40 tons	Bonaparte, 1979
<i>Shunosaurus</i>	Cetiosauridae	F, P	7–10 tons	Dong and Tang, 1984
<i>Omeisaurus</i>	Cetiosauridae	F, P	10–40 tons	Dong et al., 1989
<i>Brachiosaurus</i>	Brachiosauridae	F, H(*)	10–40 tons	Anderson et al., 1985
<i>Bothriospondylus</i>	Brachiosauridae	F	40–70 tons	Lapparent, 1943
<i>Pelorosaurus</i>	Brachiosauridae	H	7–10 tons	Mantell, 1850
<i>Pleurocoelus</i>	Brachiosauridae	F	7–10 tons	Lapparent and Zbyszewski, 1957
<i>Ultrasauros</i>	Brachiosauridae	P	40–70 tons	Jensen, 1987
<i>Chubutisaurus</i>	Brachiosauridae	F, H	10–40 tons	Corro, 1975
<i>Ischyrosaurus</i>	Brachiosauridae	H	10–40 tons	Hulke, 1874
<i>Camarasaurus</i>	Camarasauridae	M(*)	10–40 tons	Bakker, 1980
<i>Euhelopus</i>	Camarasauridae	F, P	7–10 tons	Wiman, 1929
<i>Tienshanosaurus</i>	Camarasauridae	F	7–10 tons	Young, 1937
<i>Opisthocoeleicauda</i>	Camarasauridae	F, H(*)	10–40 tons	Anderson et al., 1985
<i>Diplodocus</i>	Diplodocidae	F, H(*)	10–40 tons	Anderson et al., 1985
<i>Barosaurus</i>	Diplodocidae	F	10–40 tons	Janensch, 1961
<i>Apatosaurus</i>	Diplodocidae	F, H(*)	10–40 tons	Anderson et al., 1985
<i>Dyslocosaurus</i>	Diplodocidae	F	7–10 tons	McIntosh et al., 1992
<i>Amphicoelias</i>	Diplodocidae	F	40–70 tons	Cope, 1877
<i>Seismosaurus</i>	Diplodocidae	P	40–70 tons	Gillette et al., 1991
<i>Amargasaurus</i>	Diplodocidae	F	10–40 tons	Salgado and Bonaparte, 1991
<i>Dicraeosaurus</i>	Diplodocidae	F	10–40 tons	Janensch, 1961
<i>Rebbachisaurus</i>	Diplodocidae	F, H	10–40 tons	Lapparent, 1960



APPENDIX  
*Continued.*

Genus	Higher taxon	Method	Weight	Reference
<i>Mamenchisaurus</i>	Mamenchisauridae	F	10–40 tons	Young, 1954
<i>Titanosaurus</i>	Titanosauridae	F	40–70 tons	Anderson et al., 1985
<i>Magyarosaurus</i>	Titanosauridae	F	0.7–1 ton	Huene, 1932
<i>Laplatosaurus</i>	Titanosauridae	F, H	7–10 tons	Powell, 1979
<i>Andesaurus</i>	Titanosauridae	F	40–70 tons	Calvo and Bonaparte, 1991
<i>Saltasaurus</i>	Titanosauridae	F	7–10 tons	Bonaparte and Powell, 1980
<i>Aegyptosaurus</i>	Titanosauridae	F	10–40 tons	Stromer, 1932
<i>Alamosaurus</i>	Titanosauridae	M(*)	10–40 tons	Bakker, 1980
<i>Hypselosaurus</i>	Titanosauridae	F	7–10 tons	Lapparent, 1947
<i>Argyrosaurus</i>	Titanosauridae	F	10–40 tons	Lydekker, 1893
<i>Antarctosaurus</i>	Titanosauridae	F	40–70 tons	Anderson et al., 1985
<i>Tornieria</i>	Titanosauridae	H	10–40 tons	Raath and McInotsh, 1987
<i>Janenschia</i>	Titanosauridae	F	10–40 tons	Wild, 1991; Fraas, 1908
<i>Segnosaurus</i>	Segnosauridae	F(?)	4–7 tons	Perle, 1979
<i>Lesothosaurus</i>	(Ornithischia)	P	4–7 kg	Callison, 1987
<i>Pisanosaurus</i>	(Ornithischia)	F	7–10 kg	Bonaparte, 1976
<i>Scutellosaurus</i>	(Thyreophora)	P	7–10 kg	Callison, 1987
<i>Emausaurus</i>	(Thyreophora)	P	40–70 kg	Haubold, 1991
<i>Scelidosaurus</i>	Scelidosauridae	F, P	400–700 kg	Marsh, 1896
<i>Echinodon</i>	(Thyreophora)	P(*)	0.7–1 kg	Callison, 1987
<i>Huayangosaurus</i>	Huayangosauridae	P	1–4 tons	Galton, 1992
<i>Chialingosaurus</i>	Stegosauridae	F	1–4 tons	Young, 1959
<i>Chungkingosaurus</i>	Stegosauridae	F	1–4 tons	Dong et al., 1983
<i>Dacentrurus</i>	Stegosauridae	F, P	1–4 tons	Galton, 1985b
<i>Kentrosaurus</i>	Stegosauridae	F, H(*)	1–4 tons	Russell et al., 1980
<i>Lexovisaurus</i>	Stegosauridae	F, P	1–4 tons	Galton, 1985b
<i>Monkonosaurus</i>	Stegosauridae	P	1–4 tons	Dong, 1990b
<i>Stegosaurus</i>	Stegosauridae	M(*)	4–7 tons	Bakker, 1980
<i>Tuojiangosaurus</i>	Stegosauridae	H	4–7 tons	Dong et al., 1983
<i>Wuerhosaurus</i>	Stegosauridae	P	1–4 tons	Dong, 1990b
<i>Ankylosaurus</i>	Ankylosauridae	F	4–7 tons	Coombs, 1978a
<i>Euoplocephalus</i>	Ankylosauridae	M(*)	1–4 tons	Bakker, 1980
<i>Pinacosaurus</i>	Ankylosauridae	H	1–4 tons	Maryanska, 1977
<i>Saichania</i>	Ankylosauridae	H	1–4 tons	Maryanska, 1977
<i>Talarurus</i>	Ankylosauridae	H	0.7–1 ton	Maryanska, 1977
<i>Acanthopholis</i>	Nodosauridae	F	400–700 kg	Galton, 1983
<i>Denversaurus</i>	Nodosauridae	P	1–4 tons	Bakker, 1980
<i>Edmontonia</i>	Nodosauridae	P	1–4 tons	Carpenter, 1990
<i>Hoplitosaurus</i>	Nodosauridae	F	1–4 tons	Galton, 1983
<i>Hylaeosaurus</i>	Nodosauridae	H	0.7–1 ton	Mantell, 1841
<i>Nodosaurus</i>	Nodosauridae	P	1–4 tons	Marsh, 1889
<i>Panoplosaurus</i>	Nodosauridae	M(*)	1–4 tons	Colbert, 1962; Coombs, 1978b
<i>Sauropelta</i>	Nodosauridae	F	1–4 tons	Coombs, 1978a
<i>Silvisaurus</i>	Nodosauridae	F, P	1–4 tons	Eaton, 1960
<i>Heterodontosaurus</i>	Heterodontosauridae	F, P	1–4 tons	Santa Luca et al., 1976
<i>Drinker</i> (juv.)	Hypsilophodontidae	F	1–4 kg	Bakker et al., 1990
<i>Fulgorotherium</i>	Hypsilophodontidae	F	10–40 kg	Huene, 1932
<i>Gongbasaurus</i>	Hypsilophodontidae	P	10–40 kg	Dong, 1989
<i>Hypsilophodon</i>	Hypsilophodontidae	F	40–70 kg	Galton and Jensen, 1975
<i>Leaellynosaura</i>	Hypsilophodontidae	F	7–10 kg	Rich and Rich, 1989
<i>Orodromeus</i>	Hypsilophodontidae	M(*)	7–10 kg	Dunham et al., 1989
<i>Othnielia</i>	Hypsilophodontidae	f(*)	7–10 kg	Dodson et al., 1980
<i>Parksosaurus</i>	Hypsilophodontidae	F(*)	40–70 kg	Anderson et al., 1985
<i>Thescelosaurus</i>	Hypsilophodontidae	F(*)	100–400 kg	Anderson et al., 1985
<i>Yandusaurus</i>	Hypsilophodontidae	F, P	4–7 kg	He, 1979
<i>Tenontosaurus</i>	(Ornithopoda)	F	1–4 tons	Forster, 1990
<i>Dryosaurus</i>	Dryosauridae	F	100–400 kg	Galton, 1981
<i>Valdosaurus</i>	Dryosauridae	F	10–40 kg	Galton and Taquet, 1982
<i>Camptosaurus</i>	Camptosauridae	M(*)	400–700 kg	Bakker, 1980
<i>Iguanodon</i>	Iguanodontidae	M(*)	4–7 tons	Colbert, 1962

## APPENDIX

*Continued.*

Genus	Higher taxon	Method	Weight	Reference
<i>Ouranosaurus</i>	Iguanodontidae	F, P	1–4 tons	Taquet, 1976
<i>Muttabarrosaurus</i>	(Iguanodontia)	F	1–4 tons	Bartholomai and Molnar, 1981
<i>Rhabdodon</i>	(Iguanodontia)	F	1–4 tons	Lapparent, 1947
<i>Gilmoresaurus</i>	Hadrosauridae	P	100–400 kg	Brett-Surman, 1980
<i>Tanais</i>	Hadrosauridae	F	1–4 tons	Wiman, 1929
<i>Claosaurus</i>	Hadrosauridae	F, P	0.7–1 ton	Lull and Wright, 1942
<i>Hadrosaurus</i>	Hadrosauridae	F	1–4 tons	Galton and Jensen, 1978
<i>Kritosaurus</i>	Hadrosauridae	F	1–4 tons	Lull and Wright, 1942
<i>Maiaasaura</i>	Hadrosauridae	M(*)	4–7 tons	Dunham et al., 1989
<i>Prosaurolophus</i>	Hadrosauridae	F, P	1–4 tons	Lull and Wright, 1942
<i>Saurolophus</i>	Hadrosauridae	F	1–4 tons	Lull and Wright, 1942
<i>Lophorhothon</i>	Hadrosauridae	F	0.7–1 ton	Langston, 1960
<i>Edmontosaurus</i>	Hadrosauridae	F(*)	1–4 tons	Anderson et al., 1985
<i>“Anatosaurus”</i>	Hadrosauridae	F(*)	1–4 tons	Anderson et al., 1985
<i>Anatotitan</i>	Hadrosauridae	F, P	1–4 tons	Chapman and Brett-Surman, 1990
<i>Bactrosaurus</i>	Hadrosauridae	F, P	1–4 tons	Gilmore, 1933
<i>Corythosaurus</i>	Hadrosauridae	M(*)	4–7 tons	Colbert, 1962
<i>Hypacrosaurus</i>	Hadrosauridae	F(*)	1–4 tons	Anderson et al., 1985
<i>Lambeosaurus</i>	Hadrosauridae	F	1–4 tons	Morris, 1981
<i>Parasaurolophus</i>	Hadrosauridae	F	1–4 tons	Lull and Wright, 1942
<i>Tsintaosaurus</i>	Hadrosauridae	F	1–4 tons	Young, 1958
<i>Goyocephale</i>	Homalocephalidae	P	10–40 kg	Perle et al., 1982
<i>Homalocephale</i>	Homalocephalidae	F	10–40 kg	Maryanska and Osmolska, 1974
<i>Graviolithus</i>	Pachycephalosauridae	E	70–100 kg	Giffin, 1989
<i>Majungatholus</i>	Pachycephalosauridae	E	100–400 kg	Giffin, 1989
<i>Ornatolithus</i>	Pachycephalosauridae	E	70–100 kg	Giffin, 1989
<i>Pachycephalosaurus</i>	Pachycephalosauridae	E	0.7–1 ton	Giffin, 1989
<i>Prenocephale</i>	Pachycephalosauridae	F	70–100 kg	Maryanska and Osmolska, 1974
<i>Stegoceras</i>	Pachycephalosauridae	F	10–40 kg	Gilmore, 1924
<i>Stygomoloch</i>	Pachycephalosauridae	E	70–100 kg	Giffin, 1989
<i>Psittacosaurus</i>	Psittacosauridae	F	40–70 kg	Sereno and Shichin, 1988
<i>Bagaceratops</i>	Protoceratopsidae	F	7–10 kg	Maryanska and Osmolska, 1975
<i>Leptoceratops</i>	Protoceratopsidae	F, H(*)	100–400 kg	Anderson et al., 1985
<i>Microceratops</i>	Protoceratopsidae	F	4–7 kg	Maryanska and Osmolska, 1975
<i>Montanoceratops</i>	Protoceratopsidae	F	400–700 kg	Brown and Schlaikjer, 1942
<i>Protoceratops</i>	Protoceratopsidae	M(*)	100–400 kg	Colbert, 1962
<i>Avaceratops</i> (juv.)	Ceratopsidae	F, P	400–700 kg	Dodson, 1986
<i>Brachyceratops</i>	Ceratopsidae	F, P	100–400 kg	Lull, 1933
<i>Centrosaurus</i>	Ceratopsidae	F, P	1–4 tons	Lull, 1933
<i>Monoclonius</i>	Ceratopsidae	F, P	1–4 tons	Lull, 1933
<i>Pachyrhinosaurus</i>	Ceratopsidae	H	1–4 tons	Langston, 1975
<i>Styracosaurus</i>	Ceratopsidae	F, H(*)	1–4 tons	Anderson et al., 1985
<i>Anchiceratops</i>	Ceratopsidae	F, H	7–10 tons	Lull, 1933
<i>Chasmosaurus</i>	Ceratopsidae	P	1–4 tons	Lull, 1933
<i>Pentaceratops</i>	Ceratopsidae	M(*)	1–4 tons	Bakker, 1980
<i>Triceratops</i>	Ceratopsidae	M(*)	7–10 tons	Colbert, 1962