

Adventures in Scientific Computing: A Safari for Pure Mathematicians

Adam Larios



6 June 2019

Western Washington University
Bellingham, WA, USA

Outline

- 1 An Invitation to Computation: Some Observations
- 2 Example Problem: $A\vec{x} = \vec{b}$
- 3 Example Problem: Meshing
- 4 Example Problem: Variational Methods

Outline

- 1 An Invitation to Computation: Some Observations
- 2 Example Problem: $A\vec{x} = \vec{b}$
- 3 Example Problem: Meshing
- 4 Example Problem: Variational Methods

“The purpose of computing isn’t numbers. It is insight.”

-R.W. Hamming



- A reasonable question: Are computers beneath you?

- A reasonable question: Are computers beneath you?

[illegible]

- A reasonable question: Are computers beneath you?

```

FE30- 20 B4 FC 90 F7 60 B1 3C
#FDEDL
F0- 6C 00 JMP ($0036)
F1- 6C 00 MFP #F000
F2- 6C 00 MFP #F000
F3- 6C 00 MFP #F000
F4- 6C 00 MFP #F000
F5- 6C 00 MFP #F000
F6- 6C 00 MFP #F000
F7- 6C 00 MFP #F000
F8- 6C 00 MFP #F000
F9- 6C 00 MFP #F000
FA- 6C 00 MFP #F000
FB- 6C 00 MFP #F000
FC- 6C 00 MFP #F000
FD- 6C 00 MFP #F000
FE- 6C 00 MFP #F000
FF- 6C 00 MFP #F000


```



- A reasonable question: Are computers beneath you?

[illegible]

- A reasonable question: Are computers beneath you?

[illegible]

Question: Why learn about computers when one has mathematics?

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$

- $f'(x) = \frac{1}{1+x}$

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$
- $f'(x) = \frac{1}{1+x}$
- $f'(x) = \frac{1}{1+x^2}$

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$
- $f'(x) = \frac{1}{1+x}$
- $f'(x) = \frac{1}{1+x^2}$
- $f'(x) = \frac{1}{1+x^{2.1}}$

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$
- $f'(x) = \frac{1}{1+x}$
- $f'(x) = \frac{1}{1+x^2}$
- $f'(x) = \frac{1}{1+x^{2.1}}$

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$
- $f'(x) = \frac{1}{1+x}$
- $f'(x) = \frac{1}{1+x^2}$
- $f'(x) = \frac{1}{1+x^{2.1}}$

Exercise. **Approximate:**

- $\int_0^t 1 + x \, dx$
- $\int_0^t \frac{1}{1+x} \, dx$
- $\int_0^t \frac{1}{1+x^2} \, dx$
- $\int_0^t \frac{1}{1+x^{2.1}} \, dx$

Question: Why learn about computers when one has mathematics?

Exercise. Find $f(x)$ such that:

- $f'(x) = 1 + x$
- $f'(x) = \frac{1}{1+x}$
- $f'(x) = \frac{1}{1+x^2}$
- $f'(x) = \frac{1}{1+x^{2.1}}$

Exercise. **Approximate:**

- $\int_0^t 1 + x \, dx$
- $\int_0^t \frac{1}{1+x} \, dx$
- $\int_0^t \frac{1}{1+x^2} \, dx$
- $\int_0^t \frac{1}{1+x^{2.1}} \, dx$

Observation 1

- Exactness is a specialized tool.

Problem: Solve $x^2 - 2 = 0$.

Problem: Solve $x^2 - 2 = 0$.

Solution (?):

- $x = \pm\sqrt{2}$

Problem: Solve $x^2 - 2 = 0$.

Solution (?):

- $x = \pm\sqrt{2}$
- $x \approx \pm 1.414213562373 \dots$

Problem: Solve $x^2 - 2 = 0$.

Solution (?):

- $x = \pm\sqrt{2}$
- $x \approx \pm 1.414213562373 \dots$
- $x_0 = 1$ and then $x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}$

Problem: Solve $x^2 - 2 = 0$.

Solution (?):

- $x = \pm\sqrt{2}$
- $x \approx \pm 1.414213562373 \dots$
- $x_0 = 1$ and then $x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}$

Problem: Solve $x^2 - 2 = 0$.

Solution (?):

- $x = \pm\sqrt{2}$
- $x \approx \pm 1.414213562373 \dots$
- $x_0 = 1$ and then $x_{n+1} = x_n - \frac{x_n^2 - 2}{2x_n}$

Observation 2

- Exactness is often a *placeholder* for an algorithm.

Problem: Solve $y'' - 4y' + 13y = 0$; $y(0) = y'(0) = 1$

Problem: Solve $y'' - 4y' + 13y = 0$; $y(0) = y'(0) = 1$

Solution:

- $y(t) = e^{-2t} \sin(3t) - e^{-2t} \cos(3t).$

Problem: Solve $y'' - 4y' + 13y = 0$; $y(0) = y'(0) = 1$


Solution:

- $y(t) = e^{-2t} \sin(3t) - e^{-2t} \cos(3t).$

Problem: Solve $y'' - 4y' + 13y = 0$; $y(0) = y'(0) = 1$

Solution:


- $y(t) = e^{-2t} \sin(3t) - e^{-2t} \cos(3t).$



Problem: Solve $y'' - 4y' + 13y = 0$; $y(0) = y'(0) = 1$

Solution:

- $y(t) = e^{-2t} \sin(3t) - e^{-2t} \cos(3t).$



Observation 3

- Solutions are not always this simple...

The Incompressible Navier-Stokes Equations



Claude L.M.H. Navier

Momentum Equation

$$\underbrace{\frac{\partial}{\partial t} \mathbf{u}}_{\text{Acceleration}} + \underbrace{(\mathbf{u} \cdot \nabla) \mathbf{u}}_{\text{Advection}} = \underbrace{-\nabla p}_{\text{Pressure Gradient}} + \underbrace{\nu \Delta \mathbf{u}}_{\text{Viscous Diffusion}} + \underbrace{\mathbf{f}}_{\text{Body Force}}$$

Continuity Equation

$$\nabla \cdot \mathbf{u} = 0$$



George G. Stokes

Unknowns

$\mathbf{u} :=$ Velocity (vector)

$p :=$ Pressure (scalar)

Parameter

$\nu :=$ Kinematic Viscosity

Problem (Leray 1933)

Existence and uniqueness of strong solutions in 3D for all time. (\$1,000,000 Clay Millennium Prize Problem)

J. Fluid Mech. (1983), vol. 130, pp. 411–452

Printed in Great Britain

411

Small-scale structure of the Taylor–Green vortex

By **MARC E. BRACHET**[†], **DANIEL I. MEIRON**,
STEVEN A. ORSZAG,

Massachusetts Institute of Technology, Cambridge, MA 02139

B. G. NICKEL,

University of Guelph, Guelph, Ontario

RUDOLF H. MORF

R.C.A. Laboratories, Zurich, Switzerland

AND **URIEL FRISCH**

CNRS, Observatoire de Nice, 06-Nice, France

(Received 5 February 1982 and in revised form 14 June 1982)

The dynamics of both the inviscid and viscous Taylor–Green (TG) three-dimensional vortex flows are investigated. This flow is perhaps the simplest system in which one can study the generation of small scales by three-dimensional vortex stretching and the resulting turbulence. The problem is studied by both direct spectral numerical solution of the Navier–Stokes equations (with up to 256^3 modes) and by power-series analysis in time.




FIGURE 20. A plot of the maximum vorticity ω_{\max}^2 as a function of the distance from the walls of the impermeable cube for $t = 1-9$ (1). Here we compute ω_{\max}^2 over the faces of subcubes nested in such a way that the distance from the nearest wall of the impermeable cube is d . Thus $d = 0$ corresponds to the maximum vorticity on the faces of the impermeable cube while $d = \frac{1}{2}\pi$ corresponds to the vorticity at the centre of this cube. Observe that for early times the vorticity is concentrated near the walls of the impermeable cube. As t increases beyond 6, there is significant vorticity generation in the main body of the cube.

Commun. Math. Phys. 94, 61–66 (1984)

Remarks on the Breakdown of Smooth Solutions for the 3-*D* Euler Equations

J. T. Beale^{1*}, T. Kato^{2**}, and A. Majda^{2***}

1 Department of Mathematics, Duke University, Durham, NC 27701, USA

2 Department of Mathematics, University of California, Berkeley, CA 94720, USA

Abstract. The authors prove that the maximum norm of the vorticity controls the breakdown of smooth solutions of the 3-*D* Euler equations. In other words, if a solution of the Euler equations is initially smooth and loses its regularity at some later time, then the maximum vorticity necessarily grows without bound as the critical time approaches; equivalently, if the vorticity remains bounded, a smooth solution persists.

The motion of an ideal incompressible fluid is governed by a system of partial differential equations known as the Euler equations. For two-dimensional flow,

- Thus, computations can give mathematicians *insight*.

- Thus, computations can give mathematicians *insight*.
- Even if we don't want to use computations ourselves, it can benefit us to learn how they work (and learn to recognize when they *fail*.)

Outline

- 1 An Invitation to Computation: Some Observations
- 2 Example Problem: $A\vec{x} = \vec{b}$
- 3 Example Problem: Meshing
- 4 Example Problem: Variational Methods

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Let's use a top-of-the-line Intel i7 processor:
 $\approx 3 \times 10^{11}$ multiplications/divisions per second!

N Time

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Let's use a top-of-the-line Intel i7 processor:
 $\approx 3 \times 10^{11}$ multiplications/divisions per second!

| <u>N</u> | <u>Time</u> |
|----------|-------------|
| 100 | $1\mu s$ |

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Let's use a top-of-the-line Intel i7 processor:
 $\approx 3 \times 10^{11}$ multiplications/divisions per second!

| <u>N</u> | <u>Time</u> |
|----------|----------------|
| 100 | $1\mu\text{s}$ |
| 1000 | 0.003s |

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Let's use a top-of-the-line Intel i7 processor:
 $\approx 3 \times 10^{11}$ multiplications/divisions per second!

| <u>N</u> | <u>Time</u> |
|----------|----------------|
| 100 | $1\mu\text{s}$ |
| 1000 | 0.003s |
| 10^6 | 12 days |

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Let's use a top-of-the-line Intel i7 processor:
 $\approx 3 \times 10^{11}$ multiplications/divisions per second!

| <u>N</u> | <u>Time</u> |
|----------|----------------|
| 100 | $1\mu\text{s}$ |
| 1000 | 0.003s |
| 10^6 | 12 days |
| 10^7 | 35 years |

Problem: Solve $A\vec{x} = \vec{b}$, where A is an invertible matrix.

(Note: We don't bother to find or store A^{-1} .)

Gaussian Elimination

- Works for any matrix!
- Algorithm is “unstable”: tiny round-off errors are strongly amplified.
- For an $N \times N$ matrix, cost $\approx \frac{1}{3}N^3$ multiplications/divisions.

Let's use a top-of-the-line Intel i7 processor:
 $\approx 3 \times 10^{11}$ multiplications/divisions per second!

| <u>N</u> | <u>Time</u> |
|----------|-------------------|
| 100 | $1\mu\text{s}$ |
| 1000 | 0.003s |
| 10^6 | 12 days |
| 10^7 | 35 years |
| 10^9 | 35 million years! |

One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{cases} 5x + 6y + 4z = 3 \\ 7x + 9y + 3z = 1 \\ 2x + 1y + 8z = 2 \end{cases}$$

One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{cases} 5x + 6y + 4z = 3 \\ 7x + 9y + 3z = 1 \\ 2x + 1y + 8z = 2 \end{cases}$$

$$\begin{cases} 5x = 3 - 6y - 4z \\ 9y = 1 - 7x - 3z \\ 8z = 2 - 2x - 1y \end{cases}$$

One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{cases} 5x + 6y + 4z = 3 \\ 7x + 9y + 3z = 1 \\ 2x + 1y + 8z = 2 \end{cases}$$

$$\begin{cases} 5x = 3 - 6y - 4z \\ 9y = 1 - 7x - 3z \\ 8z = 2 - 2x - 1y \end{cases}$$

$$\begin{cases} x = \frac{1}{5}(3 - 6y - 4z) \\ y = \frac{1}{9}(1 - 7x - 3z) \\ z = \frac{1}{8}(2 - 2x - 1y) \end{cases}$$

One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{cases} 5x + 6y + 4z = 3 \\ 7x + 9y + 3z = 1 \\ 2x + 1y + 8z = 2 \end{cases}$$

$$\begin{cases} 5x = 3 - 6y - 4z \\ 9y = 1 - 7x - 3z \\ 8z = 2 - 2x - 1y \end{cases}$$

$$\begin{cases} x = \frac{1}{5}(3 - 6y - 4z) \\ y = \frac{1}{9}(1 - 7x - 3z) \\ z = \frac{1}{8}(2 - 2x - 1y) \end{cases}$$



One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{cases} 5x + 6y + 4z = 3 \\ 7x + 9y + 3z = 1 \\ 2x + 1y + 8z = 2 \end{cases}$$

$$\begin{cases} 5x = 3 - 6y - 4z \\ 9y = 1 - 7x - 3z \\ 8z = 2 - 2x - 1y \end{cases}$$

$$\begin{cases} x = \frac{1}{5}(3 - 6y - 4z) \\ y = \frac{1}{9}(1 - 7x - 3z) \\ z = \frac{1}{8}(2 - 2x - 1y) \end{cases}$$



$$\begin{cases} x_{n+1} = \frac{1}{5}(3 - 6y_n - 4z_n) \\ y_{n+1} = \frac{1}{9}(1 - 7x_n - 3z_n) \\ z_{n+1} = \frac{1}{8}(2 - 2x_n - 1y_n) \end{cases}$$

One alternative to Gaussian Elimination

$$\begin{bmatrix} 5 & 6 & 4 \\ 7 & 9 & 3 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{cases} 5x + 6y + 4z = 3 \\ 7x + 9y + 3z = 1 \\ 2x + 1y + 8z = 2 \end{cases}$$


$$\begin{cases} 5x = 3 - 6y - 4z \\ 9y = 1 - 7x - 3z \\ 8z = 2 - 2x - 1y \end{cases}$$


$$\begin{cases} x = \frac{1}{5}(3 - 6y - 4z) \\ y = \frac{1}{9}(1 - 7x - 3z) \\ z = \frac{1}{8}(2 - 2x - 1y) \end{cases}$$



$$\begin{cases} x_{n+1} = \frac{1}{5}(3 - 6y_n - 4z_n) \\ y_{n+1} = \frac{1}{9}(1 - 7x_n - 3z_n) \\ z_{n+1} = \frac{1}{8}(2 - 2x_n - 1y_n) \end{cases}$$

This is Jacobi's iteration method.





Jacobi Iteration In Detail

Problem: Solve $A\vec{x} = \vec{b}$ via

$$\text{diag}(A)\vec{x}_{n+1} + (A - \text{diag}(A))\vec{x}_n = \vec{b}.$$

Rewrite as:

$$\begin{aligned}\vec{x}_{n+1} &= (\text{diag}(A))^{-1}(\vec{b} - (A - \text{diag}(A))\vec{x}_n) \\ &= (\text{diag}(A))^{-1}(\vec{b} - A\vec{x}_n) + \vec{x}_n \\ &= \vec{x}_n + (\text{diag}(A))^{-1}(\vec{b} - A\vec{x}_n)\end{aligned}$$

Jacobi Iteration In Detail

Problem: Solve $A\vec{x} = \vec{b}$ via

$$\text{diag}(A)\vec{x}_{n+1} + (A - \text{diag}(A))\vec{x}_n = \vec{b}.$$

Rewrite as:

$$\begin{aligned}\vec{x}_{n+1} &= (\text{diag}(A))^{-1}(\vec{b} - (A - \text{diag}(A))\vec{x}_n) \\ &= (\text{diag}(A))^{-1}(\vec{b} - A\vec{x}_n) + \vec{x}_n \\ &= \vec{x}_n + (\text{diag}(A))^{-1}(\vec{b} - A\vec{x}_n)\end{aligned}$$

Write $B = (\text{diag}(A))^{-1}$. Then

$$\begin{aligned}\vec{x}_{n+1} &= \vec{x}_n + B(\vec{b} - A\vec{x}_n) \\ &= (I - BA)\vec{x}_n + B\vec{b}\end{aligned}$$

Jacobi Iteration In Detail

Problem: Solve $A\vec{x} = \vec{b}$ via

$$\text{diag}(A)\vec{x}_{n+1} + (A - \text{diag}(A))\vec{x}_n = \vec{b}.$$

Rewrite as:

$$\begin{aligned}\vec{x}_{n+1} &= (\text{diag}(A))^{-1}(\vec{b} - (A - \text{diag}(A))\vec{x}_n) \\ &= (\text{diag}(A))^{-1}(\vec{b} - A\vec{x}_n) + \vec{x}_n \\ &= \vec{x}_n + (\text{diag}(A))^{-1}(\vec{b} - A\vec{x}_n)\end{aligned}$$

Write $B = (\text{diag}(A))^{-1}$. Then

$$\begin{aligned}\vec{x}_{n+1} &= \vec{x}_n + B(\vec{b} - A\vec{x}_n) \\ &= (I - BA)\vec{x}_n + B\vec{b}\end{aligned}$$

To generalize, write $Q = B^{-1} = \text{diag}(A)$ and rearrange:

$$Q\vec{x}_{n+1} = (Q - A)\vec{x}_n + \vec{b}$$

Convergence for Iterative Methods

Problem: Solve $A\vec{x} = \vec{b}$.

Algorithm: Choose an invertible, easy-to-compute *preconditioner* matrix B .

Iterate with $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

Convergence for Iterative Methods

Problem: Solve $A\vec{x} = \vec{b}$.

Algorithm: Choose an invertible, easy-to-compute *preconditioner* matrix B .

Iterate with $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

- We can now consider other choices for $B = Q^{-1}$.
- Q is called the *splitting* of A .
- $B = Q^{-1}$ is called the *preconditioner* of A .

Convergence for Iterative Methods

Problem: Solve $A\vec{x} = \vec{b}$.

Algorithm: Choose an invertible, easy-to-compute *preconditioner* matrix B .

Iterate with $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

- We can now consider other choices for $B = Q^{-1}$.
- Q is called the *splitting* of A .
- $B = Q^{-1}$ is called the *preconditioner* of A .

Idea: If $\vec{x}_n \rightarrow \vec{x}$ for some \vec{x} , then:

$$\begin{aligned}\vec{x} &= (I - BA)\vec{x} + B\vec{b} \\ \Rightarrow \vec{x} &= \vec{x} - BA\vec{x} + B\vec{b} \\ \Rightarrow BA\vec{x} &= B\vec{b} \\ \Rightarrow A\vec{x} &= \vec{b}\end{aligned}$$

So \vec{x} is the solution to our problem!

Algorithm: $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

Q: When does the iteration converge?

Algorithm: $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

Q: When does the iteration converge?

Idea: Let error $= \vec{e}_n = \vec{x}_n - \vec{x}$. A little algebra shows,

$$e_{n+1} = (I - BA)e_n.$$

Algorithm: $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

Q: When does the iteration converge?

Idea: Let error $= \vec{e}_n = \vec{x}_n - \vec{x}$. A little algebra shows,

$$e_{n+1} = (I - BA)e_n.$$

Thus,

$$\vec{e}_{n+1} = (I - BA)\vec{e}_n = (I - BA)^2\vec{e}_{n-1} = \dots (I - BA)^n\vec{e}_0.$$

Algorithm: $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

Q: When does the iteration converge?

Idea: Let error $= \vec{e}_n = \vec{x}_n - \vec{x}$. A little algebra shows,

$$\vec{e}_{n+1} = (I - BA)\vec{e}_n.$$

Thus,

$$\vec{e}_{n+1} = (I - BA)\vec{e}_n = (I - BA)^2\vec{e}_{n-1} = \dots (I - BA)^n\vec{e}_0.$$

- Worst-case scenario: When \vec{e}_0 is an eigenvector of $I - BA$ with $|\lambda| > 1$.
- Strategy: Show spectral radius of $I - BA$, $\rho(I - BA) < 1$.

Algorithm: $\vec{x}_{n+1} = (I - BA)\vec{x}_n + B\vec{b}$.

Q: When does the iteration converge?

Idea: Let error $= \vec{e}_n = \vec{x}_n - \vec{x}$. A little algebra shows,

$$\vec{e}_{n+1} = (I - BA)\vec{e}_n.$$

Thus,

$$\vec{e}_{n+1} = (I - BA)\vec{e}_n = (I - BA)^2\vec{e}_{n-1} = \dots (I - BA)^n\vec{e}_0.$$

- Worst-case scenario: When \vec{e}_0 is an eigenvector of $I - BA$ with $|\lambda| > 1$.
- Strategy: Show spectral radius of $I - BA$, $\rho(I - BA) < 1$.
Requires spectral analysis!

Partial Answer: Gershgorin Discs

For any matrix A with eigenvalue λ , choose eigenvector \mathbf{x} so that $x_i = 1$ for some i and $|x_j| \leq 1$ for all j . Rewrite $A\mathbf{x} = \lambda\mathbf{x}$ as

$$\sum_j a_{ij}x_j = \lambda x_i = \lambda.$$

Then, since again $x_i = 1$,

$$\sum_{j \neq i} a_{ij}x_j + a_{ii} = \lambda.$$

Thus,

$$|\lambda - a_{ii}| = \left| \sum_{j \neq i} a_{ij}x_j \right| \leq \sum_{j \neq i} |a_{ij}| =: R_i.$$

Thus, every eigenvalue must lie in one of the Gershgorin discs $D_I := D(a_{ii}, R_i)$!

Theorem: The Jacobi method is convergent for any strictly diagonally dominant matrix A , i.e., $|a_{ii}| > \sum_{j \neq i} |a_{ij}|$.


Proof.

Jacobi matrix is

$$I - BA = \begin{bmatrix} 0 & -\frac{a_{12}}{a_{11}} & -\frac{a_{13}}{a_{11}} & \cdots & -\frac{a_{1n}}{a_{11}} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & -\frac{a_{i1}}{a_{ii}} & \cdots & 0 & \cdots & -\frac{a_{in}}{a_{ii}} \\ \vdots & \vdots & \vdots & & \vdots \end{bmatrix}$$

Gershgorin Discs: $D_i = \left\{ z \in \mathbb{C} : |z - 0| \leq \sum_{j \neq i} \left| \frac{a_{ij}}{a_{ii}} \right| \right\} \subset B(0, 1)$. Thus, $\rho(I - BA) < 1$!


3D Vortex Structures



Mappings

| | | |
|--|---|-----------------------------|
| Mesh generation | ← | Graph Theory |
| Iterative Methods ($A\mathbf{x} = \mathbf{b}$) | ← | Spectral Theory |
| Elliptic Curve Cryptography | ← | Number Theory |
| Finite Element Methods | ← | PDE Theory |
| Visualization | ← | Geometry |
| Parallel Computing | ← | Scheduling and Graph Theory |
| Large matrices | ← | Operator Theory |







Mappings




Outline

- 1 An Invitation to Computation: Some Observations
- 2 Example Problem: $A\vec{x} = \vec{b}$
- 3 Example Problem: Meshing**
- 4 Example Problem: Variational Methods




Meshes



Delaunay Mesh






Delaunay Mesh



Delaunay condition

- Three points p, q, r , in S form a Delaunay triangle if and only if the circumcircle of these points contains no other point of S .

Delaunay Mesh




Delaunay condition

- Three points p, q, r , in S form a Delaunay triangle if and only if the circumcircle of these points contains no other point of S .


Convex hull condition

- Three points p_0, q_0, r_0 in S_0 form a face of the convex hull of S_0 if and only if the plane passing through p_0, q_0, r_0 has all the points of S_0 lying to one side.

Efficient Delaunay Computation



Efficient Delaunay Computation




- Computational problems can lead to new ideas!


Mesh Labeling

Given $f = f(x, y)$, find $u = u(x, y)$ such that $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f$.

Write $u_{i,j} = u(x_i, y_j)$.

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \approx \frac{u_{i,j-1} + u_{i,j+1} + u_{i-1,j} + u_{i+1,j} - 4u_{i,j}}{\Delta x \Delta y}$$








Note

- Tightly “banded” structure is more efficient for computation.

A more challenging mesh









New ideas are needed!

- Efficient preconditioners.




New ideas are needed!

- Efficient preconditioners.
- New iteration methods.



New ideas are needed!





- Efficient preconditioners.
- New iteration methods.
- Ideas that exploit “sparse” structure.



New ideas are needed!

- Efficient preconditioners.
- New iteration methods.
- Ideas that exploit “sparse” structure.
- Mesh Relabeling...

Space Filling Curves



Example Problem: Linear Systems

Outline

- 1 An Invitation to Computation: Some Observations
- 2 Example Problem: $A\vec{x} = \vec{b}$
- 3 Example Problem: Meshing
- 4 Example Problem: Variational Methods

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

- Wrong way: $\mathbf{x} = A^{-1}\mathbf{b}$.

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

- Wrong way: $\mathbf{x} = A^{-1}\mathbf{b}$.
- Better way:

$$Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$$

$$\mathbf{x}_{k+1} = (I - Q^{-1}A)\mathbf{x}_k + Q^{-1}\mathbf{b}$$

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

- Wrong way: $\mathbf{x} = A^{-1}\mathbf{b}$.
- Better way:

$$Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$$

$$\mathbf{x}_{k+1} = (I - Q^{-1}A)\mathbf{x}_k + Q^{-1}\mathbf{b}$$

- Fundamental Theorem of Iterative Methods:
Iteration will converge if and only if spectral radius of $(I - Q^{-1}A)$ is < 1 .

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

- Wrong way: $\mathbf{x} = A^{-1}\mathbf{b}$.
- Better way:

$$Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$$

$$\mathbf{x}_{k+1} = (I - Q^{-1}A)\mathbf{x}_k + Q^{-1}\mathbf{b}$$

- Fundamental Theorem of Iterative Methods:
Iteration will converge if and only if spectral radius of $(I - Q^{-1}A)$ is < 1 .

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

- Wrong way: $\mathbf{x} = A^{-1}\mathbf{b}$.
- Better way:

$$Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$$

$$\mathbf{x}_{k+1} = (I - Q^{-1}A)\mathbf{x}_k + Q^{-1}\mathbf{b}$$

- Fundamental Theorem of Iterative Methods:
Iteration will converge if and only if spectral radius of $(I - Q^{-1}A)$ is < 1 .

Problem 2

- Design Q with $\rho(I - Q^{-1}A) < 1$ so that the iteration converges fast.

Iterative Methods

Problem

- Solve $A\mathbf{x} = \mathbf{b}$.

- Wrong way: $\mathbf{x} = A^{-1}\mathbf{b}$.
- Better way:

$$Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$$

$$\mathbf{x}_{k+1} = (I - Q^{-1}A)\mathbf{x}_k + Q^{-1}\mathbf{b}$$

- Fundamental Theorem of Iterative Methods:
Iteration will converge if and only if spectral radius of $(I - Q^{-1}A)$ is < 1 .

Problem 2

- Design Q with $\rho(I - Q^{-1}A) < 1$ so that the iteration converges fast.
- Now we get to play with spectral theory!

Variation/Minimization Methods

Problem (Again)

- Solve $A\mathbf{x} = \mathbf{b}$.

Variation/Minimization Methods


Problem (Again)

- Solve $A\mathbf{x} = \mathbf{b}$.

Easy 1D version

- Solve $3x = 2$.

- Let $f(x) = \frac{3}{2}x^2 - 2x$.



Variation/Minimization Methods


Problem (Again)

- Solve $A\mathbf{x} = \mathbf{b}$.

Easy 1D version

- Solve $3x = 2$.

- Let $f(x) = \frac{3}{2}x^2 - 2x$.



- Then $f'(x) = 3x - 2$.

Variation/Minimization Methods


Problem (Again)

- Solve $A\mathbf{x} = \mathbf{b}$.

Easy 1D version

- Solve $3x = 2$.

- Let $f(x) = \frac{3}{2}x^2 - 2x$.



- Then $f'(x) = 3x - 2$.
- Minimum occurs when $0 = f'(x) = 3x - 2$, i.e., $3x = 2$.

Variation/Minimization Methods


Problem (Again)

- Solve $A\mathbf{x} = \mathbf{b}$.

Easy 1D version

- Solve $3x = 2$.

- Let $f(x) = \frac{3}{2}x^2 - 2x$.



- Then $f'(x) = 3x - 2$.
- Minimum occurs when $0 = f'(x) = 3x - 2$, i.e., $3x = 2$.
- The **minimizer** is the **solution**!

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.

Convergence?

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!

Convergence?

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:

Convergence?

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:
 - Steepest Descent (Multivariable Calculus)

Convergence?

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:
 - Steepest Descent (Multivariable Calculus)
 - Conjugate Gradient (Krylov Subspaces)

Convergence?

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:
 - Steepest Descent (Multivariable Calculus)
 - Conjugate Gradient (Krylov Subspaces)

Convergence?

- Problem: Rate depends heavily on eigenvalue structure.

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:
 - Steepest Descent (Multivariable Calculus)
 - Conjugate Gradient (Krylov Subspaces)

Convergence?

- Problem: Rate depends heavily on eigenvalue structure.
- Idea: Solve $MA\mathbf{x} = M\mathbf{b}$ (M invertible) instead.

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.

- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:
 - Steepest Descent (Multivariable Calculus)
 - Conjugate Gradient (Krylov Subspaces)

Convergence?

- Problem: Rate depends heavily on eigenvalue structure.
- Idea: Solve $MA\mathbf{x} = M\mathbf{b}$ (M invertible) instead.
- New problem: Find M so that MA has a better eigenvalue structure.

Variation/Minimization Methods

Problem 1

- Solve $A\mathbf{x} = \mathbf{b}$.

Problem 2

- Minimize $\frac{1}{2}A\mathbf{x} \cdot \mathbf{x} - \mathbf{b} \cdot \mathbf{x}$.




- This will work, so long as A is symmetric and positive-definitive.
- We can now have fun with optimization techniques!
- Examples:
 - Steepest Descent (Multivariable Calculus)
 - Conjugate Gradient (Krylov Subspaces)

Convergence?

- Problem: Rate depends heavily on eigenvalue structure.
- Idea: Solve $MA\mathbf{x} = M\mathbf{b}$ (M invertible) instead.
- New problem: Find M so that MA has a better eigenvalue structure.
- **Again we can play with spectral theory!**

Conclusion

- $A\mathbf{x} = \mathbf{b}$
- $Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$





Observations

- Computers can do simple mathematics quickly.

Conclusion

- $A\mathbf{x} = \mathbf{b}$
- $Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$






Observations

- Computers can do simple mathematics quickly.
- One must find creative solutions to make mathematics simple.

Conclusion

- $A\mathbf{x} = \mathbf{b}$
- $Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$





Observations

- Computers can do simple mathematics quickly.
- One must find creative solutions to make mathematics simple.
- Computational math presents new challenges. These problems can open doors to new mathematics!

Conclusion

- $A\mathbf{x} = \mathbf{b}$
- $Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$






Observations

- Computers can do simple mathematics quickly.
- One must find creative solutions to make mathematics simple.
- Computational math presents new challenges. These problems can open doors to new mathematics!

Conclusion

- $A\mathbf{x} = \mathbf{b}$
- $Q\mathbf{x}_{k+1} = (Q - A)\mathbf{x}_k + \mathbf{b}$



Observations

- Computers can do simple mathematics quickly.
- One must find creative solutions to make mathematics simple.
- Computational math presents new challenges. These problems can open doors to new mathematics!

Thank you!