

Math 107-Lecture 26

Dr. Adam Larios

University of Nebraska-Lincoln

Taylor Series

The Taylor series for the function $f(x)$ near $x = a$ is given by:

$$f(a) + f'(a)\frac{(x-a)}{1!} + f''(a)\frac{(x-a)^2}{2!} + \dots + f^{(n)}(a)\frac{(x-a)^n}{n!} + \dots$$

Question: How can we find a Taylor series from a known one? (Taking so many derivatives can be a lot of work....)

Some simple examples (hint: substitution will mostly do it)

Example 1. Find the Taylor series near 0 for the function f below and determine for what values of x it converges.

$$f(x) = \frac{x}{1 - (x/2)}$$

Example 2. Same for g

$$g(x) = \frac{x^2}{2 + x^4}$$

Example 3. How about for h ?

$$h(x) = x^3 e^{x^2}$$

Clicker question

What is the Taylor series for the function $f(x) = \frac{1}{2-x}$ near $x = 1$?

A $\sum_{n=0}^{\infty} (x-1)^n$

B $\sum_{n=0}^{\infty} (2-x)^n$

C $\sum_{n=0}^{\infty} (1-x)^n$

D $\sum_{n=0}^{\infty} (1+x)^n$

E $\sum_{n=0}^{\infty} (2+x)^n$

We can do more. A lot more!

Inside the interval of convergence we can also multiply, integrate, and differentiate a power series **term by term**. Using this approach we can find more easily the Taylor series around the origin for many functions.

Examples 4 & 5:

- $f(x) = \ln(1 - x)$
- $g(x) = \arctan x$

Example 6: Find the first four terms of the Taylor series near the origin for

$$\ln(1 - x^2) \arctan(x^3).$$

Using series for approximation of integrals

Recall that

$$\int e^{x^2} dx$$

cannot be expressed in terms of elementary functions, hence we cannot apply the FTC to compute

$$\int_0^1 e^{x^2} dx.$$

However we can get a good approximation (how good?) for this integral by using Taylor series.