

The Orchestra of Partial Differential Equations

Adam Larios



7 March 2019

Landscape Seminar

Outline

- 1 PDEs at UNL
- 2 The Simplest Partial Differential Equation
- 3 The Diffusion Equation and Fourier Series
- 4 Putting it all together
- 5 Instability and Energy Cascades

Outline

1

PDEs at UNL

2

The Simplest Partial Differential Equation

3

The Diffusion Equation and Fourier Series

4

Putting it all together

5


Instability and Energy Cascades

PDEs at UNL

Professors



Postdocs




PDEs at UNL


Grad Students





Grad Students













Some Famous Linear PDEs I

Laplace equation $-\nabla^2 u = -\sum_{i=1}^n u_{x_i x_i} = 0$

Helmholtz's (or eigenvalue) equation $-\nabla^2 u = \lambda u$

Linear transport equation $u_t + \sum_{i=1}^n b^i u_{x_i} = u_t + u \cdot \nabla b = 0$

Liouville's equation $u_t + \sum_{i=1}^n (b^i u)_{x_i} = u_t + \nabla \cdot (bu) = 0$

Heat (or diffusion) equation $u_t = \nu \nabla^2 u$

Schrödinger's equation $u_t = i \nabla^2 u$

Kolmogorov's equation $u_t - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} = 0$

Fokker-Planck equation $u_t - \sum_{i,j=1}^n (a^{ij} u)_{x_i x_j} + \sum_{i=1}^n (b^i u)_{x_i} = 0$

Some Famous Linear PDEs II

Wave equation	$u_{tt} - c^2 \nabla^2 u = 0$
Klein-Gordon equation	$u_{tt} - c^2 \nabla^2 u + m^2 u = 0$
Telegraph equation	$u_{tt} + 2du_t - c^2 u_{xx} = 0$
General wave equation	$u_{tt} - \sum_{i,j=1}^n a^{ij} u_{x_i x_j} + \sum_{i=1}^n b^i u_{x_i} = 0$
Linear KdV	$u_t + u_{xxx} = 0$
Beam equation	$u_t + u_{xxxx} = 0$

Some Famous Nonlinear PDEs

Eikonal equation	$ \nabla u = 1$
Nonlinear Poisson equation	$-\nabla^2 u = f(u)$
p-Laplacian equation	$\nabla \cdot (\nabla u ^{p-2} \nabla u) = 0$
Minimal surface equation	$\nabla \cdot \left(\frac{\nabla u}{(1 + \nabla u ^2)^{1/2}} \right) = 0$
Monge-Ampère equation	$\det(\nabla \nabla u) = f$
Hamilton-Jacobi equation	$u_t + H(\nabla u, x) = 0$
Scalar conservation law	$u_t + \nabla \cdot (\mathbf{F}(u)) = 0$
Inviscid Burgers' equation	$u_t + uu_x = 0.$
Scalar reaction-diffusion equation	$u_t - \nabla^2 u = f(u)$
Porous medium equation	$u_t - \nabla^2 (u^\gamma) = 0$
Nonlinear wave equation	$u_{tt} - \nabla^2 u + f(u) = 0$
Korteweg-deVries (KdV) equation	$u_t + uu_x + u_{xxx} = 0$
Nonlinear Schrödinger equation	$iu_t + \nabla^2 u f(u ^2)u = 0$

Systems of partial differential equations.

- Equilibrium equations of linear elasticity

$$\mu \nabla^2 \mathbf{u} + (\lambda + \mu) \nabla(\nabla \cdot \mathbf{u}) = \mathbf{0}$$

- Evolution equations of linear elasticity

$$\mathbf{u}_{tt} - \mu \nabla^2 \mathbf{u} + (\lambda + \mu) \nabla(\nabla \cdot \mathbf{u}) = \mathbf{0}$$

- Maxwell's equations

$$\begin{cases} \mathbf{E}_t + \mu_0 \mathbf{J} = \nabla \times \mathbf{B} \\ \mathbf{B}_t = -\nabla \times \mathbf{E} \\ \nabla \cdot \mathbf{E} = q \\ \nabla \cdot \mathbf{B} = 0 \end{cases}$$

Nonlinear Systems.

- System of conservation laws

$$\mathbf{u}_t + \nabla \cdot (\mathbf{F}(\mathbf{u})) = 0$$

- Reaction-diffusion system

$$\mathbf{u}_t - \nu \nabla^2 \mathbf{u} = \mathbf{f}(\mathbf{u})$$

- Euler's equations for incompressible, inviscid flow

$$\begin{cases} \mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p \\ \nabla \cdot \mathbf{u} = 0 \end{cases}$$

- Navier-Stokes equations for incompressible, viscous flow

$$\begin{cases} \mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u} - \nu \nabla^2 \mathbf{u} = -\nabla p + \mathbf{f} \\ \nabla \cdot \mathbf{u} = 0 \end{cases}$$

Modern-Day Fact: Most equations can **only** be solved with computers






Figure: Programmers working on ENIAC, c. 1946


Numerical Solutions of PDEs




Numerical Solutions of PDEs





Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs




Numerical Solutions of PDEs



Numerical Solutions of PDEs



Numerical Solutions of PDEs



What do I study?

Turbulence: A Wide Range of Scales



Turbulence: A Wide Range of Scales



Turbulence: A Wide Range of Scales ($10^{-2}m$)






Image Credit: Animal Flight Group, Dept. of Zoology, Oxford University and Dr. John Young


Turbulence: A Wide Range of Scales ($10^8 m$)



Turbulence: A Wide Range of Scales ($10^9 m$)



Turbulence: A Wide Range of Scales ($10^{21}m$)



The Incompressible Navier-Stokes Equations

Momentum Equation



$$\underbrace{\frac{\partial \vec{u}}{\partial t}}_{\text{Acceleration}} + \underbrace{(\vec{u} \cdot \nabla) \vec{u}}_{\text{Advection}} = \underbrace{-\nabla p}_{\substack{\text{Pressure} \\ \text{Gradient}}} + \underbrace{\nu \nabla^2 \vec{u}}_{\text{Viscous Diffusion}}$$

Continuity Equation (Divergence-Free Condition)

Claude L.M.H. Navier

$$\nabla \cdot \vec{u} = 0$$



George G. Stokes

Unknowns

\vec{u} := Velocity (vector)

p := Pressure (scalar)

Parameter

ν := Kinematic Viscosity

Problem (Leray 1933)

Existence and uniqueness of strong solutions in 3D for all time. (\$1,000,000 Clay Millennium Prize Problem)

Outline

- 1 PDEs at UNL
- 2 The Simplest Partial Differential Equation
- 3 The Diffusion Equation and Fourier Series
- 4 Putting it all together
- 5 Instability and Energy Cascades

The Simplest Partial Differential Equation



$$\frac{d\rho}{dt} =$$

position = $x = x(t)$

velocity = $v = v(t, x(t)) = \frac{dx}{dt}$

density = $\rho = \rho(t, x(t))$

The Simplest Partial Differential Equation



$$\frac{d\rho}{dt} = 0$$

position = $x = x(t)$

velocity = $v = v(t, x(t)) = \frac{dx}{dt}$

density = $\rho = \rho(t, x(t))$

The Simplest Partial Differential Equation



$$\frac{d\rho}{dt} = 0$$

$$\frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + \frac{dx}{dt} \frac{\partial \rho}{\partial x}$$

$$= 0$$

position = $x = x(t)$

velocity = $v = v(t, x(t)) = \frac{dx}{dt}$

density = $\rho = \rho(t, x(t))$

The Simplest Partial Differential Equation



$$\frac{d\rho}{dt} = 0$$

$$\begin{aligned}\frac{d\rho}{dt} &= \frac{\partial \rho}{\partial t} + \frac{dx}{dt} \frac{\partial \rho}{\partial x} \\ &= \frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} = 0\end{aligned}$$

position = $x = x(t)$

velocity = $v = v(t, x(t)) = \frac{dx}{dt}$

density = $\rho = \rho(t, x(t))$

The Simplest Partial Differential Equation



$$\frac{d\rho}{dt} = 0$$

$$\begin{aligned}\frac{d\rho}{dt} &= \frac{\partial \rho}{\partial t} + \frac{dx}{dt} \frac{\partial \rho}{\partial x} \\ &= \frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} = 0\end{aligned}$$

Transport Equation

$$\rho_t + v\rho_x = 0$$

position = $x = x(t)$

velocity = $v = v(t, x(t)) = \frac{dx}{dt}$

density = $\rho = \rho(t, x(t))$

The Simplest Partial Differential Equation



position = $x = x(t)$

velocity = $v = v(t, x(t)) = \frac{dx}{dt}$

density = $\rho = \rho(t, x(t))$

$$\frac{d\rho}{dt} = 0$$

$$\begin{aligned}\frac{d\rho}{dt} &= \frac{\partial \rho}{\partial t} + \frac{dx}{dt} \frac{\partial \rho}{\partial x} \\ &= \frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} = 0\end{aligned}$$

Transport Equation

$$\rho_t + v\rho_x = 0$$

Transport Equation in \mathbb{R}^n

$$\rho_t + (\vec{v} \cdot \nabla) \rho = 0$$

$$(\vec{v} \cdot \nabla) \rho = v_1 \frac{\partial \rho}{\partial x} + v_2 \frac{\partial \rho}{\partial y} + v_3 \frac{\partial \rho}{\partial z}$$




Figure: Programmers working on ENIAC, c. 1946



Idea

What about the water itself?
What if we set $\rho = v = u$?



Idea

What about the water itself?
What if we set $\rho = v = u$?

Burgers' Equation

$$u_t + uu_x = 0$$

Burgers' Equation in \mathbb{R}^n

$$\vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = 0$$

Computer Time!




Figure: Programmers working on ENIAC, one of the first computers (c. 1946)

Outline

- 1 PDEs at UNL
- 2 The Simplest Partial Differential Equation
- 3 The Diffusion Equation and Fourier Series
- 4 Putting it all together
- 5 Instability and Energy Cascades

Diffusion Equation



Diffusion

Diffusion Equation



Diffusion

Concentration = $\theta = \theta(x, t)$


Diffusion Equation

$$\theta_t = \nu \theta_{xx}$$


Diffusion Equation in \mathbb{R}^3

$$\theta_t = \nu(\theta_{xx} + \theta_{yy} + \theta_{zz}) = \nu \nabla^2 \theta$$

Frequency




Frequency




$$\begin{aligned}u(t) = & 0.5 \cos(2t) + 0.125 \cos(8t) + 0.03125 \cos(32t) \\& + 1.0 \sin(1t) + 0.25 \sin(4t) + 0.0625 \sin(16t)\end{aligned}$$

Frequency




$$\begin{aligned}u(t) = & 0.5 \cos(2t) + 0.125 \cos(8t) + 0.03125 \cos(32t) \\& + 1.0 \sin(1t) + 0.25 \sin(4t) + 0.0625 \sin(16t)\end{aligned}$$

Frequency



$$\begin{aligned}u(t) = & 0.5 \cos(2t) + 0.125 \cos(8t) + 0.03125 \cos(32t) \\& + 1.0 \sin(1t) + 0.25 \sin(4t) + \textcolor{red}{0.3125} \sin(16t)\end{aligned}$$

Frequency



$$\begin{aligned}u(t) = & \color{red}{1.5 \cos(2t) + 0.125 \cos(8t) + 0.03125 \cos(32t)} \\& + 1.0 \sin(1t) + 0.25 \sin(4t) + 0.0625 \sin(16t)\end{aligned}$$

Fourier Series



$$u(t) = \sum_{k=0}^{\infty} (a_k \cos(kt) + b_k \sin(kt))$$

Fourier Series



$$u(t) = \sum_{k=0}^{\infty} (a_k \cos(kt) + b_k \sin(kt))$$

$$e^{ikt} = \cos(kt) + i \sin(kt)$$

$$\cos(kt) = \frac{e^{ikt} + e^{-ikt}}{2}$$

$$\sin(kt) = \frac{e^{ikt} - e^{-ikt}}{2i}$$

Fourier Series



$$u(t) = \sum_{k=0}^{\infty} (\textcolor{red}{a}_k \cos(kt) + \textcolor{red}{b}_k \sin(kt))$$

$$e^{ikt} = \cos(kt) + i \sin(kt)$$



$$\cos(kt) = \frac{e^{ikt} + e^{-ikt}}{2}$$

$$\sin(kt) = \frac{e^{ikt} - e^{-ikt}}{2i}$$

$$u(t) = \sum_{k=-\infty}^{\infty} \textcolor{red}{c}_k e^{ikt}$$

$$c_k = \frac{1}{2} (a_k - ib_k), \quad k > 0,$$

$$c_k = \frac{1}{2} (a_k + ib_k), \quad k < 0.$$



Multi-dimensional Fourier series

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx}$$

$$u(\vec{x}) = \sum_{\vec{k} \in \mathbb{Z}^n} \hat{u}_{\vec{k}} e^{i\vec{k} \cdot \vec{x}}$$

Multi-dimensional Fourier series

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx}$$

$$u(\vec{x}) = \sum_{\vec{k} \in \mathbb{Z}^n} \hat{u}_{\vec{k}} e^{i\vec{k} \cdot \vec{x}}$$


$$\hat{u}_{\vec{k}} = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(\vec{x}) e^{-i\vec{k} \cdot \vec{x}} d\vec{x}$$



(a) Original image




(a) Original image




(b) Fourier transform (magnitude)



(a) Original image




(b) Fourier transform (magnitude)




(c) Zero-out Fourier coefficients



(a) Original image



(b) Fourier transform (magnitude)



(c) Zero-out Fourier coefficients



(d) Inverse Fourier transform

Derivatives

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx}$$

Derivatives

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx}$$

$$\frac{d}{dx} u(x) = \sum_{k \in \mathbb{Z}} \cancel{i} k \hat{u}_k e^{ikx}$$

Derivatives

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx}$$

$$\frac{d}{dx} u(x) = \sum_{k \in \mathbb{Z}} ik \hat{u}_k e^{ikx}$$

$$\frac{d^2}{dx^2} u(x) = \sum_{k \in \mathbb{Z}} (-k^2) \hat{u}_k e^{ikx}$$

Derivatives

$$u(x) = \sum_{k \in \mathbb{Z}} \hat{u}_k e^{ikx}$$

$$\frac{d}{dx} u(x) = \sum_{k \in \mathbb{Z}} ik \hat{u}_k e^{ikx}$$

$$\frac{d^2}{dx^2} u(x) = \sum_{k \in \mathbb{Z}} (-k^2) \hat{u}_k e^{ikx}$$



Idea

Can the Fourier transform be used to understand differential equations?

Diffusion Equation and the Fourier Transform

$$u_t = \nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} \widehat{u}_k(t) e^{ikx}$$

Diffusion Equation and the Fourier Transform

$$u_t = \nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} \widehat{u}_k(t) e^{ikx}$$

$$u_{xx} = \sum_{k \in \mathbb{Z}} (-k^2) \widehat{u}_k e^{ikx}$$

Diffusion Equation and the Fourier Transform

$$u_t = \nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} \widehat{u}_k(t) e^{ikx}$$

$$u_{xx} = \sum_{k \in \mathbb{Z}} (-k^2) \widehat{u}_k e^{ikx}$$

$$u_t = \sum_{k \in \mathbb{Z}} \frac{d}{dt} \widehat{u}_k e^{ikx}$$

Diffusion Equation and the Fourier Transform

$$u_t = \nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} \widehat{u}_k(t) e^{ikx}$$

$$u_{xx} = \sum_{k \in \mathbb{Z}} (-k^2) \widehat{u}_k e^{ikx}$$

$$u_t = \sum_{k \in \mathbb{Z}} \frac{d}{dt} \widehat{u}_k e^{ikx}$$

Fourier Coefficients

$$\frac{d}{dt} \widehat{u}_k = -\nu k^2 \widehat{u}_k, \quad k \in \mathbb{Z}$$

Diffusion Equation and the Fourier Transform

$$u_t = \nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} \widehat{u}_k(t) e^{ikx}$$

$$u_{xx} = \sum_{k \in \mathbb{Z}} (-k^2) \widehat{u}_k e^{ikx}$$

$$u_t = \sum_{k \in \mathbb{Z}} \frac{d}{dt} \widehat{u}_k e^{ikx}$$

Fourier Coefficients

$$\frac{d}{dt} \widehat{u}_k = -\nu k^2 \widehat{u}_k, \quad k \in \mathbb{Z} \quad \Rightarrow \quad \widehat{u}_k(t) = e^{-\nu k^2 t} \widehat{u}_k(0)$$

Diffusion Equation and the Fourier Transform

$$u_t = \nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} \widehat{u}_k(t) e^{ikx}$$

$$u_{xx} = \sum_{k \in \mathbb{Z}} (-k^2) \widehat{u}_k e^{ikx}$$

$$u_t = \sum_{k \in \mathbb{Z}} \frac{d}{dt} \widehat{u}_k e^{ikx}$$

Fourier Coefficients

$$\frac{d}{dt} \widehat{u}_k = -\nu k^2 \widehat{u}_k, \quad k \in \mathbb{Z} \quad \Rightarrow \quad \widehat{u}_k(t) = e^{-\nu k^2 t} \widehat{u}_k(0)$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} e^{-\nu k^2 t} \widehat{u}_k(0) e^{ikx}$$

Computer Time Again!



Figure: Technician working on a Cray supercomputer. (c. 1986)

Outline

- 1 PDEs at UNL
- 2 The Simplest Partial Differential Equation
- 3 The Diffusion Equation and Fourier Series
- 4 Putting it all together
- 5 Instability and Energy Cascades

Burgers Equation: Well-posedness

Burgers Equation: Well-posedness

Q: What prevents the development of a singularity?

Burgers Equation: Well-posedness

Q: What prevents the development of a singularity?

$$u_t + uu_x = \nu u_{xx}$$

Burgers Equation: Well-posedness

Q: What prevents the development of a singularity?

$$u_t + uu_x = \nu u_{xx}$$

Take derivative $w = u_x$.

$$u_{xt} + u_x u_x + uu_{xx} = \nu u_{xxx}$$

$$w_t + w^2 + uw_x = \nu w_{xx}$$

If w is at a *maximum* . . .

Burgers Equation: Well-posedness

Q: What prevents the development of a singularity?

$$u_t + uu_x = \nu u_{xx}$$

Take derivative $w = u_x$.

$$u_{xt} + u_x u_x + uu_{xx} = \nu u_{xxx}$$

$$w_t + w^2 + uw_x = \nu w_{xx}$$

If w is at a *maximum* . . . $w_x = 0$ and $w_{xx} < 0$.

Burgers Equation: Well-posedness

Q: What prevents the development of a singularity?

$$u_t + uu_x = \nu u_{xx}$$

Take derivative $w = u_x$.

$$u_{xt} + u_x u_x + uu_{xx} = \nu u_{xxx}$$

$$w_t + w^2 + uw_x = \nu w_{xx}$$

If w is at a *maximum* . . . $w_x = 0$ and $w_{xx} < 0$.

$$w_t + w^2 + 0 < 0$$

$$w_t < -w^2$$

Burgers Equation: Well-posedness

Q: What prevents the development of a singularity?

$$u_t + uu_x = \nu u_{xx}$$

Take derivative $w = u_x$.

$$u_{xt} + u_x u_x + uu_{xx} = \nu u_{xxx}$$

$$w_t + w^2 + uw_x = \nu w_{xx}$$

If w is at a *maximum* . . . $w_x = 0$ and $w_{xx} < 0$.

$$w_t + w^2 + 0 < 0$$

$$w_t < -w^2$$

So w ($= u_x$) is decreasing at any maximum!

Burgers Equation: Higher dimensions

n-Dimensional Burgers

$$\vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u}$$

Burgers Equation: Higher dimensions

n-Dimensional Burgers

$$\vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u}$$

Olga Ladyzhenskaya: Maximum Principle for $\frac{1}{2}|\vec{u}|^2$.



Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers		
Navier-Stokes 2D		
Navier-Stokes 3D		

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	
Navier-Stokes 2D		
Navier-Stokes 3D		

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D		
Navier-Stokes 3D		

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D		
Navier-Stokes 3D		

Hidden Maximum Principle in 2D, $\vec{u} = (u_1, u_2)$.

$$\omega = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \quad (\text{Vorticity})$$

$$\omega_t + (\vec{u} \cdot \nabla) \omega = \nu \nabla^2 \omega$$

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D	✓	
Navier-Stokes 3D		

Hidden Maximum Principle in 2D, $\vec{u} = (u_1, u_2)$.

$$\omega = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \quad (\text{Vorticity})$$

$$\omega_t + (\vec{u} \cdot \nabla) \omega = \nu \nabla^2 \omega$$

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D	✓	
Navier-Stokes 3D		

Hidden Maximum Principle in 2D, $\vec{u} = (u_1, u_2)$.

$$\omega = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \quad (\text{Vorticity})$$

$$\omega_t + (\vec{u} \cdot \nabla) \omega = \nu \nabla^2 \omega$$

3D Vorticity, $\vec{\omega} = \nabla \times \vec{u}$.

$$\vec{\omega}_t + (\vec{u} \cdot \nabla) \vec{\omega} = (\vec{\omega} \cdot \nabla) \vec{u} + \nu \nabla^2 \vec{\omega}$$

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D	✓	
Navier-Stokes 3D	???	

Hidden Maximum Principle in 2D, $\vec{u} = (u_1, u_2)$.

$$\omega = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \quad (\text{Vorticity})$$

$$\omega_t + (\vec{u} \cdot \nabla) \omega = \nu \nabla^2 \omega$$

3D Vorticity, $\vec{\omega} = \nabla \times \vec{u}$.

$$\vec{\omega}_t + (\vec{u} \cdot \nabla) \vec{\omega} = (\vec{\omega} \cdot \nabla) \vec{u} + \nu \nabla^2 \vec{\omega}$$

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D	✓	✓
Navier-Stokes 3D	???	

Hidden Maximum Principle in 2D, $\vec{u} = (u_1, u_2)$.

$$\omega = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \quad (\text{Vorticity})$$

$$\omega_t + (\vec{u} \cdot \nabla) \omega = \nu \nabla^2 \omega$$

3D Vorticity, $\vec{\omega} = \nabla \times \vec{u}$.

$$\vec{\omega}_t + (\vec{u} \cdot \nabla) \vec{\omega} = (\vec{\omega} \cdot \nabla) \vec{u} + \nu \nabla^2 \vec{\omega}$$

Back to Navier-Stokes

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

When do we have no singularity?

	$\nu > 0$	$\nu = 0$
Burgers	✓	✗
Navier-Stokes 2D	✓	✓
Navier-Stokes 3D	???	???

Hidden Maximum Principle in 2D, $\vec{u} = (u_1, u_2)$.

$$\omega = \frac{\partial u_2}{\partial x} - \frac{\partial u_1}{\partial y} \quad (\text{Vorticity})$$

$$\omega_t + (\vec{u} \cdot \nabla) \omega = \nu \nabla^2 \omega$$

3D Vorticity, $\vec{\omega} = \nabla \times \vec{u}$.

$$\vec{\omega}_t + (\vec{u} \cdot \nabla) \vec{\omega} = (\vec{\omega} \cdot \nabla) \vec{u} + \nu \nabla^2 \vec{\omega}$$

Focus: 3D Euler Equations

Focus: 3D Euler Equations (Effective viscosity?)

$$\vec{\omega}_t + (\vec{u} \cdot \nabla) \vec{\omega} = (\vec{\omega} \cdot \nabla) \vec{u}, \quad \vec{\omega} = \nabla \times \vec{u}.$$

Focus: 3D Euler Equations (Effective viscosity?)

$$\vec{\omega}_t + (\vec{u} \cdot \nabla) \vec{\omega} = (\vec{\omega} \cdot \nabla) \vec{u}, \quad \vec{\omega} = \nabla \times \vec{u}.$$



Image Credit: J. Bell, D. Marcus, *Comm. Math. Phys.* **147**, 371-394 (1992).

Outline

- 1 PDEs at UNL
- 2 The Simplest Partial Differential Equation
- 3 The Diffusion Equation and Fourier Series
- 4 Putting it all together
- 5 Instability and Energy Cascades

Backward Diffusion

$$u_t = -\nu u_{xx}$$

Backward Diffusion

$$u_t = -\nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} e^{+\nu k^2 t} \hat{u}_k(0) e^{ikx}$$

Backward Diffusion

$$u_t = -\nu u_{xx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} e^{+\nu k^2 t} \hat{u}_k(0) e^{ikx}$$

Massively unstable!

Fourth-Order Diffusion

$$u_t = -\nu u_{xxxx}$$

Fourth-Order Diffusion

$$u_t = -\nu u_{xxxx}$$

$$u_{xxxx} = \sum_{k \in \mathbb{Z}} (k^4) \hat{u}_k e^{ikx}$$

Fourth-Order Diffusion

$$u_t = -\nu u_{xxxx}$$

$$u_{xxxx} = \sum_{k \in \mathbb{Z}} (k^4) \hat{u}_k e^{ikx}$$

$$u(x, t) = \sum_{k \in \mathbb{Z}} e^{-\nu k^4 t} \hat{u}_k(0) e^{ikx}$$

Massively stable!

Nonlinear Equations

Burgers Equation [Shock Waves, Traffic]

$$u_t + uu_x = \nu u_{xx}$$

Navier-Stokes Equations [Incompressible Fluids]

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

Nonlinear Equations

Burgers Equation [Shock Waves, Traffic]

$$u_t + uu_x = \nu u_{xx}$$

Navier-Stokes Equations [Incompressible Fluids]

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

Kuramoto-Sivashinsky (KS) Equation [Flames]

$$u_t + uu_x = -\lambda u_{xx} - u_{xxxx}$$

Nonlinear Equations

Burgers Equation [Shock Waves, Traffic]

$$u_t + uu_x = \nu u_{xx}$$

Navier-Stokes Equations [Incompressible Fluids]

$$\begin{cases} \vec{u}_t + (\vec{u} \cdot \nabla) \vec{u} = \nu \nabla^2 \vec{u} - \nabla p \\ \nabla \cdot \vec{u} = 0 \end{cases}$$

Kuramoto-Sivashinsky (KS) Equation [Flames]

$$u_t + uu_x = -\lambda u_{xx} - u_{xxxx}$$

Computer Time Once More! (Kuramoto-Sivashinsky)



Figure: Hopper Cray XE6 at NERSC, named after American computer scientist Dr. Grace Hopper, 1906-1992.

The Incompressible Navier-Stokes Equations



Claude L.M.H. Navier



George G. Stokes

Momentum Equation

$$\underbrace{\frac{\partial \vec{u}}{\partial t}}_{\textit{Acceleration}} + \underbrace{(\vec{u} \cdot \nabla) \vec{u}}_{\textit{Advection}} = \underbrace{-\nabla p}_{\textit{Pressure Gradient}} + \underbrace{\nu \nabla^2 \vec{u}}_{\textit{Viscous Diffusion}}$$

Continuity Equation (Divergence-Free Condition)

$$\nabla \cdot \vec{u} = 0$$

Unknowns

\vec{u} := Velocity (vector)

p := Pressure (scalar)

Parameter

ν := Kinematic Viscosity

Thank you!

