

# Why do we need a rule

Calculus 1

Day 11 3.3

The Quotient  
Rule

Kevin

Gonzales PhD

Today we will consider functions of the form  $\frac{f(x)}{g(x)}$ .

For example, let us consider  $F(x) = \frac{4x^4}{x^2}$ .

$$\frac{d}{dx} \left( \frac{4x^4}{x^2} \right) = \frac{d}{dx} (4x^2) = 8x$$

What if we had tried to just take the derivative of each?

$$\frac{\frac{d}{dx}(4x^4)}{\frac{d}{dx}(x^2)} = \frac{16x^3}{2x} = 8x^2$$

# Finding a rule

Let's consider the function  $Q(x) = \frac{f(x)}{g(x)}$ . Of course we have to avoid points where  $g(x) = 0$ . Rewrite as  $f(x) = Q(x)g(x)$  and use the product rule.

$$f'(x) = Q'(x)g(x) + Q(x)g'(x)$$

Replace  $Q(x) = f(x)/g(x)$ :

$$f'(x) = Q'(x)g(x) + \frac{f(x)}{g(x)}g'(x)$$

Then solve for  $Q'(x)$ :

$$Q'(x) = \frac{f'(x) - \frac{f(x)}{g(x)}g'(x)}{g(x)}$$

Multiply top and bottom to get:

$$Q'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

# Quotient Rule

## Quotient Rule

If  $u = f(x)$  and  $v = g(x)$  are differentiable, then

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

or equivalently,

$$\frac{dv}{dx} \left(\frac{u}{v}\right) = \frac{\frac{du}{dx}v - u\frac{dv}{dx}}{v^2}.$$

In words:

The derivative of a quotient is the derivative of the numerator times the denominator minus the numerator times the derivative of the denominator, all over the denominator squared.

## Example

Consider the function

$$F(x) = \frac{4 \sin(x)}{x^2 + 1}.$$

Find the derivative  $F'(x)$ .

Here  $F(x) = \frac{f(x)}{g(x)}$ , thus we use the quotient rule:

$$f'(x) = \frac{(4 \cos(x))(x^2 + 1) - (4 \sin(x))(2x)}{(x^2 + 1)^2}.$$

## Clicker Question

Differentiate the function

$$g(x) = \frac{5x^2}{x^3 + 1}.$$

a)  $\frac{10x}{3x^2}$

b)  $\frac{-5x^4 + 10x}{(x^3 + 1)^2}$

c)  $\frac{10x - 15x^4}{(x^3 + 1)}$


d)  $\frac{10x(x^3 + 1) + 5x^2(3x^2)}{(x^3 + 1)^2}$

# Clicker Question

Calculus 1  
Day 11 3.3  
The Quotient  
Rule

Kevin  
Gonzales PhD

If  $h(x) = \frac{f(x)}{g(x)}$ , what is  $h'(1)$ ? In the graph  $f(x)$  is blue and  $g(x)$  is red.



- a) Not enough information
- b)  $h'(1) = 5$
- c)  $h'(1) = -3$
- d)  $h'(1) = -4$

# Derivation of Derivative of Tangent

We have already seen that

$$\frac{d}{dx}(\tan(x)) = \frac{1}{\cos^2(x)}.$$

Now we want to use the quotient rule to derive it:

$$\begin{aligned}\frac{d}{dx}(\tan(x)) &= \frac{d}{dx} \left( \frac{\sin(x)}{\cos(x)} \right) \\ &= \frac{\cos(x) \cos(x) - (\sin(x))(-\sin(x))}{(\cos(x))^2} \\ &= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} = \frac{1}{\cos^2(x)}\end{aligned}$$

# Derivative of Cotangent

Calculus 1  
Day 11 3.3  
The Quotient  
Rule

Kevin  
Gonzales PhD

Show that:

$$\frac{d}{dx}(\cot(x)) = \frac{-1}{\sin^2(x)}$$



## Calculus 1

## Day 11 3.3

The Quotient  
Rule

Kevin

Gonzales PhD

For each of the following, calculate the derivative.

$$\textcircled{1} \quad \frac{3x^2 + 3x}{x + 1}$$

$$\textcircled{2} \quad \frac{x^3 + 1}{e^x}$$

$$\textcircled{3} \quad \frac{4^x}{2x + 1}$$