

MTH 532 Homework 10

Roy Howie

April 6, 2017

4.2 Exterior Algebra

Exercise 1

Suppose $T \in \Lambda^p(V^*)$ and $v_1, \dots, v_p \in V$ are linearly dependent, then there are constants $\{a^1, \dots, a^p\}$, not all zero, such that $a^i v_i = 0$. Without loss of generality, suppose $v_1 = a^2 v_2 + \dots + a^p v_p$. Therefore,

$$\begin{aligned} T(v_1, \dots, v_p) &= T(a^2 v_2 + \dots + a^p v_p, v_2, \dots, v_p) \\ &= a^2 T(v_2, v_2, \dots, v_p) + \dots + a^p T(v_p, v_2, \dots, v_p) \\ &= 0 + \dots + 0 \\ &= 0 \end{aligned}$$

□

Exercise 2

Suppose $\phi_1, \dots, \phi_p \in V^*$ are linearly dependent, then there are constants a^1, \dots, a^p , not all zero, such that $a^i \phi_i = 0$. That is, without loss of generality, ϕ_1 can be written as the sum $c^2 \phi_2 + \dots + c^p \phi_p$. Recall $\phi_i \wedge \phi_i = 0$, $c \wedge \phi_i = c \phi_i$ for constant c , and that the wedge product distributes over addition. Therefore,

$$\begin{aligned} \phi_1 \wedge \dots \wedge \phi_p &= (c^2 \phi_2 + \dots + c^p \phi_p) \wedge \phi_2 \wedge \dots \wedge \phi_p \\ &= c^2 \phi_2 \wedge \phi_2 \wedge \dots \wedge \phi_p + \dots + c^p \phi_p \wedge \phi_2 \wedge \dots \wedge \phi_p \\ &= 0 + \dots + 0 \\ &= 0 \end{aligned}$$

□

Exercise 6

- (a) Let $A: V \rightarrow V$ be a linear isomorphism which sends the basis $B = \{v_1, \dots, v_k\}$ to $B' = \{v'_1, \dots, v'_k\}$.

If B and B' are equivalently oriented, then $\det A$ is positive. Next, for $T \in \Lambda^k(V)$, note that $A^*T = \det(A)T$. However, A^*T was also defined to be $T(Av_1, \dots, Av_k) = T(v'_1, \dots, v'_k)$. Thus, $T(B)$ and $T(B')$ have the same sign.

Conversely, suppose $T(B)$ and $T(B')$ have the same sign. Note that $T(B') = A^*T = \det(A)T(B)$, implying that $\det A$ is positive. Therefore, B and B' are equivalently oriented.

- (b) Let $T, S \in \Lambda^k(V^*)$ be two nonzero elements. Suppose $B = \{v_1, \dots, v_k\}$ is a positively oriented ordered basis. Without loss of generality, suppose $T(B)$ is positive and $S(B)$ is negative. Let $A: V \rightarrow V$ be a linear isomorphism, then $T(B)$ and $S(AB)$ should have the same sign iff $\det A$ is negative. From **4.2.6a**, this is clearly the case, as $S(AB) = S(Av_1, \dots, Av_k) = A^*S(B) = \det(A)S(B)$. This produces a natural orientation, as we have defined two equivalence classes: positive and negative.
- (c) Let $T \in \Lambda^k(V^*)$ be nonzero. Let B and B' be two ordered bases. We define an orientation on V by saying B and B' are equivalently oriented iff $T(B)$ and $T(B')$ have the same sign. Let $A: V \rightarrow V$ be a linear isomorphism such that $A(B) = B'$.

Do the opposite of **4.2.6b**. Note $T(B') = A^*T(B) = \det(A)T(B)$. Thus, $T(B')$ and $T(B)$ have the same sign iff $\det A$ is positive, implying that B and B' are equivalently oriented. □