

Quantum Circuits with Classical and Quantum Control of Causal Orders

Alastair A. Abbott

joint work with

Julian Wechs, Hippolyte Dourdent and Cyril Branciard

Université de Genève, Geneva, Switzerland




Zurich, 28 May 2019



**UNIVERSITÉ
DE GENÈVE**

Causal Information Processing

In quantum information one usually assume a fixed, classical causal structure



■ Does this unnecessarily constrain quantum information processing?

Outline

Quantum Circuit Framework

Causally Indefinite Processes

- Quantum switch

- Process matrix framework

Quantum Circuits with Classical and Quantum Controls of Causal Order


- Classically controlled circuits

- Coherently (quantum) controlled circuits

Quantum Circuits

Quantum circuits, a.k.a. quantum combs, quantum networks

- [Chiribella, D'Ariano, Perinotti, PRL (2008) & PRA (2009)]




- Transform unknown operations $\mathcal{A}_1, \dots, \mathcal{A}_N$ into $\mathcal{W}(\mathcal{A}_1, \dots, \mathcal{A}_N)$
 - Formally they're **supermaps**
- Powerful framework for computing properties of unknown operations, distributed computation, tomography, metrology etc.

What maps can quantum circuits implement and how?

Quantum Circuits

Quantum circuits, a.k.a. quantum combs, quantum networks

- [Chiribella, D'Ariano, Perinotti, PRL (2008) & PRA (2009)]




- Transform unknown operations $\mathcal{A}_1, \dots, \mathcal{A}_N$ into $\mathcal{W}(\mathcal{A}_1, \dots, \mathcal{A}_N)$
 - Formally they're **supermaps**
- Powerful framework for computing properties of unknown operations, distributed computation, tomography, metrology etc.

What maps can quantum circuits implement and how?

Quantum Circuits with Fixed Causal Order

Most general quantum circuit can be implemented as:



\mathcal{M}_i CPTP maps and α_i ancillary systems:


- e.g., $\mathcal{M}_{n+1} : A_n^O \alpha_n \rightarrow A_{n+1}^I \alpha_{n+1}$ for $1 \leq n \leq N-1$

Useful to represent operations as matrices in Choi picture:

- $M = \mathcal{I} \otimes \mathcal{M}(|\mathbb{1}\rangle\langle\mathbb{1}|)$ with $|\mathbb{1}\rangle = \sum_i |i\rangle \otimes |i\rangle$
- \mathcal{M} completely positive $\iff M \geq 0$
- \mathcal{M} trace-preserving $\iff \text{tr}_2 M = \mathbb{1}^1$

[Chiribella, D'Ariano, Perinotti, PRA (2009)]

Quantum Circuits with Fixed Causal Order



Inverse Choi isomorphism given by the **link product**:

- $\mathcal{M}(\rho) = M^{12} * \rho^1 := \text{tr}_1[(\rho^1 \otimes \mathbb{1}^2) \cdot M^{T_1}]$
- Allows channels to be composed, etc.


For input ρ output is

$$M_{N+1} * A_N * \dots * M_2 * A_1 * M_1 * \rho = \underbrace{(M_1 * M_2 * \dots * M_{N+1})}_W * (\rho \otimes A_1 \otimes \dots \otimes A_N)$$

- The action of the circuit is characterised by W :

$$W = M_1 * M_2 * \dots * M_{N+1} = \text{Tr}_{\alpha_1 \dots \alpha_N} [M_1 \cdot M_2^{T_{\alpha_1}} \dots M_{N+1}^{T_{\alpha_N}}]$$


Quantum Circuits with Fixed Causal Order



$$W = M_1 * M_2 * \dots * M_{N+1} = \text{Tr}_{\alpha_1 \dots \alpha_N} [M_1 \cdot M_2^{T_{\alpha_1}} \dots M_{N+1}^{T_{\alpha_N}}]$$

- W is the Choi representation of the supermap describing the circuit
- The $W^{A_1 \prec \dots \prec A_N}$ that represent a circuit with causal order $A_1 \prec \dots \prec A_N$ can be characterised via SDP constraints
 - From any W satisfying constraints one can extract the implementation effectively

Beyond Quantum Circuits?



Is the fixed causal structure of quantum circuits necessary?

- Convex mixture of circuits with different orders is physical, but still a classical causal structure
- More generally, can have classical “dynamical” causal orders

[Oreshkov & Giarmatzi, NJP 2016; Abbott *et al.*, PRA 2016.]


Why not a quantum superposition of causal orders?

- What justification do we have for restricting the causal structure?
- In a quantum theory of gravity, spacetime itself would be quantum

[Oreshkov, Costa, Brukner, *Nat. Commun.* 2012]

Quantum Switch


The “**Quantum Switch**” uses a control system to apply two operations in a superposition of different orders



- A and B act on the target system and are each **only applied once**
- No well defined causal order between A and B : **causally indefinite**
- Implemented in several experiments
 - [Procopio *et al.*, Nat. Commun. 6 (2015)]; [Rubino *et al.*, Sci. Adv. 3 (2017)]; [Goswami *et al.*, PRL 121 (2018)]; [Pan *et al.*, PRL (2019)]

Quantum Switch


The “**Quantum Switch**” uses a control system to apply two operations in a superposition of different orders



- A and B act on the target system and are each **only applied once**
- No well defined causal order between A and B : **causally indefinite**
- Implemented in several experiments
 - [Procopio *et al.*, Nat. Commun. 6 (2015)]; [Rubino *et al.*, Sci. Adv. 3 (2017)]; [Goswami *et al.*, PRL 121 (2018)]; [Pan *et al.*, PRL (2019)]

Quantum Switch


The “**Quantum Switch**” uses a control system to apply two operations in a superposition of different orders



- A and B act on the target system and are each **only applied once**
- No well defined causal order between A and B : **causally indefinite**
- Implemented in several experiments
 - [Procopio *et al.*, Nat. Commun. 6 (2015)]; [Rubino *et al.*, Sci. Adv. 3 (2017)]; [Goswami *et al.*, PRL 121 (2018)]; [Pan *et al.*, PRL (2019)]

Quantum Switch


The “**Quantum Switch**” uses a control system to apply two operations in a superposition of different orders



- A and B act on the target system and are each **only applied once**
- No well defined causal order between A and B : **causally indefinite**
- Implemented in several experiments
 - [Procopio *et al.*, Nat. Commun. 6 (2015)]; [Rubino *et al.*, Sci. Adv. 3 (2017)]; [Goswami *et al.*, PRL 121 (2018)]; [Pan *et al.*, PRL (2019)]

Quantum Switch


The “Quantum Switch” uses a control system to apply two operations in a superposition of different orders



- A and B act on the target system and are each **only applied once**
- No well defined causal order between A and B : **causally indefinite**
- Implemented in several experiments
 - [Procopio *et al.*, Nat. Commun. 6 (2015)]; [Rubino *et al.*, Sci. Adv. 3 (2017)]; [Goswami *et al.*, PRL 121 (2018)]; [Pan *et al.*, PRL (2019)]


Quantum Switch

The “Quantum Switch” uses a control system to apply two operations in a superposition of different orders



- A and B act on the target system and are each **only applied once**
- No well defined causal order between A and B : **causally indefinite**
- Implemented in several experiments
 - [Procopio *et al.*, Nat. Commun. 6 (2015)]; [Rubino *et al.*, Sci. Adv. 3 (2017)]; [Goswami *et al.*, PRL 121 (2018)]; [Pan *et al.*, PRL (2019)]

Causally Indefinite Information Processing



Is the quantum switch actually useful?

- If A and B are unitary transformations:


$$\begin{aligned} \frac{1}{\sqrt{2}}(|H\rangle^c + |V\rangle^c) \otimes |\psi\rangle^t &\rightarrow \frac{1}{\sqrt{2}}(|H\rangle^c \otimes BA|\psi\rangle + |V\rangle^c \otimes AB|\psi\rangle) \\ &= |+\rangle^c \otimes \frac{1}{2}\{A, B\}|\psi\rangle^t + |-\rangle^c \otimes \frac{1}{2}[A, B]|\psi\rangle^t \end{aligned}$$

- Can then distinguish $\{A, B\} = 0$ and $[A, B] = 0$ by measuring the control
- Any quantum circuit needs to use A or B twice to do this
 - A simple query complexity advantage

Causally Indefinite Information Processing

Can generalise the quantum switch the N operations ($N!$ permutations)

- Query complexity: $O(N^2)$ advantage on a generalisation of the commutation/anticommutation problem [Araújo *et al.*, PRL 113 (2014)]
- Perfect channel discrimination [Chiribella, PRA 86 (2012)]
- Communication complexity: $O(2^N)$ advantage computing a distributed function with the quantum switch [Guérin *et al.*, PRL 117 (2016)]




Causal indefiniteness is a resource for quantum information

- But how to generalise and study more rigorously?

Process matrix formalism

The process matrix framework introduced by Oreshkov, Costa and Brukner to describe more general processes without assuming a global causal structure


- Characterises the W that represent general, consistent supermaps



$$M = W * (A_1 \otimes \cdots \otimes A_N) = \text{Tr}_{A_N^{IO}} [W(A_1^T \otimes \cdots \otimes A_N^T \otimes \mathbb{1}^{PF})]$$

- Circuits are a special case: fixed-order process

Causal Nonseparability




- Framework permits formal definition – and characterisation – of process matrices that are **causally nonseparable**
 - E.g., for two operations, **causal separability** means

$$W = q W^{A \prec B} + (1 - q) W^{B \prec A}$$

- General definition more subtle and only recently clarified and properly characterised [Wechs, AA, Branciard, NJP (2019)]
- Quantum switch can be written as a process matrix
 - Easily proven to be causally nonseparable

Beyond the Quantum Switch



“Top-down” definition means not clear if all W are physical

- Quantum switch is only noncausal process we know how to implement

Are there other classes of interesting noncausal processes?

Outline

Quantum Circuit Framework

Causally Indefinite Processes

- Quantum switch

- Process matrix framework


Quantum Circuits with Classical and Quantum Controls of Causal Order

- Classically controlled circuits

- Coherently (quantum) controlled circuits


Classically Controlled Circuits

- First generalise circuits to allow for **classical** control of causal structure
 - Encompass mixed and dynamic causal orders
 - Oreshkov & Giarmatzi proposed **quantum circuits with classical control of causal order (QC-CCs)** as most general causal process



- At each time slot t_n exactly one operation A_{k_n} is applied
- $\{M_{(k_1, \dots, k_n)}^{\rightarrow k_{n+1}}\}_{k_{n+1}}$ are instruments: $\sum_{k_{n+1}} M_{(k_1, \dots, k_n)}^{\rightarrow k_{n+1}}$ is CPTP
- Crucial requirement: **each operation applied once and only once**, irrespective of the operations themselves

Process Matrix of a QC-CC



For input ρ , when operations applied in order k_1, \dots, k_N , output is

$$\begin{aligned}
 & M_{(k_1, \dots, k_N)}^{\rightarrow F} * A_{k_N} * M_{(k_1, \dots, k_{N-1})}^{\rightarrow k_N} * \dots * M_{(k_1, k_2)}^{\rightarrow k_3} * A_{k_2} * M_{(k_1)}^{\rightarrow k_2} * A_{k_1} * M_{\emptyset}^{\rightarrow k_1} * \rho \\
 &= \underbrace{M_{\emptyset}^{\rightarrow k_1} * M_{(k_1)}^{\rightarrow k_2} * M_{(k_1, k_2)}^{\rightarrow k_3} * \dots * M_{(k_1, \dots, k_{N-1})}^{\rightarrow k_N} * M_{(k_1, \dots, k_N)}^{\rightarrow F}}_{\widetilde{W}_{(k_1, \dots, k_N, F)}} * (\rho \otimes A_1 \otimes \dots \otimes A_N)
 \end{aligned}$$


Process matrix of a QC-CC (Causally Separable)

$$W = \sum_{(k_1, \dots, k_N)} \widetilde{W}_{(k_1, \dots, k_N, F)}$$

■ Can be characterised in terms of SDP constraints

Alternative Descriptions of QC-CCs

- Conditioning can be included in operations by introducing **(classical) control system** $\llbracket (k_1, \dots, k_n) \rrbracket^{C_n} := |(k_1, \dots, k_n)\rangle\langle (k_1, \dots, k_n)|^{C_n}$




- Operations now given by the CPTP maps

$$\tilde{M}_1 := \sum_{k_1} \tilde{M}_{\emptyset}^{\rightarrow k_1} \otimes \llbracket (k_1) \rrbracket^{C_1},$$

$$\tilde{M}_{n+1} := \sum_{k_1, \dots, k_n, k_{n+1}} \tilde{M}_{(k_1, \dots, k_n)}^{\rightarrow k_{n+1}} \otimes \llbracket (k_1, \dots, k_n) \rrbracket^{C_n} \otimes \llbracket (k_1, \dots, k_n, k_{n+1}) \rrbracket^{C_{n+1}}$$

$$M_{N+1} := \sum_{k_1, \dots, k_N} M_{(k_1, \dots, k_N)}^{\rightarrow F} \otimes \llbracket (k_1, \dots, k_N) \rrbracket^{C_N}$$

Alternative Descriptions of QC-CCs



- Defining global operations $\tilde{A}_n := \bigoplus_{k_n \in \mathcal{N}} A_{k_n}$ we have

$$\tilde{M}_{N+1} * \tilde{A}_N * \tilde{M}_N * \dots * \tilde{A}_1 * \tilde{M}_1 * \rho = \underbrace{\sum_{k_1, \dots, k_N} W_{(k_1, \dots, k_N, F)}}_W * (\rho \otimes A_1 \otimes \dots \otimes A_N)$$


- Note that wlog we can take all operations to be purified isometries

$$M_{(k_1, \dots, k_{n-1})}^{\rightarrow k_n} = |V_{(k_1, \dots, k_{n-1})}^{\rightarrow k_n}\rangle\rangle\langle\langle V_{(k_1, \dots, k_{n-1})}^{\rightarrow k_n}|$$

- Suggests natural generalisation to quantum control of causal order

From Classical to Coherent Control

- Relax the control state to store only *which* operations have been performed, **but not their order**: $|\mathcal{K}_{n-1}, k_n\rangle^{C_n}$
 - Conditioning on $\mathcal{K} = \{k_1, \dots, k_{n-1}\}$ allows **different orders** to **interfere**
 - Storing full history $|(k_1, \dots, k_n)\rangle^{C_n}$ is more restrictive and included in this case by using ancillas




- Operations given by the isometries V_n with pure CJ representation

$$|\tilde{V}_{n+1}\rangle := \sum_{\substack{\mathcal{K}_{n-1} \\ k_n, k_{n+1}}} |\tilde{V}_{\mathcal{K}_{n-1}, k_n}^{\rightarrow k_{n+1}}\rangle \otimes |\mathcal{K}_{n-1}, k_n\rangle^{C_n} \otimes |\mathcal{K}_n, k_{n+1}\rangle^{C_{n+1}},$$

$$|\tilde{V}_1\rangle := \sum_{k_1} |\tilde{V}_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle \otimes |\emptyset, k_1\rangle^{C_1}, \quad |\tilde{V}_{N+1}\rangle := \sum_{k_N} |\tilde{V}_{\mathcal{N} \setminus \{k_N\}, k_N}^{\rightarrow F}\rangle \otimes |\mathcal{N} \setminus \{k_N\}, k_N\rangle^{C_N}$$

From Classical to Coherent Control

- Relax the control state to store only *which* operations have been performed, **but not their order**: $|\mathcal{K}_{n-1}, k_n\rangle^{C_n}$
 - Conditioning on $\mathcal{K} = \{k_1, \dots, k_{n-1}\}$ allows **different orders** to **interfere**
 - Storing full history $|(k_1, \dots, k_n)\rangle^{C_n}$ is more restrictive and included in this case by using ancillas




- Operations given by the isometries V_n with pure CJ representation

$$|\tilde{V}_{n+1}\rangle := \sum_{\substack{\mathcal{K}_{n-1} \\ k_n, k_{n+1}}} |\tilde{V}_{\mathcal{K}_{n-1}, k_n}^{\rightarrow k_{n+1}}\rangle \otimes |\mathcal{K}_{n-1}, k_n\rangle^{C_n} \otimes |\mathcal{K}_n, k_{n+1}\rangle^{C_{n+1}},$$

$$|\tilde{V}_1\rangle := \sum_{k_1} |\tilde{V}_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle \otimes |\emptyset, k_1\rangle^{C_1}, \quad |\tilde{V}_{N+1}\rangle := \sum_{k_N} |\tilde{V}_{\mathcal{N} \setminus \{k_N\}, k_N}^{\rightarrow F}\rangle \otimes |\mathcal{N} \setminus \{k_N\}, k_N\rangle^{C_N}$$

Coherently (Quantum) Controlled Circuits




- Each $V_{\mathcal{K}_{n-1}, k_n}^{to k_{n+1}} : \mathcal{H}^{A_{k_n}^O \alpha_n} \rightarrow \mathcal{H}^{A_{k_{n+1}}^I \alpha_{n+1}}$ embedded in larger space
- Control ensures that **each party applied once and only once**

For input $|\psi\rangle$, circuit applies transformation

$$|\tilde{V}_{N+1}\rangle\rangle * |\tilde{A}_N\rangle\rangle * |\tilde{V}_N\rangle\rangle * \cdots * |\tilde{V}_2\rangle\rangle * |\tilde{A}_1\rangle\rangle * |\tilde{V}_1\rangle\rangle * |\psi\rangle \in \mathcal{H}^{F\alpha_{N+1}}$$

with “pure link product” $|a\rangle^A * |b\rangle^B := \langle\mathbb{1}|^{A \cap B} (|a\rangle \otimes |b\rangle) = \sum_i \langle i, i |^{(A \cap B)^{\otimes 2}} (|a\rangle \otimes |b\rangle)$

QCs with Quantum Control of Causal Order




$$\begin{aligned}
 & |\tilde{V}_{N+1}\rangle * |\tilde{A}_N\rangle * |\tilde{V}_N\rangle * \dots * |\tilde{V}_2\rangle * |\tilde{A}_1\rangle * |\tilde{V}_1\rangle * |\psi\rangle \\
 &= \sum_{k_1, \dots, k_N} \underbrace{|V_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle * |V_{\emptyset, k_1}^{\rightarrow k_2}\rangle * |V_{\{k_1\}, k_2}^{\rightarrow k_3}\rangle * \dots * |V_{\{k_1, \dots, k_{N-1}\}, k_N}^{\rightarrow F}\rangle}_{|w(k_1, \dots, k_N, F)\rangle} * (|\psi\rangle \otimes |A_1\rangle \otimes \dots \otimes |A_N\rangle)
 \end{aligned}$$

Process matrix of a QC-QC

$$W = \text{Tr}_{\alpha_{N+1}} |w\rangle\langle w|, \quad \text{with} \quad |w\rangle := \sum_{k_1, \dots, k_N} |w(k_1, \dots, k_N, F)\rangle$$

- Can again be characterised in terms of SDP constraints


QC-QC Summary



- Classically controlled QC-QCs recovered as a special case
 - But in general QC-QCs can be causally nonseparable
- QC-QCs are physically realisable, e.g., with a “quantum router”
- Realisation – in terms of the \tilde{V}_n – can be effectively obtained from the any W satisfying the characterisation
- A framework within which to look for new examples and applications of noncausal processes

Example: Quantum Switch

Quantum switch is a QC-QC: $\mathcal{H}^P = \mathcal{H}^{P_t} \otimes \mathcal{H}^{P_c}$ and $\mathcal{H}^F = \mathcal{H}^{F_t} \otimes \mathcal{H}^{F_c}$



The controlled operations are

$$|V_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle = |k_1\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_1^I}, \quad |V_{\emptyset, k_1}^{\rightarrow k_2}\rangle = |\mathbb{1}\rangle^{A_{k_1}^O A_{k_2}^I}, \quad |V_{\{k_1\}, k_2}^{\rightarrow F}\rangle = |k_1\rangle^{F_c} |\mathbb{1}\rangle^{A_{k_2}^O F_t}$$

Process vector is then


$$\begin{aligned} |w_s\rangle &:= |w_{(P, A_1, A_2, F)}\rangle + |w_{(P, A_2, A_1, F)}\rangle \\ &= |V_{\emptyset, \emptyset}^{\rightarrow A_1}\rangle * |V_{\emptyset, A_1}^{\rightarrow A_2}\rangle * |V_{\{A_1\}, A_2}^{\rightarrow F}\rangle + |V_{\emptyset, \emptyset}^{\rightarrow A_2}\rangle * |V_{\emptyset, A_2}^{\rightarrow A_1}\rangle * |V_{\{A_2\}, A_1}^{\rightarrow F}\rangle \\ &= |1\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_1^I} |\mathbb{1}\rangle^{A_1^O A_2^I} |\mathbb{1}\rangle^{A_2^O F_t} |1\rangle^{F_c} + |2\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_2^I} |\mathbb{1}\rangle^{A_2^O A_1^I} |\mathbb{1}\rangle^{A_1^O F_t} |2\rangle^{F_c} \end{aligned}$$

Standard process matrix of switch recovered as $W_{\text{switch}} = |w_s\rangle\langle w_s|$

$$\text{Tr}_F W_{\text{switch}} = \underbrace{|1\rangle\langle 1|^{P_c} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{P_t A_1^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_1^O A_2^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_2^O}}_{W^{A_1 \prec A_2}} + \underbrace{|2\rangle\langle 2|^{P_c} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{P_t A_2^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_2^O A_1^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_1^O}}_{W^{A_2 \prec A_1}}$$

Example: Quantum Switch

Quantum switch is a QC-QC: $\mathcal{H}^P = \mathcal{H}^{P_t} \otimes \mathcal{H}^{P_c}$ and $\mathcal{H}^F = \mathcal{H}^{F_t} \otimes \mathcal{H}^{F_c}$



The controlled operations are

$$|V_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle = |k_1\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_1^I}, \quad |V_{\emptyset, k_1}^{\rightarrow k_2}\rangle = |\mathbb{1}\rangle^{A_{k_1}^O A_2^I}, \quad |V_{\{k_1\}, k_2}^{\rightarrow F}\rangle = |k_1\rangle^{F_c} |\mathbb{1}\rangle^{A_{k_2}^O F_t}$$

Process vector is then


$$\begin{aligned} |w_s\rangle &:= |w_{(P, A_1, A_2, F)}\rangle + |w_{(P, A_2, A_1, F)}\rangle \\ &= |V_{\emptyset, \emptyset}^{\rightarrow A_1}\rangle * |V_{\emptyset, A_1}^{\rightarrow A_2}\rangle * |V_{\{A_1\}, A_2}^{\rightarrow F}\rangle + |V_{\emptyset, \emptyset}^{\rightarrow A_2}\rangle * |V_{\emptyset, A_2}^{\rightarrow A_1}\rangle * |V_{\{A_2\}, A_1}^{\rightarrow F}\rangle \\ &= |1\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_1^I} |\mathbb{1}\rangle^{A_1^O A_2^I} |\mathbb{1}\rangle^{A_2^O F_t} |1\rangle^{F_c} + |2\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_2^I} |\mathbb{1}\rangle^{A_2^O A_1^I} |\mathbb{1}\rangle^{A_1^O F_t} |2\rangle^{F_c} \end{aligned}$$

Standard process matrix of switch recovered as $W_{\text{switch}} = |w_s\rangle\langle w_s|$

$$\text{Tr}_F W_{\text{switch}} = \underbrace{|1\rangle\langle 1|^{P_c} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{P_t A_1^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_1^O A_2^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_2^O}}_{W^{A_1 \prec A_2}} + \underbrace{|2\rangle\langle 2|^{P_c} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{P_t A_2^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_2^O A_1^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_1^O}}_{W^{A_2 \prec A_1}}$$

Example: Quantum Switch

Quantum switch is a QC-QC: $\mathcal{H}^P = \mathcal{H}^{P_t} \otimes \mathcal{H}^{P_c}$ and $\mathcal{H}^F = \mathcal{H}^{F_t} \otimes \mathcal{H}^{F_c}$



The controlled operations are

$$|V_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle = |k_1\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_1^I}, \quad |V_{\emptyset, k_1}^{\rightarrow k_2}\rangle = |\mathbb{1}\rangle^{A_{k_1}^O A_2^I}, \quad |V_{\{k_1\}, k_2}^{\rightarrow F}\rangle = |k_1\rangle^{F_c} |\mathbb{1}\rangle^{A_{k_2}^O F_t}$$

Process vector is then

$$\begin{aligned} |w_s\rangle &:= |w_{(P, A_1, A_2, F)}\rangle + |w_{(P, A_2, A_1, F)}\rangle \\ &= |V_{\emptyset, \emptyset}^{\rightarrow A_1}\rangle * |V_{\emptyset, A_1}^{\rightarrow A_2}\rangle * |V_{\{A_1\}, A_2}^{\rightarrow F}\rangle + |V_{\emptyset, \emptyset}^{\rightarrow A_2}\rangle * |V_{\emptyset, A_2}^{\rightarrow A_1}\rangle * |V_{\{A_2\}, A_1}^{\rightarrow F}\rangle \\ &= |1\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_1^I} |\mathbb{1}\rangle^{A_1^O A_2^I} |\mathbb{1}\rangle^{A_2^O F_t} |1\rangle^{F_c} + |2\rangle^{P_c} |\mathbb{1}\rangle^{P_t A_2^I} |\mathbb{1}\rangle^{A_2^O A_1^I} |\mathbb{1}\rangle^{A_1^O F_t} |2\rangle^{F_c} \end{aligned}$$

Standard process matrix of switch recovered as $W_{\text{switch}} = |w_s\rangle\langle w_s|$

$$\text{Tr}_F W_{\text{switch}} = \underbrace{|1\rangle\langle 1|^{P_c} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{P_t A_1^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_1^O A_2^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_2^O}}_{W^{A_1 \prec A_2}} + \underbrace{|2\rangle\langle 2|^{P_c} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{P_t A_2^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_2^O A_1^I} \otimes |\mathbb{1}\rangle\langle \mathbb{1}|^{A_1^O}}_{W^{A_2 \prec A_1}}$$

Beyond the Quantum Switch?

- N -partite generalisation of the quantum switch is also a QC-QC
- Do QC-QCs offer something new, or are they all “equivalent” to the switch?
- Need a better understanding of causally nonseparable resources and free operations
 - Taddei, Nery and Aolita [arXiv:1903.06180]: local operations and controlled non-signalling operations of bipartite processes
 - Composition: Possible compositions severely restricted [Guérin et al., NJP 2019], but can, e.g., concatenate switches, or insert them inside other switches

Can find QC-QCs such that $\text{Tr}_F W$ not a mixture of valid process compatible with fixed last parties, i.e.

$$\text{Tr}_F W \neq q_1 W^{\{A_1, A_2\} \prec A_3} + q_2 W^{\{A_1, A_3\} \prec A_2} + q_3 W^{\{A_2, A_3\} \prec A_1}$$

- Qualitatively different to the switch

Beyond the Quantum Switch?

- N -partite generalisation of the quantum switch is also a QC-QC
- Do QC-QCs offer something new, or are they all “equivalent” to the switch?
- Need a better understanding of causally nonseparable resources and free operations
 - Taddei, Nery and Aolita [arXiv:1903.06180]: local operations and controlled non-signalling operations of bipartite processes
 - Composition: Possible compositions severely restricted [Guérin et al., NJP 2019], but can, e.g., concatenate switches, or insert them inside other switches

Can find QC-QCs such that $\text{Tr}_F W$ not a mixture of valid process compatible with fixed last parties, i.e.

$$\text{Tr}_F W \neq q_1 W^{\{A_1, A_2\} \prec A_3} + q_2 W^{\{A_1, A_3\} \prec A_2} + q_3 W^{\{A_2, A_3\} \prec A_1}$$

- Qualitatively different to the switch

Summary & Outlook

- Quantum mechanics allows us to process information in a causally indefinite way
 - Quantum switch
 - Process matrix formalism
- Quantum circuits with classical control of causal order
 - General realisation of causally separable processes
- Quantum circuits with quantum control of causal order
 - Potential new realisable, causally nonseparable, circuits beyond the quantum switch?
 - Do QC-QCs provide new information theoretical advantages?
 - Are there other classes of physically realisable processes?

Thank you!
[paper coming soon]

Choi Isomorphism and Link Product

- $|\mathbb{1}\rangle\rangle = \sum_i |i\rangle \otimes |i\rangle$ is the “pure Choi isomorphism” of an identity channel
- Pure Choi isomorphism: for an operator A , $|A\rangle\rangle = \mathbb{1} \otimes A |\mathbb{1}\rangle\rangle$
- Mixed Choi isomorphism: for a CP map \mathcal{M} , $M = \mathcal{I} \otimes \mathcal{M}(|\mathbb{1}\rangle\rangle\langle\langle\mathbb{1}|)$
- Inverse Choi isomorphism given by the link product: $\mathcal{M}(\rho) = M * \rho$;
 $A |\psi\rangle = |A\rangle\rangle * |\psi\rangle$

Constraints for Process Matrix Validity

Recall the notation:

$${}_X W := (\text{Tr}_X W) \otimes \frac{\mathbb{1}^X}{d_X}, \quad {}_1 W := W, \quad [\sum_X \alpha_X {}_X] W := \sum_X \alpha_X {}_X W,$$

Space of valid process matrices


$$W \in \mathcal{L}^{\mathcal{N}} \Leftrightarrow \forall \chi \subseteq \mathcal{N}, \chi \neq \emptyset, \Pi_{i \in \chi} [1 - A_O^i] A_{IO}^{\mathcal{N} \setminus \chi} W = 0,$$

Space of valid process compatible with A first

$$W \in \mathcal{L}^{A_k \prec (\mathcal{N} \setminus A_k)}$$

$$\Leftrightarrow [1 - A_O^k] A_{IO}^{\mathcal{N} \setminus k} W = 0 \quad \text{and} \quad \forall \chi \subseteq \mathcal{N} \setminus k, \chi \neq \emptyset, \Pi_{i \in \chi} [1 - A_O^i] A_{IO}^{\mathcal{N} \setminus k \setminus \chi} W = 0,$$

Characterisation of QC-FOs



$$W = M_1 * M_2 * \cdots * M_{N+1} = \text{Tr}_{\alpha_1 \dots \alpha_N} [M_1 \cdot M_2^{T_{\alpha_1}} \cdots M_{N+1}^{T_{\alpha_N}}]$$

The constraint that the M_n are CPTP maps and thus satisfy

$\text{Tr}_{A_{n+1}^I \alpha_{n+1}} M_{n+1} = \mathbb{1}^{A_n^O \alpha_n}$ allow the W of QC-FOs to be characterised

QC-FOs compatible with order $P \prec A_1 \prec \cdots \prec A_N \prec F$


$$\text{Tr}_F W = W_{(N)} \otimes \mathbb{1}^{A_N^O},$$

$$\text{Tr}_{A_{n+1}^I} W_{(n+1)} = W_{(n)} \otimes \mathbb{1}^{A_n^O} \quad \forall n = 1, \dots, N-1,$$

$$\text{and } \text{Tr}_{A_1^I} W_{(1)} = \mathbb{1}^P.$$

where $W_{(n)} := \frac{1}{d_n^O d_{n+1}^O \cdots d_N^O} \text{Tr}_{A_n^O A_{\{n+1, \dots, N\}}^I} W$ are reduced process matrices

QC-CC Characterisation



Sufficient Condition for N -partite Causal Separability


$$W^{\text{sep}} = \sum_{\pi \in \Pi} \widetilde{W}_{\pi}, \quad \text{with}$$

1. $\widetilde{W}_{\pi} \geq 0$ for each permutation π of $(1, \dots, N) \equiv (A_1, \dots, A_N)$
2. For every ordered subset (k_1, \dots, k_n) (with $1 \leq n \leq N$),

$$\widetilde{W}_{(k_1, \dots, k_n)} := \sum_{\pi \in \Pi_{(k_1, \dots, k_n)}} \widetilde{W}_{\pi}, \quad \text{satisfies}$$

$$\text{Tr}_{A_{N \setminus \{k_1, \dots, k_n\}}^{IO}} \widetilde{W}_{(k_1, \dots, k_n)} = (\text{Tr}_{A_{k_n}^O} [\text{Tr}_{A_{N \setminus \{k_1, \dots, k_n\}}^{IO}} \widetilde{W}_{(k_1, \dots, k_n)}]) \otimes \mathbb{1}^{A_{k_n}^O}$$

QC-CC Characterisation



Sufficient Condition for Fourpartite Causal Separability


Valid process compatible with A first (up to norm.)

$$\begin{aligned}
 W &= \underbrace{W_{(A)}}_{\text{Valid process compatible with A first (up to norm.)}} + \underbrace{W_{(B)}}_{\text{...}} + \underbrace{W_{(C)}}_{\text{...}} + \underbrace{W_{(D)}}_{\text{...}} \\
 &= \underbrace{\widetilde{W}_{(A,B)}}_{\text{...}} + \underbrace{\widetilde{W}_{(A,C)}}_{\text{...}} + \underbrace{\widetilde{W}_{(A,D)}}_{\text{...}} + \underbrace{\dots}_{\text{...}} + \underbrace{\dots}_{\text{...}} + \underbrace{\dots}_{\text{...}} \\
 &= \underbrace{\widetilde{W}_{(A,B,C,D)} + \widetilde{W}_{(A,B,D,C)}}_{\text{...}} + \underbrace{\widetilde{W}_{(A,C,B,D)} + \widetilde{W}_{(A,C,D,B)}}_{\text{...}} + \underbrace{\widetilde{W}_{(A,D,B,C)} + \widetilde{W}_{(A,D,C,B)}}_{\text{...}} + \underbrace{\dots}_{\text{...}} + \underbrace{\dots}_{\text{...}} + \underbrace{\dots}_{\text{...}}
 \end{aligned}$$

For any CP map M_A , $(W_{(A,B)})|_{M_A}$ is valid, compatible with B first

For any CP maps M_A, M_B , $(W_{(A,B,C,D)})|_{M_A \otimes M_B}$ is valid, compatible with C first

QC-CC Characterisation



Sufficient Condition for N -partite Causal Separability


$$W^{\text{sep}} = \sum_{\pi \in \Pi} \widetilde{W}_{\pi}, \quad \text{with}$$

1. $\widetilde{W}_{\pi} \geq 0$ for each permutation π of $(1, \dots, N) \equiv (A_1, \dots, A_N)$
2. For every ordered subset (k_1, \dots, k_n) (with $1 \leq n \leq N$),

$$\widetilde{W}_{(k_1, \dots, k_n)} := \text{Tr}_{A_{N \setminus \{k_1, \dots, k_n\}}^{IO}} \sum_{\pi \in \Pi_{(k_1, \dots, k_n)}} \widetilde{W}_{\pi}, \text{ satisfies}$$

$$\widetilde{W}_{(k_1, \dots, k_n)} = (\text{Tr}_{A_{k_n}^O} \widetilde{W}_{(k_1, \dots, k_n)}) \otimes \mathbb{1}^{A_{k_n}^O}$$

QC-QC Characterisation



- As for QC-CCs, can characterise such W with SDP constraints

Characterisation of circuits with quantum control of causal order


W is the process matrix of a QC-QC iff \exists PSD matrices $W_{(\mathcal{K},\ell)} \in PA_{\mathcal{K}}^{IO} A_{\ell}^I$
 $\forall \mathcal{K} \subsetneq \mathcal{N}, \ell \in \mathcal{N} \setminus \mathcal{K}$ satisfying

$$\text{Tr}_F W = \sum_{k \in \mathcal{N}} \widetilde{W}_{(\mathcal{N} \setminus \{k\}, k)} \otimes \mathbb{1}^{A_k^O},$$

$$\forall \emptyset \subsetneq \mathcal{K} \subsetneq \mathcal{N}, \sum_{\ell \in \mathcal{N} \setminus \mathcal{K}} \text{Tr}_{A_{\ell}^I} \widetilde{W}_{(\mathcal{K}, \ell)} = \sum_{k \in \mathcal{K}} \widetilde{W}_{(\mathcal{K} \setminus \{k\}, k)} \otimes \mathbb{1}^{A_k^O},$$

$$\text{and } \sum_{\ell \in \mathcal{N}} \text{Tr}_{A_{\ell}^I} W_{(\emptyset, \ell)} = \mathbb{1}^P.$$

QC-QC Characterisation



- As for QC-CCs, can characterise such W with SDP constraints

Three-operation QC-QC Characterisation

- $\text{Tr}_F W = \widetilde{W}_{(\{B,C\},A)} \otimes \mathbb{1}^{A^O} + \widetilde{W}_{(\{A,C\},B)} \otimes \mathbb{1}^{B^O} + \widetilde{W}_{(\{A,B\},C)} \otimes \mathbb{1}^{C^O}$
- $\text{Tr}_{CI} \widetilde{W}_{(\{A,B\},C)} = \widetilde{W}_{(\{A\},B)} \otimes \mathbb{1}^{B^O} + \widetilde{W}_{(\{B\},A)} \otimes \mathbb{1}^{A^O}$, etc.
- $\text{Tr}_{BI} \widetilde{W}_{(\{A\},B)} + \text{Tr}_{CI} \widetilde{W}_{(\{A\},C)} = W_{(\{\emptyset\},A)} \otimes \mathbb{1}^{A^O}$, etc.
- $\text{Tr}_{AI} W_{(\{\emptyset\},A)} + \text{Tr}_{BI} W_{(\{\emptyset\},B)} + \text{Tr}_{CI} W_{(\{\emptyset\},C)} = \mathbb{1}^P$

QC-QCs and Causal Correlations

Can quantum circuits with quantum control of causal order violate causal inequalities?

QC-QC correlations are causal

Let W be a QC-QC with trivial spaces \mathcal{H}^P and \mathcal{H}^F . Then the correlations

$$P(a_1, \dots, a_N | x_1, \dots, x_N) = \text{Tr}[W \cdot (M_{a_1|x_1}^T \otimes \dots \otimes M_{a_N|x_N}^T)]$$

are causal for any instruments $\{M_{a_i|x_i}\}_{x_i}$.

- Can noncausal correlations be realised in nature?
 - Would require going beyond this type of generic, coherently controlled circuit
- QC-QCs nevertheless have potential for new advantages arising from indefinite causal order
 - New classes of physically realisable, causally nonseparable processes
 - Use as “quantum super-instruments”, generalising quantum testers, for transformation tasks

QC-QCs and Causal Correlations

Can quantum circuits with quantum control of causal order violate causal inequalities?

QC-QC correlations are causal

Let W be a QC-QC with trivial spaces \mathcal{H}^P and \mathcal{H}^F . Then the correlations

$$P(a_1, \dots, a_N | x_1, \dots, x_N) = \text{Tr}[W \cdot (M_{a_1|x_1}^T \otimes \dots \otimes M_{a_N|x_N}^T)]$$

are causal for any instruments $\{M_{a_i|x_i}\}_{x_i}$.

- Can noncausal correlations be realised in nature?
 - Would require going beyond this type of generic, coherently controlled circuit
- QC-QCs nevertheless have potential for new advantages arising from indefinite causal order
 - New classes of physically realisable, causally nonseparable processes
 - Use as “quantum super-instruments”, generalising quantum testers, for transformation tasks

QC-QCs and Causal Correlations

Can quantum circuits with quantum control of causal order violate causal inequalities?

QC-QC correlations are causal

Let W be a QC-QC with trivial spaces \mathcal{H}^P and \mathcal{H}^F . Then the correlations

$$P(a_1, \dots, a_N | x_1, \dots, x_N) = \text{Tr}[W \cdot (M_{a_1|x_1}^T \otimes \dots \otimes M_{a_N|x_N}^T)]$$

are causal for any instruments $\{M_{a_i|x_i}\}_{x_i}$.


- Can noncausal correlations be realised in nature?
 - Would require going beyond this type of generic, coherently controlled circuit
- QC-QCs nevertheless have potential for new advantages arising from indefinite causal order
 - New classes of physically realisable, causally nonseparable processes
 - Use as “quantum super-instruments”, generalising quantum testers, for transformation tasks

Example: New type of 3-operation QC-QC


$N = 3$, two qubit “targets” and 3-dimensional “control”

- Initial control state $|k_1\rangle^{P_c}$ determines first party
- First party acts on first target qubit $|\psi\rangle^{P_{t1}}$, while $|\phi\rangle^{P_{t2}}$ encoded as ancilla in polarisation
- Unitary U_i applied jointly on both “targets”: output of A_{k_1} may determine (dynamically, coherently) k_2

Can be represented in an “unravelled” form as:



Example: New type of 3-operation QC-QC



Controlled operations can be written

$$|V_{\emptyset, \emptyset}^{\rightarrow k_1}\rangle\rangle = |k_1\rangle^{P_c} |\mathbb{1}\rangle\rangle^{P_{t_1} A_{k_1}^I} |\mathbb{1}\rangle\rangle^{P_{t_2} \alpha_1}$$


$$|V_{\emptyset, k_1}^{\rightarrow k_2}\rangle\rangle = (\langle k_2 - k_1 - 1 |^{\alpha_2} |U_{k_1}\rangle\rangle^{A_{k_1}^O \alpha_1 A_{k_2}^I \alpha_2}) \otimes |k_2 - k_1 - 1\rangle^{\alpha_2}$$

$$|V_{\{k_1\}, k_2}^{\rightarrow k_3}\rangle\rangle = |U_{k_3+3}\rangle\rangle^{A_{k_2}^O \alpha_2 A_{k_3}^I \alpha_3}$$

$$|V_{\{k_1, k_2\}, k_3}^{\rightarrow F}\rangle\rangle = |k_3\rangle^{F_c} |\mathbb{1}\rangle\rangle^{A_{k_3}^O F_{t_2}} |\mathbb{1}\rangle\rangle^{\alpha_3 F_{t_1}}$$

$$\begin{aligned} |w_{(P, A_1, A_2, A_3, F)}\rangle\rangle &= |\mathbb{1}\rangle^{P_c} |\mathbb{1}\rangle\rangle^{P_{t_1} A_1^I} (\langle k_2 - k_1 - 1 |^{\alpha_2} |U_{k_1}\rangle\rangle^{A_{k_1}^O P_{t_2} A_{k_2}^I \alpha_2}) \\ &\otimes (\langle k_2 - k_1 - 1 |^{\alpha_2} |U_{k_3+3}\rangle\rangle^{A_{k_2}^O \alpha_2 A_{k_3}^I F_{t_2}}) |\mathbb{1}\rangle\rangle^{A_3^O F_{t_1}} |3\rangle^{F_c} \end{aligned}$$


Example: New type of 3-operation QC-QC



As before, define $W = |w\rangle\langle w|$ with $|w\rangle = \sum_{k_1, k_2, k_3} |w_{(P, k_1, k_2, k_3, F)}\rangle$

- W is easily seen to be causally nonseparable
 - Pure process, not compatible with any of A , B or C being first (after P)
- Appears qualitatively different to the quantum switch
 - Two target qubits, one of which is also used to control the causal order
- But how to prove W is fundamentally inequivalent to quantum switch?
 - Could imagine composing switches, using control of one as target for another, etc.
 - Need a more complete resource theory, e.g. generalising Taddei, Nery and

Example: New type of 3-operation QC-QC



As before, define $W = |w\rangle\langle w|$ with $|w\rangle = \sum_{k_1, k_2, k_3} |w_{(P, k_1, k_2, k_3, F)}\rangle$

- W is easily seen to be causally nonseparable
 - Pure process, not compatible with any of A , B or C being first (after P)
- Appears qualitatively different to the quantum switch
 - Two target qubits, one of which is also used to control the causal order
- But how to prove W is fundamentally inequivalent to quantum switch?
 - Could imagine composing switches, using control of one as target for another, etc.
 - Need a more complete resource theory, e.g. generalising Taddei, Nery and